- 14 Andriole GL, Kirby R. Safety and tolerability of the dual 5 α -reductase inhibitor dutasteride in the treatment of benign prostatic hyperplasia. *Eur Urol* 2003;44:82-8.
- 15 Speakman MJ, Kirby RS, Joyce A, Abrams P, Pocock R. Guideline for the primary care management of male lower urinary tract symptoms. *BJU Int* 2004;93:985-90.
- 16 Edwards JE, Moore RA. Finasteride in the treatment of clinical benign prostatic hyperplasia: a systematic review of randomised trials. *BMC Urol* 2002;2:14.
- 17 Roehrborn CG, Marks LS, Fenter T, Freedman S, Tuttle J, Gittleman M, et al. Efficacy and safety of dutasteride in the four-year treatment of men with benign prostatic hyperplasia. *Urology* 2004;63:709-15.
- 18 GlaxoSmithKline trial registration website. Study No: ARI40001, year 1. Last updated 22 December 2004. <http://ctr.gsk.co.uk/Summary/dutasteride/studylist.asp> (last accessed 18 Jul 2006).
- 19 Kaplan S, Garvin D, Gilhooly P, Koppel M, Labasky R, Milsten R, et al. Impact of baseline symptom severity on future risk of benign prostatic hyperplasia-related outcomes and long-term response to finasteride. The Pless Study Group. *Urology* 2000;56:610-6.
- 20 McConnell JD, Roehrborn CG, Bautista OM, Andriole GL Jr, Dixon CM, Kusek JW, et al. The long-term effect of doxazosin, finasteride, and combination therapy on the clinical progression of benign prostatic hyperplasia. *N Engl J Med* 2003;349:2387-98.
- 21 Chapple CR. Pharmacological therapy of benign prostatic hyperplasia/lower urinary tract symptoms: an overview for the practising clinician. *BJU Int* 2004;94:738-44.
- 22 National Institute for Clinical Excellence. *Clinical guidelines and referral protocols—referral practice: a guide to appropriate referral from general to specialist services*. London: NICE, May 2000.

(Accepted 3 August 2006)

doi 10.1136/bmj.38946.616551.BE1

Lesson of the week

Acute renal failure induced by contrast medium: steps towards prevention

R Mathew, K Haque, W Woothipoom

Acute renal failure is a well known complication of procedures that involve iodinated contrast media.¹ Despite this, contrast medium induced nephropathy accounts for about 12% of all cases of hospital acquired renal failure.² Prevention of this type of nephropathy is crucial as it is associated with prolonged hospital stay, risk of permanent renal impairment, and a more than fivefold increase in mortality.^{3,4} We report a case of acute renal failure in a woman with chronic renal disease who was investigated for metastatic breast cancer with contrast enhanced computed tomography. This case shows the importance of carrying out a risk assessment for contrast medium induced nephropathy before using procedures that involve iodinated contrast media.

Case report

An 81 year old woman with type II diabetes was admitted after a hypoglycaemic episode. She had a four week history of non-productive cough, which failed to resolve with antibiotics. She had been treated with surgery and radiotherapy for breast cancer 13 years previously. She also had stage 3 kidney disease and hypertension.

Breast examination was normal but there were signs of a left sided pleural effusion. Chest x ray showed a possible left sided coin lesion and pleural effusion. Her white blood cell count was 5.6×10^9 /litre, C reactive protein was 15 mg/litre, and the pleural aspirate was an exudate containing 38 g/litre protein and 0.5×10^9 /litre white blood cells. Plasma creatinine was 224 $\mu\text{mol/litre}$, which was not very different from her preadmission value. In view of the history of breast cancer, contrast enhanced computed tomography of the chest and abdomen was performed. No abnormalities were detected in the lungs, but a lesion, suggestive of metastasis, was detected in the left adrenal.

Serum creatinine values during admission in a woman with contrast medium induced acute renal failure

Over the next two days, the patient became increasingly nauseous and anorexic. Creatinine rose to 392 $\mu\text{mol/litre}$ (figure). Despite supportive treatment for acute renal failure she became anuric, her blood pressure began to drop, and creatinine rose to 664 $\mu\text{mol/litre}$. She was transferred to the intensive care unit, where haemofiltration was started. Her renal function started to recover over the next few days and her condition began to improve.

Discussion

Nephropathy induced by contrast medium is defined as an impairment in renal function that occurs within 72 hours of giving contrast medium.⁵ This impairment is characterised by an increase in serum creatinine of at least 44 $\mu\text{mol/litre}$ or 25% above the baseline. Creatinine typically peaks three to five days after contrast administration and returns to baseline values within two weeks. Renal replacement therapy is needed in a minority of patients, and in-hospital mortality may be as high as 62% in these cases.⁴ No specific treatment is available for contrast medium induced nephropathy and management is supportive.

Risk assessment for renal failure must be performed before procedures involving the administration of contrast media

University College
London, London
WC1E 6BT
R Mathew
medical student

North Middlesex
University Hospital,
London N18 1QX
K Haque
specialist registrar in
care of the elderly
medicine

Care of the Elderly
Medicine, North
Middlesex
University Hospital
W Woothipoom
consultant physician

Correspondence to:
R Mathew
rgmathew@
hotmail.com

BMJ 2006;333:539-40

Risk factors for contrast medium induced nephropathy

- Pre-existing renal insufficiency
- Diabetes mellitus
- Age >75 years
- Concurrent use of nephrotoxic drugs (non-steroidal anti-inflammatory drugs, aminoglycosides)
- Dehydration
- Hypotension
- Heart failure
- Cirrhosis
- Nephrotic syndrome
- Hypertension

The box shows the key risk factors for developing contrast medium induced nephropathy.⁶ Underlying impairment of renal function, particularly if secondary to diabetes, is the most important risk factor, and the associated incidence of contrast nephropathy is 12-27%. This compares with an incidence of 0-5% in patients with normal renal function.⁵ Serum creatinine should ideally be measured in all patients before giving contrast medium, but particularly if a history of renal impairment or diabetes is present. Serum creatinine, however, is a poor indicator of renal function. The best measure of renal function—and therefore risk of contrast nephropathy—is the glomerular filtration rate. The rate can be estimated from serum creatinine using the Cockcroft-Gault formula or the modification of diet in renal disease formula.⁷ It is especially useful to estimate the glomerular filtration rate in patients with borderline serum creatinine—a value of less than 60 ml/min/1.73 m² predicts increased risk of contrast nephropathy.⁸

If risk factors are present then an alternative imaging technique should be used. If however, administration of contrast medium is deemed necessary, then several steps can be taken to reduce risk. The most important of these is adequate hydration. The most effective regimen has not been defined, but one study showed that intravenous hydration with normal saline at a rate of 1 ml/kg/hour for 24 hours, beginning 12 hours before administration of the contrast medium, was associated with a significantly lower incidence of contrast nephropathy (3.7%) compared with unrestricted oral hydration (34.6%).⁹ Isotonic sodium bicarbonate has been shown to be superior to normal saline in preventing contrast nephropathy, but further evidence is needed to support this intervention.¹⁰ Nephrotoxic drugs increase the risk of developing contrast nephropathy, and it is recommended that these drugs be stopped at least 24 hours before investigation.¹¹ In addition, use of the minimum volume of low osmolar or iso-osmolar contrast media is associated with a lower incidence of contrast nephropathy compared with high osmolar media.¹²⁻¹⁴ Oral N-acetylcysteine has been used as a prophylactic treatment for contrast nephropathy, but there is currently insufficient evidence to advocate its use.^{15 16} Daily monitoring of renal function after contrast administration in high risk patients enables early diagnosis and treatment of renal failure.

We did not perform risk assessment for the development of contrast nephropathy in our patient. We did not provide the relevant clinical details when we ordered computed tomography and the necessary precautions were not taken. In retrospect, this was a serious oversight. Contrast nephropathy is highly underestimated. A retrospective study of 89 patients undergoing contrast enhanced computed tomography showed that none had creatinine concentrations documented on the request card. Consequently, steps to reduce risk were taken in only two of 10 patients at risk.¹⁷

This case emphasises the role of the clinician in preventing contrast medium induced nephropathy. A risk assessment must be performed and details of this should be provided to the radiology department. If it is deemed necessary to give contrast medium to patients with risk factors, measures must be taken to prevent renal failure from developing.

Contributors: RM and WW had the original idea. RM wrote the paper. RM and KH searched the literature. KH and WW cared for the patient. WW revised the final manuscript and is guarantor.

Funding: None.

Competing interests: None declared.

- 1 Weisbord SD, Palevsky PM. Radiocontrast-induced acute renal failure. *J Intensive Care Med* 2005;20:63-75.
- 2 Hou SH, Bushinsky DA, Wish JB, Cohen JJ, Harrington JT. Hospital-acquired renal insufficiency: a prospective study. *Am J Med* 1983;74:243-8.
- 3 Cox CD, Tsikouris JP. Preventing contrast nephropathy: what is the best strategy? A review of the literature. *J Clin Pharmacol* 2004;44:327-37.
- 4 Levy EM, Viscoli CM, Horwitz RI. The effect of acute renal failure on mortality. A cohort analysis. *JAMA* 1996;275:1489-94.
- 5 Thomsen HS, Morcos SK. Contrast media and the kidney: European Society of Urogenital Radiology (ESUR) guidelines. *Br J Radiol* 2003;76:513-8.
- 6 Barrett BJ, Parfrey PS. Clinical practice. Preventing nephropathy induced by contrast medium. *N Engl J Med* 2006;354:379-86.
- 7 Levey AS, Bosch JP, Lewis JB, Greene T, Rogers N, Roth D. A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of Diet in Renal Disease Study Group. *Ann Intern Med* 1999;130:461-70.
- 8 Mehran R, Aymong ED, Nikolsky E, Lasic Z, Iakovou I, Fahy M, et al. A simple risk score for prediction of contrast-induced nephropathy after percutaneous coronary intervention: development and initial validation. *J Am Coll Cardiol* 2004;44:1393-9.
- 9 Trivedi HS, Moore H, Nasr S, Aggarwal K, Agrawal A, Goel P, et al. A randomized prospective trial to assess the role of saline hydration on the development of contrast nephrotoxicity. *Nephron Clin Pract* 2003;93:C29-34.
- 10 Merten GJ, Burgess WP, Gray LV, Holleman JH, Roush TS, Kowalchuk CJ, et al. Prevention of contrast-induced nephropathy with sodium bicarbonate: a randomized controlled trial. *JAMA* 2004;291:2328-34.
- 11 Morcos SK. Prevention of contrast media nephrotoxicity—the story so far. *Clin Radiol* 2004;59:381-9.
- 12 Barrett BJ, Carlisle EJ. Metaanalysis of the relative nephrotoxicity of high- and low-osmolality iodinated contrast media. *Radiology* 1993;188:171-8.
- 13 Aspelin P, Aubry P, Fransson SG, Strasser R, Willenbrock R, Berg KJ, et al. Nephrotoxic effects in high-risk patients undergoing angiography. *N Engl J Med* 2003;348:491-9.
- 14 Freeman RV, O'Donnell M, Share D, Meenges WL, Kline-Rogers E, Clark VL, et al. Nephropathy requiring dialysis after percutaneous coronary intervention and the critical role of an adjusted contrast dose. *Am J Cardiol* 2002;90:1068-73.
- 15 Royal College of Radiologists. *Standards for iodinated intravascular contrast agent administration to adult patients*. London: RCR, 2005. Ref BFCR(05)7. www.rcr.ac.uk/docs/radiology/pdf/IVcontrastPrintFinal.pdf (last accessed 8 Aug 2006).
- 16 Zagler A, Azadpour M, Mercado C, Hennekens CH. N-acetylcysteine and contrast-induced nephropathy: a meta-analysis of 13 randomized trials. *Am Heart J* 2006;151:140-5.
- 17 Whittaker C, Duerden R, Rose C. *Contrast induced nephropathy in patients undergoing body CT*. London: Royal College of Radiologists. www.rcr.ac.uk/index.asp?PageID=547&AbID=786 (last accessed 28 Jun 2006).

(Accepted 21 July 2006)

doi 10.1136/bmj.38943.401852.80