In brief

Conviction for assisted suicide upheld: A Dutch appeal court has upheld the conviction of the founder of suicide support group De Einder for helping a 25 year old woman to kill herself. Jan Hilarius, aged 74, offered written instructions on suicide and medication. His sentence was reduced from 12 to four months because of his age and previous unblemished record. Dutch euthanasia laws allow only doctors to assist suicide.

Poverty in US increases: The poverty rate in US residents increased from 11.3% in 2000 to 12.7% in 2004. The population that earned less than half the poverty threshold for a family of four—\$19 307 (£10 135; €15 003)—grew by 20%. The study is to be published in October's *American Journal of Preventive Medicine* (www.ajpm-online.net, doi: 10.1016/j.amepre.2006.06.022).

Skin cancer treated with gene therapy: Two men with advanced melanoma achieved full remission for 18 months as a result of being treated with their own genetically modified, "strengthened" T cells, researchers at the US National Cancer Institute have found (*Science* 2006 Aug 31 doi: 10.1126/science.1129003). Sustained remission was not achieved in 15 other similarly treated patients.

COPD treatment has sex bias: In patients treated with long term oxygen therapy for chronic obstructive pulmonary disease (COPD), women are more likely to die of their disease than men (*American Journal of Respiratory and Critical Care Medicine* 2006;174:488-9). The study of 435 oxygen dependent patients, monitored for seven years, found that the risk of death was 54% higher in women.

NHS opts for private supplier: The NHS in England has chosen the German logistics company DHL to supply it with healthcare related goods and equipment over the next 10 years. The deal will save the NHS £1bn over a decade by increasing the volume of supplies handled by one company, said health minister Andy Burnham.

Calcium channel blockers associated with less diabetes

Susan Mayor *Barcelona*

Patients taking a treatment for high blood pressure based on a calcium channel blocker had a 34% lower risk of developing diabetes than patients using a β blocker. This result is from a sub-study of the Anglo-Scandinavian cardiac outcomes trial (ASCOT) reported this week at the World Congress of Cardiology meeting in Barcelona.

Treating patients with amlodipine, with or without the angiotensin converting enzyme inhibitor perindopril, was associated with a 34% reduction in new onset diabetes compared with treatment with the β blocker atenolol, with or without the diuretic bendroflumethiazide, the study showed.

ASCOT, the largest study of patients with hypertension yet conducted in Europe, included 19 257 hypertensive patients (160/100 mm Hg or higher

untreated, or 140/90 mm Hg or higher after treatment with one or more drug) with no previous myocardial infarction or current clinical heart disease—and three or more other risk factors for coronary heart disease.

They were randomised to antihypertensive treatment based on atenolol, with the addition of bendroflumethiazide, if required, to reach the target of 140/90 mm Hg in patients without diabetes and 130/80 mm Hg in patients with diabetes, or to amlodipine plus perindopril if the target was not reached with monotherapy.

Results showed that 1366 of the 14 120 patients who did not have diabetes at the start of the study developed the condition over the five year study period—567 (8%) in the amlodipine treated group and 799 (11%) in

patients treated with atenolol ($P=0.0001$). Diabetes was diagnosed based on criteria set by the World Health Organization (fasting glucose greater than 7 mmol/l or non-fasting glucose greater than 11.1 mmol/l). All of the patients in the substudy receiving the newer treatments (a calcium channel blocker with or without an angiotensin converting enzyme inhibitor) had a lower risk of developing diabetes, irrespective of their initial risk factors.

Reporting the findings, Dr Ajay Gupta, from the International Centre for Circulatory Health at Imperial College in London, said, "One of the most important risk factors for developing new-onset diabetes was being allocated to the beta-blocker based strategy. Patients randomised to the more modern blood pressure lowering strategy—amlodipine with or without perindopril—were 34% less likely to develop new onset diabetes. This is important because diabetes significantly increases the risk of myocardial infarction and stroke."

Troops are needed in Darfur, says UN

Peter Moszynski *London*

Hundreds of thousands of people could die in the next few weeks in the Darfur region of Sudan unless United Nations troops are allowed to take over peace keeping in the region to avoid further war, the UN has warned. Further outbreaks of fighting would mean the withdrawal of international staff from Darfur, leaving millions of vulnerable people without help, officials have said.

The warning comes as the World Health Organization's cholera task force told the *BMJ* that the failure to contain a *Vibrio cholerae* outbreak in Sudan meant that there was a serious risk of a regional epidemic, especially given the current extent of rain and flooding in Sudan and neighbouring countries.

The mandate of the African Union Mission in Sudan expires at the end of September, but UN forces will be allowed to take

over only if Sudan agrees. Currently the government in Khartoum insists that the Sudanese people "will not consent to any resolution that will violate its sovereignty."

The UN humanitarian coordinator Jan Egeland warned the UN security council, "Our entire humanitarian operation in Darfur—the only lifeline for more than three million people—is presently at risk. We need immediate action on the political front

to avoid a humanitarian catastrophe with massive loss of life."

Meanwhile, Dr Claire-Lise Chaignat, the head of the WHO's cholera task force, warned of a *V cholerae* outbreak in Sudan. WHO has already recorded almost 25 000 cases and more than 700 deaths this year in Sudan alone, and many of the worst affected areas are already out of reach to health workers.

"All the ingredients of a major epidemic are in place. You have already vulnerable populations, weakened by malnutrition and exposed to the elements, you have displaced people crowded together in unsanitary conditions, you have extensive rainfall and flooding, and you have the presence of the cholera bacillus."

Dr Chaignat explained, "The outbreak has already spread from south Sudan across the country into Darfur and the east and now risks affecting the neighbouring countries. It appears to have already crossed into Ethiopia and could soon emerge in Chad. This disease doesn't respect borders."

She said that such outbreaks were easily preventable.

Refugees at waterhole in the dry river Wadi Kaca in Darfur