Israel permits ova donation from woman who is not being treated

A lesbian woman in her 30s who, for medical reasons, may not be able to become pregnant has been permitted by Israel's health ministry to donate ova to her partner. After an ovum is fertilised with sperm from an anonymous donor the partner will carry the pregnancy.

Such arrangements will automatically become permissible if a bill sponsored by the ministry is passed by the Knesset (parliament). A legal adviser to the ministry, Mira Huebner, said she hoped the bill, which has been five years in the making, will be approved during the Knesset's upcoming autumn session.

The bill sets down strict rules for the donation of ova by women who are not undergoing fertility treatments. If passed, it will be one of the most comprehensive laws on ova donation in the world.

For decades Israeli law has specified that only women who are undergoing fertility treatment can donate ova, with unpaid donors doing so for altruistic reasons.

Judy Siegel-Itzkovich *Jerusalem*

Drug eluting stents cause more deaths than plain stents

The safety of drug eluting stents has been called into question by two meta-analyses of their long term effects. The results of the studies were presented at the World Congress of Cardiology in Barcelona last week.

Both studies showed that the risk of death and of Q wave myocardial infarction, a clinical surrogate of stent thrombosis, was higher with drug eluting stents than with bare metal or uncoated stents.

The use by cardiologists of the first generation of drug eluting stents—the sirolimus (Cypher) and the paclitaxel (Taxus) stents—has risen rapidly since they were introduced. Currently around six million people

worldwide have had a drug eluting stent fitted.

Their widespread uptake came after early results showed that the rate of re-stenosis was zero at six months after implantation of drug eluting stents but 26% after implantation of bare metal stents (*New England Journal of Medicine* 2002;346: 1773-80).

However, four years later case reports of late stent thrombosis and delayed healing in patients fitted with drug eluting stents prompted two Swiss researchers to review the evidence and conduct meta-analyses.

Lisa Hitchen *London*

Reports from the World Congress of Cardiology are at www.worldcardio2006.org.

Bird flu's deadly effects may give clues to treatment

The massive inflammation caused by the H5N1 strain of the avian influenza virus explains why the virus is so deadly, but it also provides clues to better treatments, experts say.

In a letter published online ahead of print publication on 10 September in *Nature Medicine* (www.nature.com/nm, doi: 10.1038/nm1477) the authors document how the immune systems of 18 Vietnamese people who contracted the infection in 2004/5 became stoked up to an extraordinary degree by the virus.

Among the striking observations was a massive release of inflammatory cytokines, causing immune cells to proliferate and provoking inflammation. Some of these cytokines were present at amounts hundreds or thousands of times higher than in patients with ordinary flu.

These observations explain how the infection typically kills patients through pneumonia and multiorgan failure, the researchers say.

In theory, using drugs to dampen down patients' immune systems might help, they say. But so far such methods had not proved particularly effective in practice.

Michael Day *London*

Statin prescribing fell after drug became available over the counter

Roger Dobson *Abergavenny*

The number of GPs' prescriptions of statins fell by 10% in the United Kingdom after low dose simvastatin was made available over the counter in August 2004.

Around 600 000 fewer statin prescriptions were issued in the second quarter of 2005 after the switch than would have been predicted by the previous trend, says a paper that assessed the effect of the policy change.

The authors of the paper, which was published online ahead of print publication on 5 September in *Pharmacoeconomics and Drug Safety* (www3.interscience.wiley.com; doi: 10.1002/pds.1314), say that the drop was not just in prescriptions of low dose statins but in those of higher dose ones too, and they caution that some patients may have been switched to less aggressive treatment. They warn too of a substantial number of people whose use of statins is now not recorded.

"Our results suggest that the new policy allowing OTC [over the counter] sales of 10 mg simvastatin has had a significant impact on the total volume of statin prescriptions in the UK," wrote the researchers, from Montreal's McGill University.

"This shift towards less aggressive statin therapy may reduce the potential health benefits associated with its use. This may have direct clinical implications in this era where 'intensive' statin therapy is increasingly being shown to have additional health benefits.

"The observed decrease in the

rate of statin prescriptions could indicate that there is a non-trivial segment of the population that is receiving treatment that is no longer reflected in these databases. These missing data have important implications for future observational studies of cardiovascular disease."

The authors examined numbers of statin prescriptions in the UK general practice research database from the first quarter of 2001 to the second quarter of 2005. The database has information on around 3.5 million patients.

Their results show that the number of statin prescriptions in the database fell for the first time in the first quarter of 2005, after several years of increase.

"It is a very interesting paper. I suspect the explanation is more complex than the authors suggest, and it may be that a combination of factors is at work," said Nicholas Brooks, president of the British Cardiovascular Society.

"GPs have been writing large numbers of statin prescriptions, and we know from the quality and outcomes framework data that they are meeting their targets for primary and secondary prevention, so sooner or later it was inevitable that we would see a plateau. However, that does not explain the apparent fall. A greater awareness of statin prescribing guidelines, coupled with pressure from primary care trusts to save money, and the change to over the counter status, may all have had an effect."