In brief

FDA fines Red Cross: The US Food and Drug Administration fined the American Red Cross \$4.2m (£2.3m; €3.3m) last week for violating federal rules on blood safety, including failing to ask donors screening questions and to follow testing protocols.

Doctors are encouraged to teach: Doctors should be given more opportunities to undergo teacher training, and medical teaching should be recognised as a core professional activity equivalent to research and care of patients, recommends the BMA's board of medical education in a report published this week, *Doctors as Teachers* (see www.bma.org.uk).

Patients in Burundi are detained for unpaid bills: Hundreds of patients are illegally detained in Burundi's hospitals, sometimes for months, because they can't afford their medical bills, says a report from Human Rights Watch, *A High Price to Pay: The Detention of Poor Patients in Hospitals*. Many are women who unexpectedly needed caesarean deliveries (<http://hrw.org/reports/2006/burundi0906/>).

Dutch committee to oversee mercy killing of newborns: Joep Hubben, professor of health law at Groningen University, has been appointed by the Dutch government to chair a committee to investigate rare cases where doctors "end the lives" of severely ill newborn babies who are suffering hopelessly and unbearably. The move follows years of lobbying by the profession (*BMJ* 2005;330:560).

Emotional difficulties are disproportionate among Aboriginal children: 24% of Australian Aboriginal children aged 4-17 are at high risk of clinically significant emotional or behavioural difficulties, whereas the percentage among non-Aboriginal Australian children is 15%, it was reported at this week's world congress of the International Association for Child and Adolescent Psychiatry and Allied Professions in Melbourne.

Nicotine content in smoke from US cigarettes has increased

Janice Hopkins Tanne *New York*

The amount of nicotine inhaled by cigarette smokers and by bystanders in secondhand smoke has risen by 10% over the past seven years, says a report from the Massachusetts Department of Public Health.

"Although per capita consumption of cigarettes has declined, the amount of nicotine consumed per cigarette has increased. Concurrently, the amount of nicotine present in second-hand smoke has also increased," the report says.

Massachusetts is one of only three US states that require tobacco

companies to submit information every year on the nicotine content of cigarettes sold in the state. The state began requiring the information in 1998. This is the first report of changes in nicotine content from 1998 to 2004.

The information the state received showed major increases in nicotine yield for 92 of the 116 brands of cigarettes manufactured by all three major tobacco companies, Lorillard, Philip Morris, and RJ Reynolds. Yield is the amount of nicotine in the smoke a smoker inhales, not the amount in the cigarette.

"We want healthcare providers to know that smokers are getting more nicotine than in the past and may need additional help in trying to quit," said the Massachusetts health commissioner Paul Cote, Jr.

The department tests for nicotine yields using a method that it considers more accurate than the previous "smoking machine" tests. The department's method "better simulates the smoking behavior of the typical smoker under typical smoking conditions."

"Similar increases were found for each type of cigarette tested (full flavor, light, light/medium, and ultra-light)," the report says. P

Change in Nicotine Yields 1998-2004 is available at <http://mass.gov/dph>.

Experts devise strategy to fight new TB strain

Peter Moszynski *Johannesburg*

Delegates at a conference in Johannesburg last week drew up a seven point action plan to counter the threat of extensively drug resistant tuberculosis (TB), a newly emergent strain that seems to be resistant to nearly all existing drugs.

The new strain was already present on every continent, but a recent outbreak in South Africa was particularly worrying, experts told the meeting, which was hosted by the South African Medical Research Council, with support from the World Health Organization and the US Centers for Disease Control and Prevention. The outbreak in South Africa was of an extraordinarily

A patient with extensively drug resistant tuberculosis in Ethiopia

virulent form and occurs in a region with extremely high HIV co-infection.

WHO said that multidrug resistant strains of the disease are already widespread worldwide—mainly, it says, because of healthcare workers' improper

treatment regimens and failure to ensure that patients complete the whole course of treatment. "Essentially, drug resistance arises in areas with poor TB control programmes," WHO said.

Paul Nunn, of WHO's Stop TB campaign, said that there is now "tremendous international concern" over the "spectre of untreatable TB" and the "extraordinarily high mortality rate demonstrated in the recent outbreak in KwaZulu/Natal."

Sputum samples from 536 people with TB in the town of Tugela Ferry were collected between January 2005 and March 2006. A total of 221 (41%) participants had multidrug resistant TB, of whom 53 had the extensively drug resistant strain. Fifty two of these 53 patients died, within an average of 25 days.

Dr Nunn said the immediate priority was to establish the extent of infection. "We know about the problems in KwaZulu/Natal because the area has comparatively good health facilities. We don't know how serious the disease is in other parts of Africa where there is almost no surveillance."

Karin Weyer, of the South African Medical Research Council, said, "Among the actions required are urgent and rapid surveys in high risk countries to assess the full extent of XDR-TB [extensively drug resistant TB] globally. (See editorial, p 559.) P

See www.who.int/tb/en/ for more information.

Seven point action plan on extensively drug resistant tuberculosis

- Conduct rapid surveys of the strain
- Enhance laboratory capacity
- Improve technical capacity of clinical and public health managers to effectively respond to outbreaks
- Implement infection control precautions
- Increase research support for development of anti-TB drugs
- Increase research support for development of rapid diagnostic tests
- Promote universal access to antiretrovirals under joint TB and HIV activities