REFERENCES AND REVIEWS

SYSTEMIC MANIFESTATIONS DUE TO ALLERGY—W. G. Crook, W. W. Harrison, S. E. Crawford, and B. S. Emerson. *Pediatrics*—Vol. 27:790 (May) 1961.

Allergy was found to cause systemic signs and symptoms in 50 children. Fatigue, pallor, infraorbital circles, nasal congestion, and mental and nervous symptoms were seen in most. Headache, abdominal pain, muscular aching, and a variety of other symptoms occurred in some. Incriminated allergens included foods (especially milk and chocolate) and inhalants. Although two-thirds of the patients had other allergies, such as asthma and hay fever, the systemic manifestations occurred independently and were due usually to different allergens. The criteria for diagnosis included: (1) relief of symptoms when the probable cause was eliminated, (2) precipitation of symptoms by reintroducing the cause, and (3) positive skin tests (inhalants). These cases, and those reported in the literature, suggest that systemic manifestations due to allergy are common, that they can be readily recognized, and that often they can be dramatically relieved by the elimination of a food from the child's diet.

* * *

METHOD FOR STUDY OF ARTERIAL PROFILE—O. J. Pollak and P. S. Morrow. *Arch. Path.*—Vol. 71:576 (May) 1961.

A method was devised to study the profile of coronary arteries, and to evaluate the location and degree of stenosis. Plaster imprints of longitudinally opened arteries were prepared, the casts were segmented with a saw, contours of the segments were traced, the tracings were enlarged with

a pantograph, and the width and height of excursions were measured. The technique does not interfere with chemical or histological studies of the arteries. Application of this method in comparative studies of arteries from persons of varying age, different sex or ethnic origin, and of arteries of subjects who had died of various diseases may help to advance our knowledge of atherosclerosis.

* * *

EFFECT OF PARATHYROIDECTOMY ON INCIDENCE OF RENAL CALCULI—M. G. McGeown. *Lancet*—Vol. 1:586 (March 18) 1961.

There were 68 cases of hyperparathyroidism amongst 405 patients (16.9 per cent) suffering from calcareous renal stones. Fifty-six patients were followed for periods of one to five years, and the effect of parathyroidectomy on stone recurrence was studied. In the postparathyroidectomy period 99 stones occurred in 401 patient-years, but there were only 8 in 158 postparathyroidectomy patient-years ($P. < 0.001$). When each patient was treated as his own control, there was a significant difference between the observed and the expected figures ($P. < 0.001$) for the five postparathyroidectomy years. The decrease in the incidence of stone recurrence justifies the search for hyperparathyroidism in all patients with calcareous calculi.

* * *

DIETARY LIPIDS, THROMBOSIS, AND CLOT-LYSIS—R. F. Scott, K. Alousi, and W. A. Thomas. *Arch. Path.*—Vol. 71:594 (May) 1961.

We have previously demonstrated in rats that diets containing large quantities of common fatty substances, such as butter (plus cholesterol, propylthiouracil and bile salts), will produce thromboses and infarcts, particularly in the

(Continued on Page 60)

when your older patient is out of focus

LIPO-NICIN PTZ

CAPSULES AND ELIXIR T.M.

provides emotional and mental improvement
and a brighter outlook for your aging patients
a safe and efficient cerebral stimulant and vasodilator

LIPO-NICIN PTZ capsules

Each capsule contains:

PTZ (Pentamethylene Tetrazole).....	100 mg.
Nicotinic Acid	100 mg.
Niacinamide	5 mg.
Vitamin C	100 mg.
Thiamine-HCl	25 mg.
Riboflavin	2 mg.
Pyridoxine	3 mg.
1-Glutamic Acid	50 mg.

AVERAGE DOSAGE: One capsule three times daily.

AVAILABLE in bottles of 100 capsules.

CAUTION: Most persons experience a flushing and tingling sensation after taking a higher potency niacin-containing compound. As a secondary reaction some will complain of nausea and other sensations of discomfort. This reaction is transient and is rarely a cause for discontinuance of the drug if the patient is forewarned to expect the reaction.

WARNING: Contraindicated in the presence of epilepsy.

Write for Free Samples and Literature.

THE BROWN PHARMACEUTICAL COMPANY
2500 W. 6th Street, Los Angeles 57, Calif.

LIPO-NICIN PTZ elixir (ELIXIR OF PORT WINE)

Each teaspoonful (5 cc) contains:

PTZ (Pentamethylene Tetrazole).....	100 mg.
Nicotinic Acid	100 mg.
Ascorbic Acid	100 mg.
1-Glutamic Acid	50 mg.
Niacinamide	5 mg.
Riboflavin (as Riboflavin 5'-	
Phosphate Sodium)	2 mg.
D-Pantothenyl Alcohol	5 mg.
In a port wine vehicle. Alcohol 20%	

AVERAGE DOSAGE: One teaspoonful three times daily or as prescribed by physician.

AVAILABLE in Pint bottles.