Capsaicin exhibits neuroprotective effects in a model of transient global cerebral ischemia in Mongolian gerbils

¹Simona Pegorini, ¹Daniela Braidà, ¹Chiara Verzoni, ¹Chiara Guerini-Rocco, ²Gian Giacomo Consalez, ²Laura Croci & ^{*,1}Mariaelvina Sala

¹Department of Pharmacology, Chemotherapy and Medical Toxicology, University of Milan, Via Vanvitelli 32, 20129 Milan, Italy and ²San Raffaele Scientific Institute, Via Olgettina 58, 20132 Milan, Italy

1 Capsaicin, the irritant principle of hot peppers, is a vanilloid agonist known to activate the transient receptor potential channel vanilloid subfamily member 1 (VR1), recently reported to be involved in neurodegeneration. The present study investigated the role of VR1 in a model of global cerebral ischemia in gerbils.

2 Over the dose range tested, capsaicin (0.01, 0.025, 0.05, 0.2 and 0.6 mg kg⁻¹), given 5 min after recirculation, dose-dependently antagonized the ischemia-induced electroencephalographic total spectral power decrease and restored relative frequency band distribution evaluated 7 days after ischemia.

3 Capsaicin, at all tested doses, fully prevented ischemia-induced hyperlocomotion evaluated 1 day after ischemia.

4 Capsaicin dose-dependently antagonized ischemia-induced memory impairment evaluated in a passive avoidance task, 3 days after ischemia.

5 Capsaicin showed a dose-dependent hypothermic effect evaluated for 2 h after recirculation.

6 At 7 days after ischemia, a progressive survival of pyramidal cells in the CA1 subfield in capsaicin-treated gerbils, with a maximum of 80%, at a dose of 0.2 mg kg⁻¹, was obtained.

7 The selective VR1 antagonist, capsazepine (0.01 mg kg⁻¹), reversed capsaicin-induced protective effects, in a competitive manner.

8 These results suggest that the neuroprotective effect of capsaicin may be attributable, at least in part, to VR1 desensitization and provide a valuable target for development of interventional pharmacological strategies.

British Journal of Pharmacology (2005) **144**, 727–735. doi:10.1038/sj.bjp.0706115

Published online 24 January 2005

Keywords: Capsaicin; vanilloid agonist; vanilloid antagonist; ischemia; EEG; memory; motor activity; rectal temperature; CA1; neuroprotection

Abbreviations: ANOVA, analysis of variance; CPZ, capsazepine; EEG, electroencephalogram; TRPV1, transient receptor potential channel vanilloid subfamily member 1; VR1, vanilloid receptor 1

Introduction

Capsaicin (*N*-(3-methoxy-4-hydroxy-benzyl)-8-methyl-6-*trans*-nonenamide or *N*-vanillyl-8-methyl-6-*trans*-nonenamide), the irritant principle of hot peppers, is a vanilloid agonist known to activate the transient receptor potential channel vanilloid subfamily member 1 (TRPV1), named vanilloid receptor 1 (VR1) (Gunthorpe *et al.*, 2002; van der Stelt & Di Marzo, 2004) and to possess analgesic (Perkins & Campbell, 1992), anti-inflammatory (Kim *et al.*, 2003) and hypolocomotor effects (Di Marzo *et al.*, 2001).

TRPV1 is believed to function as a molecular integrator of noxious stimuli (Szallasi & Blumberg, 1999; Cortright & Szallasi, 2004). However, the recent discovery in both human and rat brains of TRPV1, where it is unlikely to be gated by noxious stimuli, has suggested additional functional roles for this channel (Sasamura & Kuraishi, 1999; Hayes *et al.*, 2000;

Mezey *et al.*, 2000) in physiological processes such as satiety, cognition and motor control (Szallasi *et al.*, 1995). There is also some evidence of involvement of TRPV1 in neurodegeneration: (i) Pathological changes in brain temperature or pH, for example after a severe stroke, may influence TRPV1 activity (Marinelli *et al.*, 2002) suggesting an important role in events occurring after ischemia (like modulation of glutamate release and consequent activation of excitatory amino-acid receptors); (ii) TRPV1 is present in brain regions (hippocampus) that are highly susceptible to neurodegenerative insults, suggesting that this ion channel might contribute to the cellular processes involved in neuronal death (Mezey *et al.*, 2000); and (iii) capsaicin and the vanilloid antagonist capsazepine (CPZ), peripherally administered, have been shown to exhibit neuroprotection against ouabain-induced excitotoxicity in rats through a mechanism not fully identified (Veldhuis *et al.*, 2003).

In addition, TRPV1 can be activated by endogenously generated compounds such as anandamide and lipoxigenase products, which accumulate during brain injury (Marinelli

*Author for correspondence; E-mail: mariaelvina.sala@unimi.it
Published online 24 January 2005

et al., 2002). In this context, capsaicin has been shown to stimulate the biosynthesis of endocannabinoids (Di Marzo *et al.*, 2001), which may afford neuroprotection.

In the current study, we investigated the possible neuroprotectant effect of capsaicin *in vivo* using a model of transient global cerebral ischemia in the gerbil. Since capsaicin can be tested only up to 1.0 mg kg^{-1} due to its toxic effects *in vivo* (Di Marzo *et al.*, 2001), we employed a wide range of doses (from 0.01 to 0.6 mg kg^{-1}) below 1 mg kg^{-1} . To quantify the ischemic damage, we measured, from 1 h to 7 days after recirculation, different parameters known to be hardly influenced by global cerebral ischemia: electroencephalographic (EEG) spectral power, spontaneous motor activity, memory function and hippocampal CA1 neuronal count. We measured motor activity on day 1, since ischemia-induced hyperlocomotion is maximal 24 h after occlusion (Araki *et al.*, 1986), memory function on day 3 according to Sala *et al.* (1997), and EEG spectral power and neuronal counts on day 7 since EEG decrease has been related to pronounced damage of neurons on day 7 (Suzuki *et al.*, 1983; Hunter *et al.*, 1995; Peruche *et al.*, 1995). Rectal temperature was also monitored during the first 2 h after recirculation.

The role of TRPV1 was studied using the vanilloid antagonist CPZ peripherally administered.

Methods

Animals

Male Mongolian gerbils (*Meriones unguiculatus*) (Charles River, Calco, Como, Italy) weighing 60–80 g were housed singly in standard laboratory conditions: air-conditioned room ($22 \pm 2^\circ\text{C}$), 12-h light/12-h dark illumination cycle, free access to food and water.

The gerbils were allowed to acclimatize themselves to the environment for a period of 1 week prior to the surgical implantation of cortical electrodes. All gerbils were submitted to EEG electrodes implantation and then divided in different groups on the basis of the treatment received. Each animal was submitted to all tests.

All procedures were approved under Italian Governmental decree No. 94/2000-A.

Surgical procedure

Gerbils were anesthetized with an intraperitoneal injection of chloral hydrate (450 mg kg^{-1} , Sigma, St Louis, MO, U.S.A.) dissolved in saline and given at a volume of 9 ml kg^{-1} . Four electrodes (Bilaney, Dusseldorf, Germany) for EEG recordings were implanted, as described elsewhere (Sala *et al.*, 1997), on the right and left of the parieto-occipital cortex according to the coordinates (anterior +2, posterior -3, lateral 2, ventral 1.6 from bregma) of a brain atlas (Loskota *et al.*, 1974). A further electrode was inserted into the nasal bone as ground. The five electrodes were connected to a pedestal (Bilaney, Dusseldorf, Germany) and fixed with an acrylic cement (Palavit, New Galetti & Rossi, Milan, Italy). After the chronic implantation of the electrodes, 1 week was allowed for recovery from surgery before experiments started.

EEG recording

Freely moving awake gerbils were acclimatized in a sound-attenuated Faraday chamber and then their EEG was recorded for 1 h a day, for 3 days, to determine the basal total and relative spectral power. Spectral powers between 0 and 25 Hz ($0.2\text{--}4.0 \text{ Hz } \delta$, $4.2\text{--}8.0 \text{ Hz } \theta$, $8.2\text{--}13 \text{ Hz } \alpha$, $13.2\text{--}25 \text{ Hz } \beta$) were evaluated using a resolution of 0.2 Hz. The signals were recorded on paper and processed for fast Fourier transform spectral analysis. EEG recordings were also made during ischemia and 7 days after. Each 1 h spectral power was calculated as the mean of six 1 min recordings taken at 10 min intervals.

Cerebral ischemia

After basal EEG recordings, each gerbil was again lightly anesthetized with 2,2,2-tribromoethanol (200 mg kg^{-1} , 10 ml kg^{-1} ; Sigma Aldrich, St Louis, MO, U.S.A.). Body temperature was kept at 37°C throughout surgery with a heating lamp during 10-min ischemia, which was induced by common carotid occlusion as previously described (Braidà *et al.*, 2000). The ischemia was verified qualitatively on paper by the complete flattening of the EEG. A group of animals (sham-operated) was submitted to the same surgical procedure except for carotid clamping.

Body temperature

The experiments were performed in the sound-attenuated Faraday chamber. Controls, sham-operated and ischemic gerbils were trained to the temperature measurements for 5 days before thermoregulator reactions were tested as described by Sala *et al.* (1997). Briefly, after a 1-h acclimatization period in the test room, maintained at 22°C , body temperature was monitored with a rectal thermistor probe (PRA-22002-A, Ellab, Roedovre, Denmark) inserted 3 cm into the colon. After a 30-s equilibration period, the temperature was recorded to the nearest 0.1°C on a CTD 85-M Thermometer (Ellab, Roedovre, Denmark). During temperature measurements, gerbils were unrestrained and were held gently by hand at the base of the tail. Temperature was measured three times (every 10 min) before induction of ischemia (basal) and again at 30, 60, 90 and 120 min after recirculation.

Locomotor activity

Spontaneous motor activity was evaluated as previously described (Braidà *et al.*, 2000) in an activity cage ($43 \text{ cm long} \times 43 \text{ cm wide} \times 32 \text{ cm high}$; Ugo Basile, Varese, Italy) placed in a sound-attenuating room. The cage was fitted with two parallel horizontal infrared beams located 2 cm from the floor. Cumulative horizontal movement counts were recorded for 30 min, every 5 min, 1 day after ischemia.

Passive avoidance task

On the third day after ischemia, each gerbil was submitted to the passive avoidance task. The apparatus (Ugo Basile, Varese, Italy) consisted of a box divided by a guillotine door into two compartments of the same size ($22 \text{ cm long} \times 22 \text{ cm wide} \times 21 \text{ cm high}$) in which the floor had a grid of stainless

steel rod. One compartment was lit with a 10 W electric bulb, and the other compartment was dark. The step-through type passive avoidance task was used as described by Katoh *et al.* (1992), with some modifications. Adaptation was carried out 10 min before the training, and retention 1 day after the training. For the adaptation, a gerbil was placed in the light compartment, and allowed to explore both compartments by leaving the guillotine door open for 5 min. After 10 min, training started during which each gerbil was placed in the light compartment and allowed to enter the dark. When the gerbil entered the dark compartment, the door was automatically closed and an unavoidable scrambled foot shock (1.5 mA) was delivered for 5 s. Each gerbil was then allowed to stay there for 10 s. The procedure was immediately repeated twice. For the retention test, gerbils were placed in the light compartment and the latency to re-enter the dark compartment was recorded up to a maximum of 180 s.

Histology

At 7 days after the ischemic injury, all the gerbils were anesthetized with an overdose of chloral hydrate 5% and transcardially perfused with 4% paraformaldehyde (Sigma Aldrich, St Louis, MO, U.S.A.) for the histological determination as previously described (Braidia *et al.*, 2003). Then, brains were removed and placed in the same fixative overnight. Brains were embedded in paraffin wax. Five serial 5- μ m coronal hippocampal sections at 1.5, 1.7 and 1.9 mm caudal to bregma were cut using a microtome (Leica, Mod. RM2125RT, Solms, Germany) and stained with cresyl violet (Sigma Aldrich, St Louis, MO, U.S.A.). Neurons with a normal appearance in the pyramidal cell layer of the CA1 sector were counted blind (from coded slides) in each section for each group.

Treatment

Gerbils submitted to ischemia were divided in 12 groups of five animals each, receiving s.c. vehicle + vehicle, vehicle + capsaicin (0.01, 0.025, 0.05, 0.2 and 0.6 mg kg⁻¹) (Tocris Cookson Ltd, U.K.), CPZ (0.01 mg kg⁻¹) (Tocris Cookson Ltd, U.K.) + vehicle and CPZ (0.01 mg kg⁻¹) + capsaicin (0.01, 0.025, 0.05, 0.2 and 0.6 mg kg⁻¹). Capsaicin was injected 5 min after recirculation, while CPZ was given 15 min before carotid occlusion. Vehicle was given 5 min after or 15 min before ischemia. Both drugs were dissolved in an appropriate vehicle (Tween 80, ethanol, saline, 1:1:8) and injected in a volume of 10 ml kg⁻¹. A further group, sham-operated, received the same volume of vehicle.

Statistical analysis

Results were presented as mean $\pm$ s.e.m. Total EEG spectral data were expressed as mean of the percent difference vs preischemic value while relative power was expressed as percent of total. For the temperature results, data were expressed as the means of $^{\circ}$ C deviation from the baseline. EEG, motor activity, latency time and neuronal count were analyzed by one-way analysis of variance (ANOVA) for multiple comparisons, followed by Tukey's test. Temperature data were analyzed by repeated measures analysis of variance (two-way ANOVA) with drug treatment as a between-subjects

factor and time as a within-subjects factor. The *post hoc* individual comparisons were performed with Bonferroni's test or Student's *t*-test. The accepted level was $P < 0.05$. All statistical analyses were performed using software Prism version 4 (GraphPad, U.S.A.).

Results

The physiological parameters, such as food and water consumption and body weight, were stable throughout the study in all groups (data not shown).

EEG

A quantitative EEG analysis of gerbils treated with increasing doses of capsaicin is given in Figure 1, in which significant between-group differences were observed in terms of percent vs preischemic value of mean total spectral power ($F_{(11,48)} = 32.913$, $P < 0.0001$, ANOVA), evaluated on day 7. *Post hoc* analysis showed that, in comparison with sham-operated values, the vehicle group had a decrease of EEG power of about 80%. Increasing doses of capsaicin progressively antagonized the ischemia-induced EEG flattening. At higher doses (0.2 and 0.6 mg kg⁻¹), a complete recovery was obtained. CPZ *per se* (0.01 mg kg⁻¹) did not induce any change in comparison with the vehicle group (data not shown). Thus, this dose was chosen to study a possible antagonism. Pretreatment with CPZ only partially antagonized the protective effect observed with capsaicin alone at the doses of 0.2 and 0.6 mg kg⁻¹.

There was a drug treatment effect in the relative distribution of power over frequency bands, except for β ($F_{(12,52)} = 164.00$, $P < 0.0001$, ANOVA for δ) ($F_{(12,52)} = 93.26$, $P < 0.0001$, ANOVA for θ) ($F_{(12,52)} = 74.19$, $P < 0.0001$, ANOVA for α) ($F_{(12,52)} = 0.96$, NS, ANOVA for β) calculated 7 days after

Figure 1 Cortically derived EEG total spectral power evaluated as the difference ($\Delta\%$) from the preischemic value in freely moving awake gerbils given increasing doses (mg kg⁻¹) of capsaicin 5 min after recirculation either singly or in combination with CPZ at a dose of 0.01 mg kg⁻¹ administered s.c. 15 min before bilateral carotid occlusion. Veh = vehicle. Each column represents the mean ($\pm$ s.e.m.) of five animals. ^a $P < 0.001$ compared with sham; ^b $P < 0.001$ compared with capsaicin 0.2 and 0.6; ^c $P < 0.001$ compared with vehicle; ^d $P < 0.05$, ^e $P < 0.001$ in comparison with CPZ + capsaicin (one-way ANOVA followed by Tukey's test).

Table 1 Effect of capsaicin and CPZ on spectral power distribution evaluated 7 days after ischemia, in gerbils

Drug (dose (mg kg ⁻¹ s.c.))	δ (0.2–4.0 Hz)	θ (4.2–8.0 Hz)	α (8.2–13 Hz)	β (13.2–25 Hz)
Sham	45 ± 3.0	44 ± 2.0	10 ± 0.4	1.0 ± 0.3
Vehicle ^a	62 ± 2.0***	32 ± 2.2**	5 ± 1.7**	1.0 ± 0.6
Capsaicin (0.01) ^b	65 ± 1.9***	30 ± 0.8**	4 ± 0.5**	1.0 ± 0.22
Capsaicin (0.025) ^b	58 ± 1.2***	33 ± 1.6**	8 ± 0.9	1.0 ± 0.3
Capsaicin (0.05) ^b	50 ± 1.6**	39 ± 1.2*	10 ± 0.9 ^s	1.0 ± 0.3
Capsaicin (0.2) ^b	38 ± 3.0 ^{sss}	49 ± 3.0 ^{sss}	13 ± 0.8 ^s	1.0 ± 0.3
Capsaicin (0.6) ^b	40 ± 2.0 ^{sss}	48 ± 1.0 ^{ss}	11 ± 0.8 ^s	1.5 ± 0.4
CPZ (0.01)	63 ± 1.0***	31 ± 2.5**	5 ± 1.5**	1.0 ± 0.5
CPZ (0.01) ^c + capsaicin (0.01)	62 ± 1.0***	31 ± 1.0**	2 ± 0.3**	1.0 ± 0.2
CPZ (0.01) ^c + capsaicin (0.025)	62 ± 1.5***	30 ± 2.0**	7 ± 0.9*	1.0 ± 0.4
CPZ (0.01) ^c + capsaicin (0.05)	60 ± 2.0***###	32 ± 2.0***	7 ± 1.0*	1.0 ± 0.2
CPZ (0.01) ^c + capsaicin (0.2)	58 ± 1.5***###	30 ± 1.6***###	11 ± 0.6 ^s	1.0 ± 0.2
CPZ (0.01) ^c + capsaicin (0.6)	50 ± 1.0***###	37 ± .01***###	12 ± 1.0 ^s	1.0 ± 0.3

^aVehicle was given 15 min before carotid occlusion or 5 min after recirculation.

^bCapsaicin was administered s.c. 5 min after recirculation.

^cCPZ was injected s.c. 15 min before carotid occlusion.

Each value represents mean (±s.e.m.) of five animals. CPZ = capsazepine. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ compared with sham-operated group; ^s $P < 0.05$, ^{ss} $P < 0.01$, ^{sss} $P < 0.001$ compared with the vehicle group and CPZ alone; # $P < 0.05$, ### $P < 0.001$ compared with capsaicin alone, same dose (ANOVA followed by Tukey's test).

recirculation. *Post hoc* analyses revealed that ischemia significantly increased the relative EEG power density in the δ , θ and α frequency band in the vehicle-treated group (Table 1). Increasing doses of capsaicin progressively restored the power density distribution. No differences were found between vehicle- and CPZ-alone-treated gerbils, while the antagonist significantly reversed the protective effect of capsaicin starting from the dose 0.05 mg kg⁻¹ shown for δ and θ bands.

Spontaneous locomotor activity

Significant between-group modifications in spontaneous locomotor activity were found ($F_{(11,48)} = 23.638$, $P < 0.0001$, ANOVA) (Figure 2). As expected, the vehicle group showed an increase of motility in comparison to the sham-operated group (Tukey's test). The *post hoc* Tukey's test showed that capsaicin completely antagonized ischemia-induced hyperlocomotion at all tested doses. CPZ (0.01 mg kg⁻¹), which *per se* did not affect ischemia-induced hyperlocomotion (data not shown), reversed the protective effect of capsaicin at doses of 0.01 and 0.025 mg kg⁻¹ (Tukey's test) but it failed to block the protective effect of capsaicin shown at higher doses.

Passive avoidance

The mean latency time revealed a significant difference between groups ($F_{(11,48)} = 35.22$, $P < 0.0001$, ANOVA) (Figure 3). *Post hoc* analyses revealed that vehicle-treated gerbils exhibited a significant impairment as shown by the decrease of mean escape latency in comparison with the sham-operated group. In the range of doses between 0.01 and 0.2 mg kg⁻¹, capsaicin produced a progressive increase of the latency. No effect was observed with the highest dose. When CPZ was given in combination with increasing doses of capsaicin, a complete antagonism was obtained (Tukey's test).

Figure 2 Effect of increasing doses (mg kg⁻¹) of capsaicin given s.c. 5 min after recirculation either singly or in combination with CPZ at a dose of 0.01 mg kg⁻¹ administered s.c. 15 min before bilateral carotid occlusion on spontaneous motor activity evaluated for 30 min, 1 day after ischemia in gerbils. Veh = vehicle. Each column represents the total number of horizontal counts (mean ± s.e.m.) of five animals. ^a $P < 0.001$ compared with sham; ^b $P < 0.001$ compared with corresponding capsaicin alone; ^c $P < 0.01$, ^d $P < 0.001$ compared with vehicle (one-way ANOVA followed by Tukey's test).

CPZ alone did not modify the ischemia-induced mean escape latency (data not shown).

Body temperature

A repeated measures ANOVA revealed a significant dose ($F_{(11,144)} = 55.62$, $P < 0.0001$, ANOVA), time ($F_{(3,144)} = 622.60$, $P < 0.0001$, ANOVA) and interaction effect ($F_{(33,144)} = 55.62$, $P < 0.0001$, ANOVA) of rectal temperature between groups evaluated at 30, 60, 90 and 120 min after recirculation (Figure 4). *Post hoc* test revealed a nonsignificant decrease of

body temperature in both sham-operated and vehicle-treated groups at all tested intervals. Increasing doses of capsaicin progressively decreased the mean rectal temperature at 30 min. At 60 min, a hypothermic effect was still present for capsaicin 0.2 and 0.6, while at 90 min this effect was shown only for the highest dose. No significant hypothermic effect at 120 min was observed. Pretreatment with CPZ, which *per se* did not affect this parameter (data not shown), completely antagonized capsaicin-induced hypothermic effect in the range between

0.025 and 0.2 mg kg⁻¹. The effect produced by the highest dose of capsaicin was reversed by CPZ starting from 60 min.

Histology

Histological examination of the hippocampus at 7 days after recirculation following 10 min of occlusion showed severe alteration in the CA1 region (Figure 5b) in comparison to the sham-operated animals (Figure 5a). Most of the pyramidal cells were darkly stained and shrunk. Increasing doses of capsaicin led to a progressive survival of neurons (data not shown), except for the highest (0.6 mg kg⁻¹). Maximal protection was obtained with capsaicin at 0.2 mg kg⁻¹ (Figure 5c). Pretreatment with CPZ significantly antagonized capsaicin-induced neuroprotective effect (Figure 5d). ANOVA revealed a significant difference in the number of survived neurons in the CA1 region ($F_{(7,32)} = 45.89$, $P < 0.0001$) (Figure 6). There was a mean loss of 80% of neuronal cells in vehicle-treated gerbils in comparison with the sham-operated group. A progressive survival in the animals given increasing doses of capsaicin compared to the vehicle group was obtained in the range of doses between 0.02 and 0.2 mg kg⁻¹, with a peak of 75%.

Discussion

We report here a full protection afforded by the VR1 agonist capsaicin on global cerebral ischemia in the gerbil. The neuroprotection was quantified in terms of complete recovery of total and relative spectral power, spontaneous motor activity, memory function and hippocampal CA1 neuronal density. These parameters are known to be affected by global cerebral ischemia as previously reported (Suzuki *et al.*, 1983;

Figure 3 Escape latency evaluated in the passive avoidance task 3 days after ischemia. Increasing doses (mg kg⁻¹) of capsaicin were given s.c. 5 min after recirculation either singly or in combination with CPZ at a dose of 0.01 mg kg⁻¹ administered s.c. 15 min before bilateral carotid occlusion. Veh = vehicle. Each column represents the mean ($\pm$ s.e.m.) of five animals. ^a $P < 0.01$, ^b $P < 0.001$ compared with sham; ^c $P < 0.001$ compared with corresponding capsaicin alone; ^d $P < 0.001$ compared with vehicle (one-way ANOVA followed by Tukey's test).

Figure 4 Effect of increasing doses (mg kg⁻¹) of capsaicin given s.c. 5 min after recirculation either singly or in combination with CPZ at a dose of 0.01 mg kg⁻¹ administered s.c. 15 min before bilateral carotid occlusion on changes from baseline in rectal temperature. Veh = vehicle. Rectal temperatures were measured starting from 5 min after recirculation. Each column represents the mean ($\pm$ s.e.m.) of five gerbils. ^a $P < 0.05$, ^b $P < 0.01$, ^c $P < 0.001$ compared with sham; ^d $P < 0.001$ compared with corresponding capsaicin alone (two-way ANOVA followed by Bonferroni's test).

Figure 5 Photomicrographs of the hippocampal CA1 region of gerbils with or without 10-min ischemia, evaluated 7 days after ischemia. (a) Sham-operated animal. (b) Ischemic animal treated with vehicle 5 min after recirculation. (c) Ischemic animal treated with capsaicin (0.2 mg kg^{-1}) 5 min after recirculation. (d) Ischemic animal treated with CPZ (0.01 mg kg^{-1}) and capsaicin. Bar = $50 \mu\text{m}$.

Figure 6 Effect of increasing doses of capsaicin (mg kg^{-1}) on neuronal count evaluated 7 days after recirculation in the CA1 region of the hippocampus of sham-operated or ischemic gerbils. Capsaicin was given s.c. 5 min after recirculation. For the antagonism, capsaicin (0.2 mg kg^{-1}) was given in combination with CPZ at a dose of 0.01 mg kg^{-1} given s.c. 15 min before bilateral carotid occlusion. Each column represents the mean ($\pm \text{s.e.m.}$) of five hippocampal sections from the same coronal plane for each animal. $n=5$ for each group. ^a $P<0.05$, ^b $P<0.001$ compared with sham; ^c $P<0.05$, ^d $P<0.01$, ^e $P<0.001$ compared with vehicle; ^f $P<0.001$ compared with capsaicin alone (one-way ANOVA followed by Tukey's test).

Araki *et al.*, 1986; Hunter *et al.*, 1995; Peruche *et al.*, 1995; Sala *et al.*, 1997). In addition, both cognitive and sensorimotor deficits can be found in some human pathologies such as heart attack and coronary artery bypass surgery (Hunter *et al.*, 1998). Furthermore, the histopathology seen in the gerbil is similar to that observed in the hippocampal CA1 region of human brain following cardiac arrest (Hunter *et al.*, 1995). Capsaicin prevented all these mentioned neuropathological consequences of ischemia, suggesting its therapeutic potential.

For all behavioral parameters, a dose-dependent effect was obtained except for spontaneous motor activity, as capsaicin completely reduced ischemia-induced hyperlocomotion, at all tested doses. We can suggest that, even if tested on day 1 after treatment, capsaicin *per se* might exhibit hypolocomotor activity by increasing biosynthesis and release of anandamide, as previously found in rats by Di Marzo *et al.* (2001), using a similar range of doses.

Severe interference with cerebral blood flow has a pronounced effect on electrical activity, which is reflected in EEG flattening. The decrease in EEG spectral power observed 7 days after ischemia has been related to a severe neuronal damage (delayed death), since the cells are replaced by astrocytes in the hippocampal CA1 subfield (Peruche *et al.*, 1995). The protection induced by capsaicin indicates that the compound may contrast the cascade of pathological events that lead to neuronal death. The hippocampus is a preferred area for global ischemia-induced damage and the CA1 sector is the most selectively vulnerable to the reduced cerebral blood flow (Schmidt-Kastner & Freund, 1991). The survival of CA1 neurons obtained 7 days after ischemia with capsaicin (0.2 mg kg^{-1}) may account for the protection against EEG flattening in capsaicin-treated gerbils.

Results obtained from the EEG power spectral analysis indicate changes in the relative spectral composition of vehicle ischemic gerbils. There was a shift toward the lower frequency band, causing an increase in the relative power of the δ band.

The increase in relative power of this frequency band is considered a sign of synchronization of neuronal activity (Frigeni *et al.*, 2004). The increase was associated with a decrease in relative power of the θ and α frequency bands. Comparable EEG alterations have been reported for different kinds of focal brain ischemia and different animals species (Gloor *et al.*, 1977; Hossmann & Schuier, 1980; Kataoka *et al.*, 1987). Following treatment with capsaicin, as opposed to vehicle treatment, there was a significant recovery of spectral power distribution toward basal values, suggesting the presence of recovering neuronal activity.

Capsaicin promoted an important protection against motor hyperactivity. This paradoxical behavior may indicate a reduction of the ischemic animal's ability to form spatial maps (O'Neill & Clemens, 2000) because of the loss of CA1 neurons. These findings are further sustained by memory results. Capsaicin prevented ischemia-induced impairment of memory retention as shown by the longer mean escape latency, in comparison to vehicle-treated ischemic gerbils. It is widely reported that a brief period of cerebral ischemia induces neurological dysfunction in laboratory animals, which is reflected in the deterioration of memory function due to a more vulnerability of CA1 pyramidal cells (LaPoncin-Lafitte *et al.*, 1981; Kiyota *et al.*, 1985).

The protective effect of capsaicin was accompanied by a clear dose-dependent hypothermic effect. Presently, postischemic hypothermia provides the best long-term functional and histological protection after global ischemia (Freund *et al.*, 1990; Colbourne & Corbett, 1995; Colbourne *et al.*, 1999b; Hickey *et al.*, 2000). For instance, prolonged mild hypothermia persistently reduces CA1 zone delayed neuronal death even when cooling begins up to 12 h postischemia (Colbourne *et al.*, 1999a). Hypothermia-induced improvement in postischemic functional recovery has been recently demonstrated, which correlated with decreased production of ascorbyl radicals, considered as a marker of oxidative stress either *in vitro* or *in vivo* (Sharma *et al.*, 1994; Vergely *et al.*, 1998) and in isolated rat heart model (Gambert *et al.*, 2004). Capsaicin given s.c. from 0.1 to 5 mg kg⁻¹ has been reported to cause a fall of body temperature in several animal species (Sasamura & Kuraishi, 1999), acting directly on thermoregulatory neurons of the hypothalamus (Kobayashi *et al.*, 1998; Osaka *et al.*, 2000). Our data suggest that VR1 is required for the hypothermic effect in agreement with Caterina *et al.* (2000) who found no decrease in body temperature in VR1^{-/-} mice when compared with wild-type mice. However, no antagonism on rectal temperature with CPZ (30 mg kg⁻¹) pretreatment in mice given capsaicin (0.3 mg kg⁻¹) i.v. (Di Marzo *et al.*, 2000) was observed. The different animal species, route of administration, doses and experimental protocol used in the present study probably account for this discrepancy.

The protective effect of capsaicin is supported by previous findings in which the VR1 agonist, given i.p. at a dose of 1 mg kg⁻¹, led to neuroprotection against ouabain-induced *in vivo* excitotoxicity in Wistar rats (Veldhuis *et al.*, 2003). The proposed mechanism suggested by the authors was a VR1 desensitization or endocannabinoids biosynthesis stimulation.

In our ischemic model, pretreatment with CPZ, at a dose that *per se* was ineffective, reversed, in a competitive manner, total spectral power, behavioral correlates and body temperature, suggesting that the protective effect of capsaicin was VR1 mediated.

The mechanism by which capsaicin exhibits its protective effect is still unclear. However, it has been reported that during brain injury, *N*-acyl-ethanolamines (including anandamide) and other membrane breakdown products accumulate in the brain (Hansen *et al.*, 2002). These compounds activate VR1 with subsequent Ca²⁺ influx (Szallasi & Blumberg, 1999), glutamate release (Marinelli *et al.*, 2002) and substantial contribution to neuronal excitotoxicity (apoptosis) (Maccarrone *et al.*, 2000; Hail, 2003). Conversely, exogenous compounds capable of quickly desensitizing VR1, such as capsaicin, might exert neuroprotective action by preventing VR1 activation by endogenous stimuli. In this respect, pretreatment with CPZ may prevent VR1 desensitization. A block of capsaicin-induced desensitization has been reported to occur also centrally (Hajos *et al.*, 1987). It can be argued that a simple dose of capsaicin (0.2 mg kg⁻¹) may not be enough to produce vanilloid receptor desensitization. It is well documented that anandamide binds with great affinity to cannabinoid receptor type 1 (CB1) leading to a neuroprotection, most likely *via* a dominant presynaptic inhibition of glutamate release (Shen & Thayer, 1998; Nagayama *et al.*, 1999; Sinor *et al.*, 2000). In addition, a great amount of the endovanilloid ligand anandamide accumulates considerably in response to ischemia (Hansen *et al.*, 2002), which causes a homologous desensitization of the capsaicin-induced response and *vice versa* in human VR1-transfected cells (Smart *et al.*, 2000). Thus, an additive effect of capsaicin and anandamide in producing a massive increase of Ca²⁺, leading to desensitization, can be hypothesized.

The antagonism obtained with CPZ does not agree with the findings of Ray *et al.* (2003), who found that CPZ *per se* protected against cell death in an oxygen glucose deprivation model of organotypic hippocampus slices culture derived from both VR1 knockout and wild-type mice, *via* a non-VR1-mediated mechanism. In addition, a high dose of CPZ (10 mg kg⁻¹) led to neuroprotection as previously reported in an *in vivo* rat model of excitotoxicity (Veldhuis *et al.*, 2003). The low dose and the different model of injury used in the present study probably account for this discrepancy. Experiments are in progress to verify if higher doses of CPZ lead to neuroprotection in our model. A possible explanation for the lack of a protective effect of CPZ may be the presence in the central nervous system of an as yet undefined CB_n/VR_n receptor, as already suggested by different authors (Brooks *et al.*, 2002; Hajos & Freund, 2002; Veldhuis *et al.*, 2003), sensitive to capsaicin and CPZ at a very low dose, which might play a role in neuroprotection.

It is widely reported that large increases in Ca²⁺ influx during and following brain ischemia contribute to both acute and delayed neuronal death. On the other hand, moderate increase in Ca²⁺ influx may activate cell survival programs and promote resistance to hypoxia or ischemia, increasing the antiapoptotic protein, protein kinase B, and the MAP kinase ERK levels. It has been recently reported that calcimycin or ionomycin Ca²⁺ preconditioning greatly reduced cell death in CA1-CA3 and dentate neurons following 7 days after simulated ischemia induced by oxygen and glucose deprivation (Bickler & Fahlman, 2004). It can be argued that our single dose of capsaicin moderately increased Ca²⁺ influx that served as potent preconditioning stimulus *via* VR1 activation, inducing a tolerance of subsequent larger elevation in Ca²⁺ influx. However, since capsaicin was given after recirculation, it is unlikely that it worked as preconditioning stimulus.

Since cytotoxic and vasogenic edema appears during and after bilateral carotid occlusion in gerbils (Hakamata *et al.*, 1997), we cannot exclude the possibility that other mechanisms may have contributed to the protection afforded by capsaicin. For example, capsaicin has potent anti-inflammatory properties by inhibiting the activity of cyclooxygenase-2 and the expression of inducible nitric oxide synthase protein in lipopolysaccharide-stimulated murine peritoneal macrophages (Kim *et al.*, 2003) and *in vivo* via inhibition of nuclear factor- κ B, a transcription factor essentially involved in reactive oxygen species-induced apoptosis (Sancho *et al.*, 2002), through a mechanism not VR1 mediated. Furthermore, the possibility that capsaicin's neuroprotection could involve a release of neuropeptides from sensory neurons cannot be ruled out. A number of studies have shown that capsaicin-induced desensitization

results in a transient depletion of some neuropeptides such as calcitonin gene-related peptide (CGRP), substance P and somatostatin storage (Shin & Hong, 2004). An inhibition of neurogenic inflammation, motor and cognitive deficits development following a traumatic brain injury in rats after depletion of the above neuropeptides has been shown (Nimmo *et al.*, 2004).

In summary, we have demonstrated that capsaicin protected gerbil's brain from global ischemia. This effect was blocked by CPZ, suggesting a VR1 mediation. These findings contribute to develop new exogenous VR1 agonists as valuable targets for therapy against brain injury.

This research was supported by a grant (FIRST 2002) from the Ministry of Scientific Research and Technology.

References

- ARAKI, H., NOJIRI, M., KAWASHIMA, K., KIMURA, M. & AIHARA, H. (1986). Behavioral, electroencephalographic and histopathological studies on mongolian gerbils with occluded common carotid arteries. *Physiol. Behav.*, **38**, 89–94.
- BICKLER, P.E. & FAHLMAN, C.S. (2004). Moderate increases in intracellular calcium activate neuroprotective signals in hippocampal neurons. *Neuroscience*, **127**, 673–683.
- BRAIDA, D., PEGORINI, S., ARCIDIACONO, M.V., CONSALAZ, G.G., CROCI, L. & SALA, M. (2003). Post-ischemic treatment with cannabidiol prevents electroencephalographic flattening, hyperlocomotion and neuronal injury in gerbils. *Neurosci. Lett.*, **346**, 61–64.
- BRAIDA, D., POZZI, M. & SALA, M. (2000). CP 55,940 protects against ischemia-induced electroencephalographic flattening and hyperlocomotion in Mongolian gerbils. *Neurosci. Lett.*, **296**, 183–191.
- BROOKS, J.W., PRYCE, G., BISOGNO, T., JAGGAR, S.I., HANKEY, D.J., BROWN, P., BRIDGES, D., LEDENT, C., BIFULCO, M., RICE, A.S., DI MARZO, V. & BAKER, D. (2002). Arvanil-induced inhibition of spasticity and persistent pain: evidence for therapeutic sites of action different from the vanilloid VR1 receptor and cannabinoid CB(1)/CB(2) receptors. *Eur. J. Pharmacol.*, **439**, 83–92.
- CATERINA, M.J., LEFFLER, A., MALMBERG, A.B., MARTIN, W.J., TRAFTON, J., PETERSEN-ZEITZ, K., KOLTZENBURG, M., BASBAUM, A.I. & JULIUS, D. (2000). Impaired nociception and pain sensation in mice lacking the capsaicin receptor. *Science*, **288**, 306–313.
- COLBOURNE, F. & CORBETT, D. (1995). Delayed postischemic hypothermia: a six month survival study using behavioral and histological assessments of neuroprotection. *J. Neurosci.*, **15**, 7250–7260.
- COLBOURNE, F., LI, H. & BUCHAN, A.M. (1999a). Indefatigable CA1 sector neuroprotection with mild hypothermia induced 6 h after severe forebrain ischemia in rat. *J. Cereb. Blood Flow Metab.*, **19**, 742–749.
- COLBOURNE, F., SUTHERLAND, G.R. & AUER, R.N. (1999b). Electron microscopic evidence against apoptosis as the mechanism of neuronal death in global ischemia. *J. Neurosci.*, **19**, 4200–4210.
- CORTRIGHT, D.N. & SZALLASI, A. (2004). Biochemical pharmacology of the vanilloid receptor TRPV1. An update. *Eur. J. Biochem.*, **271**, 1814–1819.
- DI MARZO, V., BREIVOGEL, C., BISOGNO, T., MELCK, D., PATRICK, G., TAO, Q., SZALLASI, A., RAZDAN, R.K. & MARTIN, B.R. (2000). Neurobehavioral activity in mice of *N*-vanillyl-arachidonylamide. *Eur. J. Pharmacol.*, **406**, 363–374.
- DI MARZO, V., LASTRES-BECKER, I., BISOGNO, T., DE PETROCELLIS, L., MILONE, A., DAVIS, J.B. & FERNANDEZ-RUIZ, J.J. (2001). Hypolocomotor effects in rats of capsaicin and two long chain capsaicin homologues. *Eur. J. Pharmacol.*, **420**, 123–131.
- FREUND, T.F., BUZSAK, I.G., LEON, A. & SOMOGYI, P. (1990). Hippocampal cell death following ischemia: effects of brain temperature and anesthesia. *Exp. Neurol.*, **108**, 251–260.
- FRIGENI, V., MIRAGOLI, L., GROTTI, A. & LORUSSO, V. (2004). Neurotolerability of contrast agents in rats with brain ischemia induced by transient middle cerebral artery occlusion: EEG evaluation. *Invest. Radiol.*, **36**, 1–8.
- GAMBERT, S., BES-HOUTMANN, S., VANDROUX, D., TISSIER, C., VERGELY-VANDRIESSE, C., ROCHETTE, L. & ATHIAS, P. (2004). Deep hypothermia during ischemia improves functional recovery and reduces free-radical generation in isolated reperfused rat heart. *J. Heart Lung Transplant.*, **23**, 487–491.
- GLOOR, P., BALL, G. & SCHAUL, N. (1977). Brain lesions that produce delta waves in the EEG. *Neurology*, **27**, 326–333.
- GUNTHORPE, M.J., BENHAM, C.D., RANDALL, A. & DAVIS, J.B. (2002). The diversity in the vanilloid (TRPV) receptor family of ion channels. *Trends Pharmacol. Sci.*, **23**, 183–191.
- HAIL JR, N. (2003). Mechanisms of vanilloid-induced apoptosis. *Apoptosis*, **8**, 251–262.
- HAJOS, M., JANCOS, G. & ENGBERG, G. (1987). Capsaicin-induced excitation of locus coeruleus neurons. *Acta Physiol. Scand.*, **129**, 415–420.
- HAJOS, N. & FREUND, T.F. (2002). Distinct cannabinoid sensitive receptors regulate hippocampal excitation and inhibition. *Chem. Phys. Lipids*, **121**, 73–82.
- HAKAMATA, Y., HANYU, S., KUROIWA, T. & ITO, U. (1997). Brain edema associated with progressive selective neuronal death or impending infarction in the cerebral cortex. *Acta Neurochir. Suppl. (Wien)*, **70**, 20–22.
- HANSEN, H.S., MOESGAARD, B., PETERSEN, G. & HANSEN, H.H. (2002). Putative neuroprotective actions of *N*-acyl-ethanolamines. *Pharmacol. Ther.*, **95**, 119–126.
- HAYES, P., MEADOWS, H.J., GUNTHORPE, M.J., HARRIES, M.H., DUCKWORTH, D.M., CAIRNS, W., HARRISON, D.C., CLARKE, C.E., ELLINGTON, K., PRINJHA, R.K., BARTON, A.J., MEDHURST, A.D., SMITH, G.D., TOPP, S., MURDOCK, P., SANGER, G.J., TERRETT, J., JENKINS, O., BENHAM, C.D., RANDALL, A.D., GLOGER, I.S. & DAVIS, J.B. (2000). Cloning and functional expression of a human orthologue of rat vanilloid receptor-1. *Pain*, **88**, 205–215.
- HICKEY, R.W., FERIMER, H., ALEXANDER, H.L., GARMAN, R.H., CALLAWAY, C.W., HICKS, S., SAFAR, P., GRAHAM, S.H. & KOCHANAK, P.M. (2000). Delayed, spontaneous hypothermia reduces neuronal damage after asphyxial cardiac arrest in rats. *Crit. Care Med.*, **28**, 3511–3516.
- HOSMANN, K.A. & SCHUIER, F.J. (1980). Experimental brain infarcts in cats. *Stroke*, **11**, 583–592.
- HUNTER, A.J., GREEN, A.R. & CROSS, A.J. (1995). Animal models of acute ischaemic stroke: can they predict clinically successful neuroprotective drugs? *Trends Pharmacol. Sci.*, **16**, 123–128.
- HUNTER, A.J., MACKAY, K.B. & ROGERS, D.C. (1998). To what extent have functional studies of ischaemia in animals been useful in the assessment of potential neuroprotective agents? *Trends Pharmacol. Sci.*, **19**, 59–66.

- KATAOKA, K., GRAF, R. & ROSNER, G. (1987). Differentiation between cortical and subcortical lesions following focal ischemia in cats by multimodality evoked potentials. *J. Neurol. Sci.*, **79**, 117–127.
- KATOH, A., ISHIBASHI, C., SHIOMI, T., TAKAHARA, Y. & EIGYO, M. (1992). Ischemia-induced irreversible deficit of memory function in gerbils. *Brain Res.*, **577**, 57–63.
- KIM, C.S., KAWADA, T., KIM, B.S., HAN, I.S., CHOE, S.Y., KURATA, T. & YU, R. (2003). Capsaicin exhibits anti-inflammatory property by inhibiting I κ B- α degradation in LPS-stimulated peritoneal macrophages. *Cell. Signal.*, **15**, 299–306.
- KIYOTA, Y., HAMAJI, K., MIYAMOTO, M. & NAGAO, M. (1985). Effect of idebenone (CV-2619) on memory impairment observed in passive avoidance task in rats with cerebral embolization. *Jpn. J. Pharmacol.*, **37**, 300–302.
- KOBAYASHI, A., OSAKA, T., NAMBA, Y., INOUE, S., LEE, T.H. & KIMURA, S. (1998). Capsaicin activates heat loss and heat production simultaneously and independently in rats. *Am. J. Physiol.*, **275**, R92–R98.
- LAPONCIN-LAFITTE, M., GROSDÉMOGUE, C., ROYBILLON, C., POTRAT, P., LESPINASSE, P. & RAPIN, J.R. (1981). Short-term memory and cerebral ischemia: pharmacological application. *Eur. Neurol.*, **20**, 265–269.
- LOSKOTA, W.J., LOMAX, P.L. & VERITY, M.A. (1974). *A Stereotaxic Atlas of the Mongolian Gerbil (Meriones uingulatus)*, pp. 1–157. Ann Arbor: Ann Arbor Science Publishers.
- MACCARRONE, M., LORENZON, T., BARI, M., MELINO, G. & FINAZZI-AGRO, A. (2000). Anandamide induces apoptosis in human cells via vanilloid receptors. Evidence for a protective role of cannabinoid receptors. *J. Biol. Chem.*, **275**, 31938–31945.
- MARINELLI, S., VAUGHAN, C.W., CHRISTIE, M.J. & CONNOR, M. (2002). Capsaicin activation of glutamatergic synaptic transmission in the rat locus coeruleus *in vitro*. *J. Physiol.*, **543**, 531–540.
- MEZEY, E., TOTH, Z.E., CORTRIGHT, D.N., ARZUBI, M.K., KRAUSE, J.E., ELDE, R., GUO, A., BLUMBERG, P.M. & SZALLASI, A. (2000). Distribution of mRNA for vanilloid receptor subtype 1 (VR1), and VR1-like immunoreactivity, in the central nervous system of the rat and human. *Proc. Natl. Acad. Sci. U.S.A.*, **97**, 3655–3660.
- NAGAYAMA, T., SINOR, A.D., SIMON, R.P., CHEN, J., GRAHAM, S.H., JIN, K. & GREENBERG, D.A. (1999). Cannabinoids and neuroprotection in global and focal cerebral ischemia and in neuronal cultures. *J. Neurosci.*, **15**, 2987–2995.
- NIMMO, A.J., CERNAK, I., HEATH, D.L., HU, X., BENNETT, C.J. & VINK, R. (2004). Neurogenic inflammation is associated with development of edema and functional deficits following traumatic brain injury in rats. *Neuropeptides*, **38**, 40–47.
- O'NEILL, M.J. & CLEMENS, J.A. (2000). Preclinical models of neurologic and psychiatric disorders. In: *Current Protocols in Neuroscience*, ed. Crawley, J.N., Gerfen, C.R., McKay, R., Rogawski, M.A., Sibley, D.R. & Skolnick, P. pp. 9.5.1–9.5.25. New York: Wiley.
- OSAKA, T., KOBAYASHI, A., NAMBA, Y., INOUE, S. & KIMURA, S. (2000). Lack of integrative control of body temperature after capsaicin administration. *Korean J. Intern. Med.*, **15**, 103–108.
- PERKINS, M.N. & CAMPBELL, E.A. (1992). Capsazepine reversal of the antinociceptive action of capsaicin *in vivo*. *Br. J. Pharmacol.*, **107**, 329–333.
- PERUCHE, B., KLAASSENS, H. & KRIEGLSTEIN, J. (1995). Quantitative analysis of the electrocorticogram after forebrain ischemia in the rat. *Pharmacology*, **50**, 229–237.
- RAY, A.M., BENHAM, C.D., ROBERTS, J.C., GILL, C.H., LANNEAU, C., GITTERMAN, D.P., HARRIES, M., DAVIS, J.B. & DAVIES, C.H. (2003). Capsazepine protects against neuronal injury caused by oxygen glucose deprivation by inhibiting I(h). *J. Neurosci.*, **23**, 10146–10153.
- SALA, M., LEONE, M.P., LAMPUGNANI, P., MATTURRI, L. & GORI, E. (1997). Polydeoxyribonucleotide (defibrotide) protects against post-ischemic behavioural, electroencephalographic and neuronal damage in the gerbil. *Eur. J. Pharmacol.*, **328**, 143–152.
- SANCHO, R., LUCENA, C., MACHO, A., CALZADO, M.A., BLANCO-MOLINA, M., MINASSI, A., APPENDINO, G. & MUNOZ, E. (2002). Immunosuppressive activity of capsaicinoids: capsiate derived from sweet peppers inhibits NF- κ B activation and is a potent antiinflammatory compound *in vivo*. *Eur. J. Immunol.*, **32**, 1753–1763.
- SASAMURA, T. & KURAISHI, Y. (1999). Peripheral and central actions of capsaicin and VR1 receptor. *Jpn. J. Pharmacol.*, **80**, 275–280.
- SCHMIDT-KASTNER, R. & FREUND, T.F. (1991). Selective vulnerability of the hippocampus in brain ischemia. *Neuroscience*, **40**, 599–636.
- SHARMA, M.K., BUETTNER, G.R., SPENCER, K.T. & KERBER, R.E. (1994). Ascorbyl free radical as a real-time marker of free radical generation in briefly ischemic and reperfused hearts – an electron paramagnetic resonance study. *Circ. Res.*, **74**, 650–658.
- SHEN, M. & THAYER, S.A. (1998). Cannabinoid receptor agonists protect cultured rat hippocampal neurons from excitotoxicity. *Mol. Pharmacol.*, **54**, 459–462.
- SHIN, H.K. & HONG, K.W. (2004). Importance of calcitonin gene-related peptide, adenosine and reactive oxygen species in cerebral autoregulation under normal and diseased conditions. *Clin. Exp. Pharmacol. Physiol.*, **31**, 1–7.
- SINOR, A.D., IRVIN, S.M. & GREENBERG, D.A. (2000). Endocannabinoids protect cerebral cortical neurons from *in vitro* ischemia in rats. *Neurosci. Lett.*, **278**, 157–160.
- SMART, D., GUNTHORPE, M.J., JERMAN, J.C., NASIR, S., GRAY, J., MUIR, A.I., CHAMBERS, J.K., RANDALL, A.D. & DAVIS, J.B. (2000). The endogenous lipid anandamide is a full agonist at the human vanilloid receptor (hVR1). *Br. J. Pharmacol.*, **129**, 227–230.
- SUZUKI, R., YAMAGUCHI, T., KIRINO, T., ORZI, F. & KLATZO, I. (1983). The effects of 5-min ischemia in Mongolian gerbils: I. Blood–brain barrier, cerebral blood flow, and local cerebral glucose utilization changes. *Acta Neuropathol.*, **60**, 207–216.
- SZALLASI, A. & BLUMBERG, P.M. (1999). Vanilloid (Capsaicin) receptors and mechanisms. *Pharmacol. Rev.*, **51**, 159–212.
- SZALLASI, A., NILSSON, S., FARKAS-SZALLASI, T., BLUMBERG, P.M., HOKFELT, T. & LUNDBERG, J.M. (1995). Vanilloid (capsaicin) receptors in the rat: distribution in the brain, regional differences in the spinal cord, axonal transport to the periphery, and depletion by systemic vanilloid treatment. *Brain Res.*, **703**, 175–183.
- VAN DER STELT, M. & DI MARZO, V. (2004). Endovanilloids. Putative endogenous ligands of transient receptor potential vanilloid 1 channels. *Eur. J. Biochem.*, **271**, 1827–1834.
- VELDHUIS, W.B., VAN DER STELT, M., WADMAN, M.W., VAN ZADELHOFF, G., MACCARRONE, M., FEZZA, F., VELDINK, G.A., VLIEGENTHART, J.F., BAR, P.R., NICOLAY, K. & DI MARZO, V. (2003). Neuroprotection by the endogenous cannabinoid anandamide and arvanil against *in vivo* excitotoxicity in the rat: role of vanilloid receptors and lipoxygenases. *J. Neurosci.*, **15**, 4127–4133.
- VERGELY, C., MAUPOIL, V., BENDERITTER, M. & ROCHETTE, L. (1998). Influence of the severity of myocardial ischemia on the intensity of ascorbyl free radical release and on postischemic recovery during reperfusion. *Free Radic. Biol. Med.*, **24**, 470–479.

(Received July 22, 2004

Revised November 4, 2004

Accepted November 24, 2004)