Short cuts

What's new in the other general journals

People with AIDS are increasingly dying of unrelated diseases

People with AIDS are living longer, thanks to highly active retroviral treatment, and an increasing proportion are surviving their infection long enough to die of unrelated illnesses such as cardiovascular disease. This is certainly true in New York city, where recent research shows that more than a quarter of people with AIDS die from something else—mostly drug overdose, heart disease, or a cancer unrelated to HIV infection.

This proportion increased by a third (from 19.8% to 26.3%, $P = 0.015$) between 1999 and 2004. The authors and a linked editorial agree there is a growing need for doctors to take a more holistic view of people with AIDS, possibly by focusing less on their infection and more on their general health. Both suggest shifting the care for these patients towards primary care, in which people with AIDS would benefit from standard practices, such as advice about quitting smoking and other drugs, cancer screening, and chronic disease management for hypertension and diabetes.

In the five years of this study, annual mortality in people with AIDS fell faster for deaths related to HIV than for deaths from other causes, but a low CD4 count was an effective predictor for both. Injecting drug users had higher mortality than gay men.

Ann Int Med 2006;145:397-406

Modern management of pre-eclampsia seems safer for babies

Many pregnancies complicated by pre-eclampsia now end in deliberate preterm delivery of the infant. Delivery cures the pre-eclampsia and may prevent a stillbirth, but it exposes the infant to all the risks of being born too early. Has the trend towards earlier intervention saved babies, or simply shifted their deaths from the womb to the postnatal period?

Observational data from Norway's almost complete register of births is encouraging. In a study of all first singleton births between 1967 and 2003, the risk of stillbirth fell substantially in pre-eclamptic pregnancies, and the risks of death between birth and 5 years of age stayed about the same. In the same period, inductions before 37 weeks rose from 8% of pre-eclamptic pregnancies to nearly 20%.

Although trends can be hard to interpret, the authors think it likely that earlier intervention in pre-eclampsia has prevented stillbirths, and improvements in postnatal care have kept mortality down after delivery. The difficult decisions made daily by Norwegian obstetricians seem to have worked in favour of the babies, but there's still the possibility of a net increase in neonatal morbidity and childhood disability, outcomes that were impossible to measure in this study.

JAMA 2006;296:1357-62

Inflammatory marker might predict older women's risk of fracture

Further evidence that inflammation is somehow linked to bone metabolism, and

by extension to bone mass and risk of fracture, comes from a cohort study of 444 typical Australian women older than 65 years. The women were recruited initially for a study looking at the prevalence of osteoporosis. At the start of the study they gave baseline blood samples, which the researchers tested for high sensitivity C reactive protein, a marker of systemic inflammation. Over a median follow-up of 5.5 years, 102 women had a fracture. The researchers found a link between baseline serum concentrations of the marker and fractures, such that each increase by one standard deviation was associated with a 23% rise in fracture risk (unadjusted hazard ratio 1.23, 95% CI 1.01 to 1.51). The link was unexplained by differences in bone mineral density, history of fractures, markers of bone turnover (C telopeptide and bone specific alkaline phosphatase), diet, lifestyle, drug treatments, or other illness. Perhaps high sensitivity C reactive protein is involved in remodelling processes that weaken bone, write the authors. If so, it could be a useful marker for risk of fracture in older women, a possibility that deserves further study.

JAMA 2006;296:1353-5

Invasive treatment works best for acute coronary syndrome

In 1999 Scandinavian researchers showed that for people with an acute coronary syndrome, early angiography followed by revascularisation if necessary works better than a more conservative approach using drugs. The first trial lasted only two years, however. To find out if the benefits lasted any longer, the Scandinavian researchers

continued to follow-up their 2457 patients using the Swedish, Danish, and Norwegian national registers that record deaths and hospital admissions.

Patients who had an early angiogram coupled with revascularisation, as appropriate, were still doing better five years later than patients treated with drugs alone, largely because they had had fewer heart attacks (12.9% *v* 17.7%; relative risk 0.73, 95% CI 0.6 to 0.89).

The invasive strategy had no impact on long term risk of death, however, and the other benefits were confined to men, non-smokers, and patients classified as moderate or high risk at the start of the trial. In this subgroup, the invasive strategy reduced the risk of death or myocardial infarction from 28.1% to 21.2% at five years (relative risk 0.75, 0.64 to 0.89) but again had no significant effect on deaths alone.

A linked editorial (pp 966-9) concludes that there seems to be clinically important long term advantage to early angiography and revascularisation for these patients, so long as they are stratified first by risk. National guidelines from Europe and America agree.

Lancet 2006;368:998-1004

Trials of bone marrow cells for damaged hearts yield modest, mixed results

Despite promising studies in animals, convincing evidence that bone marrow cells can help repair damaged hearts in humans remains elusive. The three latest trials reported mixed results, and a linked editorial (pp 1274-7) says we still have a long way to go before cellular treatments for ventricular dysfunction and heart attack become mainstream, if they ever do.

Part of the problem is the complexity of the treatments being tested. These trials evaluated three different cell populations, two from bone marrow and one from circulating blood. The most successful trial (and the largest) reported a small improvement in left ventricular ejection fraction in patients who had been given an intracoronary infusion of progenitor cells from bone marrow three to seven days after successful reperfusion treatment for an acute myocardial infarction (5.5% improvement *v* 3.0% in the placebo group, *P*=0.01). The benefits were evident at four months, but it's unclear how long they will last. A second trial reported no improvement in left ventricular function after intracoronary infusion of mononuclear cells from the bone marrow after myocardial infarction. The third compared bone marrow cells with cells derived from circulating blood of patients with healed myocardium. The authors found a small benefit from the bone marrow cells only.

Progress may be slow, but we should not give up, says the editorial. Bigger trials with longer follow-up will help refine the

treatments and perhaps target them better at people with the most to gain.

New Engl J Med 2006;355:1199-209, 1210-21, 1222-32

Tenth of older US residents have valvular heart disease

Valvular heart disease has been largely ignored as a public health problem since the decline of rheumatic heart disease, and the epidemiology of valve problems in modern industrialised populations has received little attention. To fill some of the gaps, researchers from the United States trawled through all population based studies held by the National Heart, Lung, and Blood Institute to derive a cohort of nearly 12 000 normal US adults who had been offered an echocardiogram. In all, 5.5% of them had valvular heart disease. Adjusting for the national age and sex distribution—which was different from the distribution in their cohort—the authors estimate that 2.5% of US citizens have valvular heart disease, most commonly mitral regurgitation. A sister study of all echo cardiograms done in Olmsted County, Minnesota, between 1990 and 2000 found a prevalence of 1.8%. The difference is probably due to undiagnosed disease in the general population.

Both studies found that prevalence of all types of valve disease rose significantly with age, reaching 13% for people aged 75 years or older. They also found that valve disease was associated with an increased risk of death (relative risk 1.36, 95% CI 1.15 to 1.62). These figures point to a large and growing burden of disease in the US, which can only get worse as the population ages.

Lancet 2006;368:1005-11

Many men who have sex with men describe themselves as straight

When researchers asked a random sample of almost 4200 male New Yorkers about their sexual orientation, 91.3% said they

were straight and 3.7% said they were gay. Towards the end of each lengthy telephone survey, the researchers asked them how many men and how many women they had had sex with in the previous year. Almost one in 10 (9.4%) of the “straight” men reported having sex with at least one man but no women. Put another way, about 12% of the 2898 men who had answered both questions had had exclusively male relationships in the past year, but most of them described themselves as straight. Does it matter?

As a group, the men who had homosexual relationships but described themselves to researchers as straight had fewer partners than the men who identified themselves as gay. But they were less likely to have used a condom during their last encounter (adjusted prevalence ratio 0.5, 95% CI 0.3 to 1.0) and less likely to have had a test for HIV than men who identified themselves as gay (0.6, 0.4 to 0.9). These findings have implications for doctors assessing HIV risk and for sexual health strategies aimed at gay men, say the authors. Both may need to be refocused on specific sexual activity, rather than self reported sexual orientation.

Ann Int Med 2006;145:416-25

Overseas aid misses world's poorest women and children

In 2004, rich countries and donor agencies gave \$2bn (£1.0bn; €1.6bn) to help poor countries improve the health of mothers, newborn babies, and young children, according to recent estimates. The 60 poorest countries with the highest rates of infant and child mortality got just \$1.4bn, or \$3.10 for each child. Data given by donors to the Organisation for Economic Development and Cooperation show that only 2% of overseas development aid is earmarked for women, newborn babies, and children. Health economics researchers from London warn that it's nowhere near enough to meet the millennium development goals for child survival and maternal health. Their best guesses put the shortfall at about \$7bn a year. The deadline is 2015.

Overall, Africa receives about half of this meagre funding, and South East Asia about a fifth, but the researchers found considerable variation between countries in these and other regions. They also found that countries with the greatest need don't necessarily receive the most aid, presumably because of the complex geopolitical, strategic, and historical factors that determine where the money goes.

Donor countries and agencies have already promised to increase their aid, but whether the funds will find their way to the world's neediest women and children remains to be seen.

Lancet 2006;368:1077-87

Alison Tonks *associate editor*
atonks@bmj.com