reviews

BOOKS • CD ROMS • ART • WEBSITES • MEDIA • PERSONAL VIEWS • SOUNDINGS

ART

Leonardo da Vinci: Experience, Experiment and Design

An exhibition at the Victoria and Albert Museum, London, until 7 January 2007
www.vam.ac.uk/leonardo

Rating: ★★★★★

In the broad sweep of Renaissance intellectual curiosity, few if any thinkers wielded their cerebral brooms through as many rooms or compartments of human endeavour as Leonardo da Vinci (1452-1519). His painting of an enigmatic young woman, celebrated for centuries, now arguably depicts the most recognised half smile in the world. Many of his drawings, such as the naked man inscribed in a circle, are universally recognised as being his work and have become icons representing our human quest for a rational understanding of how we and our world work.

Drawn solely from British collections, this exhibition of 62 drawings by Leonardo

demonstrates his insatiable curiosity and how he explored and developed ideas, using pen and paper to try to fathom the underlying mechanisms of how animals and machines function, rather than simply recording their external appearance. They are primarily dynamic sketches, drawn quickly in order to capture and investigate animal movement and physical forces, not finished drawings intended to be mounted and framed and presented to patrons. Most of the sheets are also covered with his handwritten observations and conclusions, which are as perceptive as his sketches.

Vertical and Horizontal Sections of the Human Head and Eye, drawn around 1489, shows that Leonardo followed the traditional view that mental activity occurred in the cerebral ventricles, rather than in the cerebral cortex. He thought that the eye was the primary means for obtaining knowledge about the external environment and that visual knowledge was the only kind that could be verified. He often shows morphological analogies from botany or engineering in his drawings; in this case, his cross section of an onion alludes to the layered membranes that cover the eye and brain.

Drawn almost three decades later around 1508, *Vessels of the Arm with Comparison of Vessels in the Old and the Young* is a drawing showing the straight course taken by the superficial blood vessels of a young person's arm alongside a much smaller sketch of the tortuous course of those of an

Men at Work, Digging, Carrying, Pulling etc (c 1509)

old man's arm. During the winter of 1507-8 Leonardo had dissected the body of a 100 year old man, paying particular attention to the cadaver's vascular system. From his earlier observations that meandering channels slowed down the flow of water in rivers and their branches, he concluded that death had resulted from silted-up blood vessels, exemplifying his ability to synthesise knowledge from disparate disciplines.

Men at Work, Digging, Carrying, Pulling etc is one of two sheets filled with sketches of gangs of little men in motion, drawn in black chalk around 1509, included in the exhibition. It exhibits Leonardo's desire to consider all these different types of human movement in terms of mechanical laws. He carried notebooks with him, in order to capture his observations, including workers busily constructing new roads and buildings in the rapidly expanding Italian cities of the Renaissance; however, the figures sketched on this larger sheet of paper were probably drawn from memory.

"We see that someone who takes up weight with one of his arms naturally thrusts the other arm outside himself in an opposite direction, and if this is not sufficient to ensure his equilibrium, he bends so as to impose as much of his weight as suffices to counteract the acquired weight," is typical of the notes Leonardo jotted.

It is seldom that we have the opportunity to study a genius at work, particularly by looking at drawings which are fresh to the public as are those shown in London.

Colin Martin independent consultant in healthcare communication, London
Cmpubrel@aol.com

Items reviewed are rated on a 4 star scale
(4=excellent)

Left: *Vertical and Horizontal Sections of the Human Head and Eye* (c 1489). Right: *Vessels of the Arm with Comparison of Vessels in the Old and the Young* (c 1508)

PERSONAL VIEW

“Did you ever suffer from any mental illness?”

At one of my psychotherapy sessions the first thing my client asked me, after the usual greetings and smiles, was, “Did you ever suffer from any mental illness?” I had been engaged in therapy with this particular client for a few weeks. We had a good rapport, and in terms of the therapy I felt we were making better than average progress. But nothing in the previous sessions had prepared me for this question. I had never had the opportunity to discuss the nature and degree of self disclosure that would be appropriate to a particular context. I had always and only understood self disclosure as a personal matter: no rules, no guidelines, only discretion. I have not had any diagnosable mental illness; but now, faced with this opening question and put on the spot, I felt a salvo of thoughts shooting through my mind. How should I answer this question? I was apprehensive of losing our rapport, the engagement in therapy, and progress in future sessions. Will my client engage better if I am able to deliver a positive response? Is this type and degree of identification necessary? In my mind I thus intellectualised the situation, and, while struggling to maintain my confident and unruffled appearance, I reverted to using that old therapist’s trick of asking the Socratic question: “I wonder why you ask me that?” “Just like that,” came the answer, accompanied by an uncertain smile. I am certain my client was now wondering whether this had been an appropriate question. Sensing this discomfort, I waffled on, albeit defensively, about stress and individual vulnerability, hoping that my status as the trusted therapist, the all knowing and good doctor, was undamaged despite my not having been “empathic” enough.

Looking back, I wonder about this ethical quagmire of self disclosure. What constitutes empathy? How much of it does the therapist need? How is it measured, and how does it need to be expressed for a good interaction—a therapeutic relationship? What degree of self disclosure is appropriate? When does it become violation of a boundary? What constitutes excessive? What is professional and what personal?

It is not that I have not witnessed senior psychiatrists and therapists, my consultants and trainers, breathing magic into a session by disclosing some personal detail about their life experience or friends, families, or partners. However, having had a mental illness is a more deeply personal issue, and some of my colleagues would agree that sharing such experience with a patient helps

the therapist to relate better to the patient, to be more trusted. Even a simple disclosure such as, “You seem to have flu—I was off sick two weeks back with it, and it’s so annoying,” generates a personal touch that may leave the patient leaving your consulting chamber more satisfied. Once I saw one of my colleagues suddenly bring to life a therapy session after disclosing the ordeal suffered when a family member was given the diagnosis of an ominously advanced carcinoma. I have also seen the effect of a consoling hand on a shoulder or an empathic, assuring hand on an anxious and tremulous one.

But I continue to wonder at this. Is it necessary? If so, with whom and when, and how much? Speaking hypothetically, how important is it for achieving therapeutic benefit to disclose that my wife has diabetes or my daughter has epilepsy? Natural curiosity gets the better of you when you are trusting important details about yourself to a stranger. And if, say, I used Prozac, should I make

everyone I prescribe it to, and who may a bit uncertain about taking it, aware that it worked and that the side effects didn’t last long? And if it didn’t work for me, should I stop prescribing it? For three years now I have attended induction sessions at the beginning of every six months of my basic psychiatric training, and never have the answers to such questions or even the questions themselves figured there. I had discussed the issue with my trainers, the summary of whose answers is, “Use your discretion.”

But is my judgment free from error? And do all doctors have a similar degree of discretion? So many factors would need to be taken into account. I would imagine that the age and sex of my patient matters, how he or she presents and behaves in consultation, his or her past experiences, vague resemblances, attire, presenting complaint, and degree of distress. Whether somebody was accompanying the patient might also be a factor, as well as non-verbal communication and the patient’s understanding of his or her needs. Self disclosure and tactile assurance may be seen as minor matters, but the boundary between our personal and professional identities is not watertight. I wonder how important it is to get this aspect of the therapeutic relationship right; it is so easy to get it wrong.

Arnob Chakraborti senior house officer, psychiatry, South Staffordshire NHS Trust
ac97dmc@yahoo.co.uk

Will my client engage better if I am able to deliver a positive response

SOUNDINGS

Illness behaviour

We summarise the notes of patients joining the practice and put them on to the computer. The odd thing is that, often, if you have summarised the notes yourself, you almost seem to know the patient as they walk in through the door for the first time. In particular, you have a feel for their illness behaviour.

In general practice there is a high “noise to signal” ratio of symptoms that signify self limiting rather than significant disease. Frequently, particularly with an elderly population, a general practitioner may be told of 10 or more symptoms in one consultation. Each symptom potentially has an algorithm or guideline associated with it. And these patients are not a small subgroup of “somatisers.” They will be normal patients, usually slightly beyond middle age, who are becoming aware of their mortality and are no longer dismissing symptoms as minor, as they did 20 years ago.

Looking back, I suspect that the NHS survived because general practitioners investigated or referred hardly anything. They were, in David Haslam’s memorable phrase, “the risk sink of the NHS.” They soaked up these symptoms at minimal cost. Sometimes this may have been merely because they were poor clinicians, but often it was because they knew their patients.

I realise that my own practice of investigation or referral is very much influenced by the knowledge of prior illness behaviour. It is only one of a myriad of other variables that come into the decision, but it is an important one. When I see a discharge summary of a patient admitted through casualty with alarming symptoms I am not surprised that all the scans were normal. I may know that, although the patient is delightful and hard working, the whole family “catastrophises” symptoms. Their described symptoms are never of moderate severity.

It would be difficult to include illness behaviour in any algorithm, or indeed, to explain your inaction under scrutiny if a patient turns out to have serious disease. Yet if such considerations are lost in future, if a patient is always assessed by a new doctor without knowledge of prior health seeking behaviour, I worry that the NHS will collapse under the weight of undiluted algorithmic investigation—and that people will spend much of their lives in radiology departments.

Kevin Barraclough general practitioner, Painswick, Gloucestershire