⁷ Jackson, P., *et al.*, *British Medical Journal*, 1970, 4, 295.⁸ Livingstone, E. S., *Canadian Medical Association Journal*, 1971, 105, 1065.⁹ Marshall, S., and Lyon, R. P., *Journal of the American Medical Association*, 1972, 219, 1753.¹⁰ Moss, W. M., *Fertility and Sterility*, 1972, 23, 33.¹¹ Wallace, D. M., and Riddle, P., *British Medical Journal*, 1971, 9, 100.¹² Dodds, D. J., *Journal of the American Medical Association*, 1972, 220, 1498.¹³ Pugh, R. C. B., and Hanley, H. G., *British Journal of Urology*, 1969, 41, 340.¹⁴ Pryor, J. P., *et al.*, *British Medical Journal*, 1971, 1, 272.¹⁵ Singh, S. M., personal communication, 1973.¹⁶ Wig, D. N., and Singh, S., *Indian Journal of Medical Research*, 1972, 60, 1386.¹⁷ Wolfers, H., *British Medical Journal*, 1970, 4, 297.¹⁸ Yeates, W. Keith, personal communication, 1972.¹⁹ Rolnick, H. C., *Journal of Urology*, 1954, 72, 911.²⁰ Alken, C. E., and Ferner, H., *Urologia Internationalis*, 1960, 10, 66.²¹ Blandy, J. P., *Operative Surgery*, ed. C. Rob and R. Smith, Vol. 12, 2nd edition. London, Butterworths, 1971.²² Blandy, J. P., *British Journal of Hospital Medicine*, 1973, 9, 319.²³ Cornes, J. S., *Lancet*, 1973, 1, 721.²⁴ Jackson, L. N., *Lancet*, 1973, 1, 770.²⁵ Halim, A., and Blandy, J. P., *British Medical Journal*, 1973, 3, 110.

Contemporary Themes

Thoughts in Mind of a Person with Cancer

G. W. MILTON

British Medical Journal, 1973, 4, 221-223

Summary

The paper is based on the personal experience of many patients suffering from cancer. It analyses in a discursive way thoughts which may be present in the minds of cancer patients. The paper also briefly discusses the techniques used to minimize the common fears of the cancer patient.

Introduction

Of all the words in the English language probably none causes such personal dread as the word cancer, a word often believed synonymous with a death sentence.

This paper is based on a personal experience of more than 1,000 patients with malignant melanoma, and much experience with other malignant diseases. It is an attempt to condense my understanding of their thoughts at different stages of the disease.

Stages of Thought

The thoughts of a patient with malignancy pass through several stages, the first being when he realizes that he has a disease which could possibly be malignant; the second when the malignancy is established but the condition is thought to be curable by the patient, his relatives, and the doctors; the third is when the disease is obviously no longer curable—recognized first by the doctor and before very long by the patient and his relatives; and the fourth and last stage is that of the dying patient.¹ Though the thoughts of the cancer patient generally fall into these stages there are, of course, exceptions because there are many different ways of coping with bad news. The stages may also overlap or be arrested by the patient who is unwilling to adjust to the realization of the progress of his disease.

Presentation of cancer may occur in one of two ways, and the mode of presentation will influence to some extent the patient's initial reaction to his knowledge of his disease. In one type the patient has what appears to him to be a minor complaint; one which has caused him no pain, no loss of strength or vitality and he presents to his doctor confident that with some trivial treatment he will be permanently cured. In the second method the patient has suffered from a variety of symptoms for weeks or months which have aroused his suspicions that all is not well—for example, steady loss of weight and appetite or chronic gnawing pain which causes him gradually to suspect that he may have a malignant disease.

All of us in times of great distress need some minimal time to become mentally adjusted to misfortune; a sort of acclimatization period. With a malignant disease which develops gradually this acclimatization is part of the developing process. If the patient has no idea that he has a malignant disease then the sudden "death sentence" (as he believes) can quite distort his judgment until he has had time to adjust. The transition of a healthy person into stage one of the mental adjustment to cancer may therefore be either gradual or sudden.

Stage One

The first reaction of the patient in this stage begins when the "icy fingers" of contemplated death touch him. His initial reaction is therefore one of fear. He is frightened of several things—for instance, of dying and the unknown, of being dead, of leaving loved ones and happy associations. Another fear is that of the effects of treatment, even if successful. Few surgeons seem to fully appreciate the dread that many patients feel at the thought of a mastectomy, colostomy, or radical lymph node dissection. The patient's mind is therefore torn by conflicting fears, of the disease and death on the one hand, and mutilation from treatment on the other. Additionally, the patient may also be most apprehensive least he should reveal himself to be a coward and may try by various deceptions to conceal his terror.

While these dreads are developing in his mind another feeling may arise; a suspicion relating to his next of kin and his doctors. "Are they telling me the truth?" The wonderful account in *Othello* shows how the most trivial evidence can be accepted as concrete proof of bad news to the suspicious mind. The patient, once his suspicions have been fully aroused, will analyse every word said by the surgeon, intern, ward sister, and even hospital

University of Sydney, Melanoma Clinic, Sydney Hospital, Sydney, N.S.W., Australia

G. W. MILTON, M.B., F.R.A.C.S., Professor of Surgery

porters, to try to find out if any remark indicates a good or bad omen. Suspicion is based on fear and fear increases the suspicion and so the mind of a person in the early stages of malignant process is torn by fear, doubt, and suspicion, all of which feed one on another.

Stage Two

As the patient moves from stage one into stage two when the diagnosis becomes established another emotion becomes grafted on to the previous ones. This could best be expressed as "Why me?" Most people think that misfortunes will always happen to someone else. In other words as the diagnosis of malignancy becomes established it strikes home to the patient for the first time that the dreaded disease of cancer is not something that affects other people, "This time it's *me*." This is a terrible personal blow and it requires a lot of adjustment because the natural response to this problem "Why me?" tends to be one of self pity and resentment, which are the next emotions felt by the patient with malignancy. "Why should I get this?" "I never did any harm to anybody, not like some people I know," "I never have any luck." The egocentricity as shown by self pity often leads the patient into being both irascible and unreasonable, "They don't understand what I have to face," "You don't really care what happens to me."

Final Stages

When the disease recurs after treatment and the patient realizes that attempts at cure have failed he begins to move emotionally into stage three, and a feeling which began in earlier stages becomes more powerful. This is a feeling of loneliness, which consists of both insulation and isolation from the community. Many writers have expressed this in a variety of ways. Solzhenitzyn² described the tumour as growing like a wall behind the patient; the patient's family and loved ones are on one side of the wall and the patient himself is *alone* on the other side. Dr. Johnson in one of his essays³ wrote that "every inhabitant of the earth must walk downward to the grave alone and unregarded, without any partner of his joy or grief, without any interested witness of his misfortune or success." This feeling of loneliness as malignancy develops becomes very intense and is often made a great deal worse by inept management of the dying patient.¹ The loneliness will also vary from time to time; during the day so long as the patient is in company and has something to occupy his mind he may well be able to suppress his loneliness, but at night, particularly if he has pain and the rest of the house is asleep, the feeling of isolation becomes almost overpowering.

The mental state of a patient with malignant disease is often very seriously injured by a tantalizing process when the disease appears to be cured and he is in high spirits, and then follows a relapse and a further attempt at cure, followed by another relapse. This situation is the very essence of the torture of Tantalus as each hopeful episode is shattered by the prospect that his life is to be amputated. Such periods of hope and despair require great understanding on the part of his doctor and relatives. It is not unusual in these periods for the patient to develop symptoms which are similar to those of any anxiety state—for example, headaches. There are specific fears suffered by the cancer patient and these gradually increase in intensity as hope of cure begins to diminish. They include a fear of loss of dignity and of pain.

FEARS

Lack of dignity is often not discussed by the patient but it lurks in his mind because he is unfamiliar with death except possibly in extreme old age. The senile patient may have been converted in the mind of those near him from a person to a "thing" and the

patient with cancer may believe this is a function of dying rather than senility. The idea of becoming a thing could have been implanted in his mind by past hospital experience. He may have been in a ward and seen a terminal patient almost ignored by nurses, doctors, and visitors. The ward sister, while standing at the end of the bed of such a dying patient, may have issued a staccato directive to a junior nurse, "This patient must be turned every two hours, nurse," and move on without a backward glance. The doctors whisper to the sister at the end of the bed "Nothing to be done here, sister" and hurry past. Not a reassuring sight for one expecting the ordeal himself. The fear of pain is very real, as nearly everyone knows of a relative or friend who was supposed to have died in agony from cancer.

Throughout the whole process of mentally adjusting to the facts of cancer the patient is often plagued with depression. On rare occasions some patients lapse into a state of total indifference. This seems to resemble the mental attitude which occurs in primitive peoples preceding death from witchcraft, and will be discussed in detail elsewhere.

DEFENCE MECHANISMS

If the above represents the thoughts that go on in the mind of a patient with malignant disease it is important to understand that patients cope with their thoughts by a variety of techniques. Basically, there are three defence mechanisms—one is denial;⁴ two is withdrawal; three is accepting reality.

Denial of the facts of death is extremely common in our century, we do it both collectively and individually. To kill a man in warfare we need to depersonalize him into an initial—for example, for Vietcong—V.C. or use a slang expression that converts men into objects (gook, wog). The medical profession is certainly not immune to this, as in some medical articles animals are "sacrificed" and patients "succumb." When confronted with real death we tend to be horrified. Realistically simulated death and brutality on the other hand are obviously highly entertaining, a fact well understood by the entertainment industry. It follows that as a massive denial of death has been part of most of our lives, when the dread word cancer is mentioned one of the standard methods of coping with it from an individual point of view is denial. The denial is apparent by a variety of behavioural patterns shown by the patient. He may accept that the doctors have told him the truth as they see it but equally will be adamant that they are wrong. Secondly, he may show considerable bonhomie, and the young intern may be amazed with the patient who is facing up to his situation incredibly well. He is jocular, cheerful, will make fun of various methods of treatment by describing chemotherapeutic agents as cocktails, soup, or some other flippant phrase. His attendance at the clinic will be accompanied by much giggling from the nurses as he chats to everyone. It is worthwhile to recognize that this state of false jocularity may suddenly break down and if it does there is a metamorphosis of the patient's appearance and attitude from one of excessive good cheer to one of indifference and lethargy, referred to above.

A fourth defence mechanism which the patient may use is to accept the diagnosis but reject the consequences. This is not the same as denial, but its effects may be far worse for the patient and his family, because such a person is a "sitting duck" to be "skinned" of all his money as he vainly seeks magic cures from all who profess to be able to cure cancer. The common remark of many such patients is, "well, the cure can't do any harm and, who knows, it may do me some good. It's worth a try."

Coping with Cancer

Coping with cancer is a problem which has to be faced by a group of people—patient, nurses, and doctor in the middle, family and associates around the periphery. If the above is a true account of the emotions felt by a patient the following is an

outline of the principles of diminishing these unfavourable emotions as much as possible. The patient who initially has no idea of his malignancy but who nevertheless is going to require surgery for its treatment must, I believe, be told the cause of his symptoms and the justification for his treatment, but it is very important for him to have a period of acclimatization in which to adjust to the fact that he has cancer. It follows that if at the first encounter the doctor thinks his patient has cancer, then when the problem of diagnosis is discussed the doctor should imply that malignancy is a possible diagnosis. The patient then has some diagnostic tests and when he returns for the results he should be asked to come with his wife so that if the final blow has to be coped with they are together at the interview. I believe the latter to be important because the key to excessive suffering in this situation comes from suspicion, fear, loneliness, and self pity. If the patient is aware that his doctor is saying one thing to him and something else to his wife the obvious inference is that his wife is being told the truth which is too unpleasant for him to hear. The surgeon must see the patient with his wife alone and not surrounded by a formidable entourage of white coats and serious faces. The surgeon must have time enough to answer questions and be gentle enough to feel some sympathy for the man and his family.

The key to management therefore is gentle understanding, time, and honesty; it is not a long face and "I am so sorry for you." The patient's specific fears are dealt with one by one in discussion and positive things are stressed rather than negative. For example, ways of relieving pain are outlined if the patient

starts to worry about pain. Above all the care of the patient with cancer must be combined with understanding the thoughts passing through his mind and discussing them frankly with him and his family.

The patient must not feel that the surgeon or physician to whom he turned for treatment when his disease appeared curable has now abandoned him, "There is nothing more I can do for you, go back to your family doctor." There is a great deal that the specialist can do in the way of emotional support and this may be very valuable. Abandonment of the incurable patient by the specialist has far-reaching effects: not only does the patient and his family feel disappointed, even betrayed, but also the junior staff in training from interns to nurses learn that the "great man" abandons an incurable patient and may put this into practice in their own working life.

Kindly understanding by a team which includes the specialist, the family doctor, nurses, and the patient and his family, will not eliminate the suffering of the cancer patient but will diminish it—a modest achievement but one worth obtaining.

References

- ¹ Milton, G. W., *Medical Journal of Australia*, 1972, 2, 177.
- ² Solzhenitzyn, Alexander, *Cancer Ward*. Harmondsworth, Penguin Books, 1972.
- ³ Johnson, Samuel, *Rasselas, in Poems and Selected Prose*, ed. B. H. Bronson, p. 180. New York, Holt, Rinehart and Winston Inc., 1958.
- ⁴ Maddison, D., *Archives of the Foundation of Thanatology*, 1970, 2, 000.

Medicine in Old Age

Role of Day Hospital Care

J. C. BROCKLEHURST

British Medical Journal, 1973, 4, 223-225

Geriatric medicine initiated an early form of progressive patient care in hospital—through acute admission ward, to rehabilitation ward, and continuing care ward. The geriatric day hospital is a logical extension of this system and forms a bridge between the hospital and the community. The ever-increasing numbers and proportion of elderly people, particularly the very old, in our society inevitably increase the incidence of disabling disease. The provision of day care for the treatment and support of elderly disabled people has been an attempt, attended with a good deal of success, to prevent the admission to hospital of more and more of these old people.

Social Day Care

Day care for the elderly includes a wide variety of social and hospital provisions which must be clearly distinguished. Social

day care is provided either by local authorities or by voluntary organizations (for example, old peoples welfare committees)—or occasionally by thoughtful employers. Social day care itself offers a whole spectrum of facilities.¹ Anderson's definition of social day care is as follows: "Clubs and social centres exist for all old people, to increase social contact and to give scope and facilities for new pursuits in retirement".

Several main types of day care are now available. Day care centres, most usually instituted by local authorities, provide a substantial amount of personal care to at least a small proportion of clients. Referral is usually through general practitioners or social workers. There are a wide range of services, including bathing, chiropody, meals, and provision of low priced foods. Psychogeriatric day care centres are similar to day care centres but require a greater degree of supervision. Their purpose is more that of a crèche than a club. Since it is now the Department of Health and Social Security's policy to support old people suffering from dementia, who are neither physically handicapped nor pose behavioural problems, in the community the importance of psychogeriatric day care centres is likely to increase.²

Both these types of social day centres may be provided by local authorities under the 1968 Health Services and Public Health Act and both require transport to bring the clients to them.

Social centres or social clubs come in many varieties; some are open seven days a week, others only one half day a fortnight.

Department of Geriatric Medicine, University Hospital of South Manchester, M20 8LR

J. C. BROCKLEHURST, M.D., F.R.C.P., Professor