Thymectomy for Myasthenia Gravis

SIR,—In your leading article (2 September, p. 543) you discussed the evidence for the existence of a transmissible agent responsible for neonatal myasthenia gravis. Dr. A. D. Korczyn offered an alternative suggestion (30 December, p. 793) implicating anticholinesterase drugs. If the latter were true a higher incidence of neonatal myasthenia would occur in women taking high doses of medication or for more prolonged periods, or both. This, however, is not the case. Furthermore, several cases of neonatal myasthenia gravis have been reported where the mother was not receiving medication.¹ In addition, drug requirements decrease in approximately one-third of pregnant women. Other reports provide data that do not support Dr. Korczyn's suggestions.²

It has become increasingly evident that thymectomy is followed by improvement and stable remissions in a significant percentage of cases.³ Yet no attempt has been made to investigate differences in the incidence of neonatal myasthenia between the pre-thymectomy and postthymectomy groups. In any such analysis it is important to consider the clinical classification of the disease⁴ because: (1) children born to women with ocular myasthenia gravis do not develop neonatal myasthenia;¹ (2) thymectomy is not generally performed in ocular myasthenia, and therefore this category should be excluded; (3) the severity of the disease (that is, the clinical classification at the peak of symptoms) influence the prognosis both before and after thymectomy; and (4) symptoms during pregnancy are variable and do not necessarily reflect the severity of the disease.

Review of the reported series shows a wide discrepancy in the incidence (4-50%).¹ Among the lowest (4%) is the one reported by Fraser⁵ consisting predominately of patients who had undergone thymectomy. A previous report from this institution⁶ recorded 10 out of 69 (14%) cases of neonatal myasthenia. We have now examined the incidence of neonatal myasthenia among 800 myasthenic women (190 postthymectomy). The incidence of neonatal myasthenia gravis in women who had not undergone thymectomy was twice that in the thymectomy group. Our late colleague Dr. K. E. Osseman⁶ indicated that 26% of pregnancies in myasthenic mothers resulted in spontaneous abortion. Our present review indicates that spontaneous abortions are more common in the non-thymectomy group. Several of the more severe cases of myasthenia, particularly in the non-thymectomy group, had their pregnancies interrupted for medical reasons, and in any comparison of the incidence of neonatal myasthenia these factors must be considered.

Our preliminary data confirm that the variability of the symptoms during pregnancy does not appear to be related to the incidence of neonatal myasthenia gravis. Mothers who gave birth to children with neonatal myasthenia were primarily those in whom the disease reached the more severe forms either before or after the pregnancy. It is obvious that when this factor is taken into consideration the difference is even more pronounced. The effects of thymectomy on myasthenia may be delayed for months or even years.³ All cases of neonatal myasthenia in our thymectomy group occurred in women

who had not as yet reached complete remission.

These data seem to indicate that the presence of the thymus and possibly a thymic factor play a role in neonatal myasthenia gravis.—We are, etc.,

ANGELOS E. PAPATESTAS
GABRIEL GENKINS
ALLAN E. KARK

Mount Sinai Hospital and School of Medicine,
City University,
New York

- 1 Namba, T., Brown, S. B., and Grob, D., *Pediatrics*, 1970, 45, 488.
- 2 Stern, G. M., Hall, J. M., and Robinson, D. C., *British Medical Journal*, 1964, 2, 284.
- 3 Papatestas, A. E., Alpert, L. I., Osseman, K. E., Osseman, R. S., and Kark, A. E., *American Journal of Medicine*, 1971, 50, 465.
- 4 Osseman, K. E., *Myasthenia Gravis*. New York, Grune and Stratton, 1958.
- 5 Fraser, D., and Turner, J. W. A., *Proceedings of the Royal Society of Medicine*, 1963, 56, 379.
- 6 Osseman, K. E., in *Medical, Surgical and Gynecologic Complications of Pregnancy*, ed. J. J. Rovinsky and A. F. Guttmacher. Baltimore, Williams and Wilkins, 1965.

Subclinical Brucellosis

SIR,—Dr. Eirian Williams (31 March, p. 791) writes that in brucellosis the term "subclinical" "should be used with care." What is meant by this statement? Subclinical is a precise and self-explanatory term to describe a perfectly obvious state. Why then should it be used with care? If Dr. Williams does not recognize subclinical infection or doubts that it exists, then he is simply ignoring the observations of others; the subclinical state in brucellosis is well documented.¹⁻⁵

Further on in his article Dr. Williams admits that in symptomless veterinary surgeons who are continually exposed to brucella antigen all the usual serological tests are positive in high dilution. With the use of what term other than subclinical does he describe their infection?—I am, etc.,

R. J. HENDERSON

Public Health Laboratory,
Royal Infirmary,
Worcester

- 1 Cayton, H. R., Osborne, A. D., and Sylvester, D. G. H., *Lancet*, 1967, 2, 836.
- 2 Drole, P., *Archives of Internal Medicine*, 1932, 50, 373.
- 3 Foley, B. V., Clay, M. M., and O'Sullivan, D. J., *Irish Journal of Medical Science*, 1970, 3, 457.
- 4 Foley, B. V., and O'Flynn, J. G. A., *Irish Veterinary Journal*, 1971, 25, 42.
- 5 Henderson, R. J., *Lancet*, 1967, 2, 353.

Toxoplasma gondii Oocysts in the Faeces of Naturally Infected Cat

SIR,—In the course of a general survey of gastrointestinal parasites in domestic animals we found *Toxoplasma gondii* oocysts in one stray cat (out of 250 examined). Reports about naturally infected cats¹⁻⁴ are extremely rare, and this finding (the first case described in Italy) is reported because of its considerable epidemiological importance to man.

A 4-month-old male stray cat in apparently good health was captured in the outskirts of Bologna. A dye test (Aagaard's method⁵) showed a titre of < 1:10. His faeces were mushy and dark brown and microscopic examination showed many *Isospora felis* and *T. gondii* oocysts after concentration in sodium nitrate and sugar solution (specific gravity 1300). *T. gondii* oocysts kept in 2.5% potassium dichromate sporulated after 16 days. The sporulated oocysts were

washed in saline in a centrifuge and injected intraperitoneally into 5 dye-test-negative mice (< 1:10). Oocysts were also given orally to two cats (2-month-old), one of them (cat A) showing a dye test titre of < 1:10 and the other (cat B) of 1:50.

After four weeks all the inoculated mice had a dye test of > 1:250 and the microscopic examination of their brains showed typical toxoplasma cysts. The faeces from cat A were negative for *T. gondii* throughout the whole observation period (45 days) but a dye test on the 16th day reached 1:250. This fell to 1:50 on the 26th day and remained unchanged. On the third day occasional sporulated oocysts were found in the faeces of cat B, these apparently representing a transit of the ingested inoculum. The faeces became negative on the following day. On the 27th day new, abundant, and unsporulated *T. gondii* oocysts appeared. They persisted for three days. Later, however, the faeces remained negative. The original dye-test titre of 1:50 reached > 1:1,250 on the 16th day and remained so till the animal was killed (45th day). *T. gondii* cysts were isolated in mice injected with brain material from cat B.

These data confirm that cats may be responsible for the natural spread of *T. gondii* oocysts. The oocyst displays its infectivity after some days at sporont maturation (16 days in the present case) but sometimes seems to be already infective at emission (transit oocysts from contaminated food). The marked rise of the dye test titre associated with emission of oocysts in cat B (already positive before the experimental infection) seems to show that cats who have previously had *T. gondii* infection may become spreaders again by a renewed oral load of parasites. The results of the dye test in cats excreting *T. gondii* oocysts do not seem to represent a reliable indication for epidemiological investigations on cats, because their range may vary from comparatively high values (cat B) down to nothing (naturally infected stray cat).—We are, etc.,

S. PAMPIGLIONE
G. POGLAYEN
B. ARNONE

Cattedra di Parassitologia,
Facoltà di Medicina Veterinaria,
Università di Bologna,
Italy

F. DE LALIA

Istituto di Malattie Infettive,
Università di Milano,
Centro Studi sulla Toxoplasmosi,
Milano, Italy

- 1 Jacobs, L., in *Advances in Parasitology*, vol. 5, ed. B. Dawes. London, Academic Press, 1967.
- 2 Wallace, G. D., *Journal of Infectious Diseases*, 1971, 124, 227.
- 3 Janitschke, K., and Kühn, D., *Berliner und Münchener Tierärztliche Wochenschrift*, 1972, 85, 46.
- 4 Wermer, J. K., and Walton, B. C., *Journal of Parasitology*, 1972, 58, 1148.
- 5 Berengo, A., de Lalla, F., Bechelli, G., and Cavallini-Sampieri, L., *Minerva Medica*, 1969, 60, 377.

Smoking and Ischaemic Heart Disease

SIR,—It is stated in the interesting paper by Dr. Nicholas Wald and others (31 March, p. 761) that "there is a well recognized correlation between smoking and atherosclerosis." The data cited in support show, however, that there is an association between smoking and clinical ischaemic heart disease. It is at least possible that these are not two identical propositions.

It is true that death from myocardial ischaemia is unlikely in the absence of coronary atheroma but, as Morris¹ has said, many are susceptible to the latter but far