Annotations

Pain and analgesia in the newborn

The traditional view that neonates do not feel pain is now being questioned. Early studies showed what was thought to be a decorticate response in babies with no localisation to painful stimuli and this, together with the belief that the painful experience was not remembered, implied that pain was unimportant.¹ As a consequence and because of certain anxieties about the handling of drugs (particularly opiates) in the newborn there has been a reluctance to prescribe analgesia for these patients.² Data now suggest that infants do feel pain.^{1 3 4} In this review the underlying neurobiology of pain development in neonates is described, followed by a review of clinical studies indicating that the newborn baby responds (possibly adversely) to pain. We conclude that the arguments in favour of the use of analgesia have become overwhelming.

Neurobiology of pain development

The limited evidence available shows that innervation of the peripheral tissue and the basic connections between primary sensory neurones and the cells in the dorsal horn of the spinal cord occurs early in fetal development. In this review we concentrate on those events that seem to occur later in fetal life or in the early postnatal period. In rats the central neurophysiological effects of unmyelinated C fibres, the group of afferents that are particularly concerned in chronic and inflammatory pain, mature gradually over the first two to three postnatal weeks.^{5 6} This maturation is associated with several anatomical and chemical changes. The concentrations of neuropeptides found within the C fibres (such as substance P, somatostatin, and vasoactive intestinal polypeptide) increase considerably during the early postnatal period.⁷ Furthermore, at birth binding sites for substance P are in high concentrations and distributed diffusely over the grey matter, and only become defined in specific laminae and reduced in number as the rat matures.⁸ Whether there is a similar delayed development of C fibres in man is not known, but there is a pronounced increase in the concentrations of the above neuropeptides around the time of birth.^{7 9}

The activity of interneuronal pathways in the dorsal horn, especially relevant to pain processing,

also develops postnatally. The spinal cord lacks inhibitory control at birth and although the synaptic linkage between afferents and dorsal horn cells is still weak, receptive fields are large and single stimuli can often evoke long lasting excitation that lasts for several minutes; repeated stimuli can build up considerable background activity in the cells.⁵ The result is that otherwise weak cutaneous inputs are made more effective centrally. The flexor reflex, a useful measure of nociceptive function at the spinal cord level, is highly exaggerated in neonates.^{10 11} Thresholds are lower and the flexion withdrawal lasts longer. Repeated skin stimulation results in considerable hyperexcitability or sensitisation, which is particularly pronounced before 35 weeks' gestation in the human and the eighth postnatal day in rats.¹¹ Nevertheless there is a marked reduction in the threshold of this reflex after skin injury in the newborn that is analogous to the tenderness or hyperalgesia experienced in adults.¹² Inhibitory pathways descending from the brain stem are ineffective at birth,¹³ perhaps because their neurotransmitters (5-hydroxytryptamine and noradrenaline) appear only postnatally.¹⁴

Development of opioid sensitivity

Another important postnatal event is the maturation of the opioid/opiate system in the spinal cord. No analgesic effects of morphine can be detected in the rat pup until the seventh postnatal day and adult sensitivity is only reached at the 14th postnatal day, exactly coinciding with the appearance of high affinity mu-receptor binding in the rat cord. K-receptors and ketocyclazocine analgesia mature earlier at the seventh to 10th postnatal days.¹⁵ Delta receptors develop last at the 12th postnatal day but the mu:delta ratio does not reach 1:1 until the 26th postnatal day.¹⁶ Little is known about opiate receptor development in the human cord, but it apparently begins before birth. Morphine and other opioids are effective analgesics in newborn infants born both prematurely and at full term, although dose response studies have not been undertaken.^{1 4} Enkephalin appears postnatally (first postnatal day) in the rat and is found later than most other peptides in the human fetal dorsal horn at 12–14 weeks.¹⁷

Assessment of pain in the newborn infant

Three types of response to painful stimuli have been studied in the newborn human infant.

AUTONOMIC RESPONSES

Changes in heart rate, blood pressure, transcutaneous partial pressure oxygen, and palmar sweating are common autonomic responses to pain. In newborn babies, lancing of the heel produces sustained increases in heart rate in full term infants but the responses are less clear in the preterm baby.¹⁸ The palmar sweating response to this stimulus does not appear until 37 weeks' gestation.¹⁹ Local anaesthesia prevents the heart rate and blood pressure changes associated with circumcision.²⁰

BEHAVIOURAL RESPONSES

Diffuse body movement and purposeful withdrawal have been quantified in response to painful stimuli²¹ and have been used to show that the preterm infant is, if anything, supersensitive to painful stimuli when compared with the full term infant.¹¹ These responses can be reduced by local analgesia (M Fitzgerald, C Millard, N McIntosh, unpublished observations). Facial expressions have also been linked with painful stimuli but vary with the behavioural state.²² Crying (? primitive communication) has been evaluated clinically and by spectrographic analysis. A painful cry is different from that indicating hunger or fear in a full term baby, but the differentiation is difficult in the preterm infant.²³

HUMORAL AND METABOLIC RESPONSES

Anand *et al* studied the response to surgical operations in neonates.²⁴ A pronounced release of catecholamines, growth hormone, glucagon, aldosterone, and other corticosteroids was seen when insulin secretion was suppressed. These responses were associated with the breakdown of fat and the release of carbohydrates resulting in increased concentrations of glucose, lactate, pyruvate, ketone bodies, and non-esterified fatty acids in the blood, as well as an increase in the breakdown of protein.⁴ Potent anaesthetic agents (for example, fentanyl and halothane) inhibited these stress responses²⁵ and preliminary data suggested that the stress response might be associated with increased postoperative morbidity and mortality.¹⁴

Conclusions

Laboratory studies have shown that the basic connections required for painful stimuli to be transmitted to the central nervous system are present in the newborn, despite the considerable organisation

and maturation (particularly of control systems) that occurs postnatally. Clinical studies of the pain responses mounted by newborn babies have confirmed that these pathways are functional, albeit somewhat variable. Furthermore, in some cases these behavioural changes persisted after the painful procedure, implying that the experience was remembered.²⁶ This behaviour may disrupt both the infants' adaptive responses to feeding and the mother/infant reactions.²⁷ Anaesthetic agents reduce the stress response after neonatal operations. It is difficult at this stage to evaluate whether a profound stress response is good or bad. Stability of physiology is important in the preterm infant: fluctuations of arterial oxygen pressure may be potentially dangerous to the eye,²⁸ and variability in blood pressure has been linked with the occurrence and extension of intraventricular haemorrhage.²⁹ The suggestion, however, that there are fewer side effects of operation when potent anaesthetic agents are used,²⁴ and that postoperative mortality and morbidity are reduced, mean that the judicious provision of analgesia is important, independent of whether the pain is remembered.

References

- McGrath PJ, Unruh AM, eds. *Pain in children and adolescents. Pain research and clinical management*. Vol 1. Amsterdam: Elsevier, 1987.
- Purcell-Jones G, Dormon F, Sumner E. Paediatric anaesthetists' perceptions of neonatal and infant pain. *Pain* 1988;**33**:181-7.
- Fitzgerald M. Pain and analgesia in neonates. *Trends in Neuroscience* 1987;**10**:344-6.
- Anand KJS, Hickey PR. Pain and its effects in the human neonate and fetus. *N Engl J Med* 1987;**317**:1321-9.
- Fitzgerald M. The postnatal development of cutaneous afferent fibre input and receptive field organization in the rat dorsal horn. *J Physiol* 1985;**364**:1-18.
- Fitzgerald M. The development of activity evoked by fine diameter cutaneous fibres in the spinal cord of the newborn rat. *Neurosci Lett* 1988;**86**:161-6.
- Marti E, Gibson SJ, Polak JM, *et al*. Ontogeny of peptide and amino-containing neurons in motor, sensory and autonomic regions of rat and human spinal cord. *J Comp Neurol* 1987;**266**:332-59.
- Charlton CG, Helke CJ. Ontogeny of substance P receptors in the rat spinal cord: quantitative changes in receptor number and differential expression in specific loci. *Developmental Brain Research* 1986;**29**:81-6.
- Charnay Y, Chayvialle JA, Pradayrol L, Bouvier R, Paulin C, Dubois PM. Ontogeny of somatostatin-like immunoreactivity in the human foetus and spinal cord. *Developmental Brain Research* 1987;**36**:63-73.
- Ekholm J. Postnatal changes in cutaneous reflexes and in the discharge pattern of cutaneous articular sense organs. *Acta Physiol Scand* 1967;**297**(suppl 7):1-130.
- Fitzgerald M, Shaw A, McIntosh N. Postnatal development of the cutaneous flexor reflex: comparative study of preterm infant and newborn rat pups. *Dev Med Child Neurol* 1988;**30**:520-6.
- Fitzgerald M, Millard C, McIntosh N. Hyperalgesia in premature infants. *Lancet* 1988;**i**:292.
- Fitzgerald M, Koltzenburg M. The functional development of descending inhibitory pathways in the dorsolateral funiculus of

- the newborn rat spinal cord. *Developmental Brain Research* 1986;**24**:261–70.
- ¹⁴ Bregman BS. Development of serotonin immunoreactivity in the rat spinal cord and its plasticity after neonatal spinal cord lesions. *Developmental Brain Research* 1987;**34**:245–64.
 - ¹⁵ Barr GA, Paredes W, Erickson KL, Zukin RS. K-opioid receptor-mediated analgesia in the developing rat. *Developmental Brain Research* 1986;**29**:145–52.
 - ¹⁶ Wohltmann M, Roth BL, Coscia CJ. Differential postnatal development of mu and delta opiate receptors. *Developmental Brain Research* 1982;**3**:679–84.
 - ¹⁷ Charnay Y, Paulin C, Dray F, Dubois PM. Distribution of enkephalin in human fetus and infant spinal cord: an immunofluorescence study. *J Comp Neurol* 1984;**223**:415–23.
 - ¹⁸ Owens ME, Todt EH. Pain in infancy: neonatal reaction to a heel lance. *Pain* 1984;**20**:77–86.
 - ¹⁹ Harpin CA, Rutter N. Development of emotional sweating in the newborn infant. *Arch Dis Child* 1982;**57**:691–5.
 - ²⁰ Williamson PS, Williamson ML. Physiological stress reduction by a local anesthetic during newborn circumcision. *Pediatrics* 1983;**71**:36–40.
 - ²¹ Frank LS. A new method to quantitatively describe pain behaviour in infants. *Nurs Res* 1986;**35**:28–31.
 - ²² Grunau RVE, Craig KD. Pain expression in neonates: facial action and cry. *Pain* 1987;**28**:395–410.
 - ²³ Michelsson K, Jarvenpaa A-L, Rinne A. Sound spectrographic analysis of pain cry in preterm infants. *Early Hum Dev* 1983;**8**:141–9.
 - ²⁴ Anand KJS, Brown MJ, Causon RC, Christofides ND, Bloom SR, Aynsley-Green A. Can the human neonate mount an endocrine and metabolic response to surgery? *J Pediatr Surg* 1985;**20**:41–8.
 - ²⁵ Anand KJS, Sippell WG, Aynsley-Green A. Randomised trial of fentanyl anaesthesia in preterm neonates undergoing surgery: effects on the stress response. *Lancet* 1987;**i**:243–8.
 - ²⁶ Marshall RE, Stratton WC, Moore JA, Boxerman SB. Circumcision. I. Effects upon newborn behaviour. *Infant Behaviour and Development* 1980;**3**:1–14.
 - ²⁷ Marshall RE, Porter FL, Rogers AG, Moore JA, Anderson B, Boxerman SB. Circumcision. II. Effects upon mother-infant interaction. *Early Hum Dev* 1982;**7**:367–74.
 - ²⁸ Venus V, Patel KC, Pratap KS, Konchigeri H, Vidyasagar D. Transcutaneous PO₂ monitoring during pediatric surgery. *Crit Care Med* 1981;**9**:714–16.
 - ²⁹ Goodard J, Lewis RM, Armstrong DL, Zeller RA. Moderate rapidly induced hypertension as a cause of intraventricular hemorrhage in the newborn beagle model. *J Pediatr* 1980;**96**:1057–60.

M FITZGERALD

Department of Anatomy and Developmental Biology,
University College,
London WC1E 6BT
and

N MCINTOSH

Department of Child Life and Health,
University of Edinburgh,
17 Hatton Place,
Edinburgh EH9 1NW