The classical caesarean section would avoid both of these problems, although it is associated with more postoperative morbidity and makes a repeat caesarean section very likely. For these reasons it is not applicable to every patient in progressive preterm labour with a breech presentation, but it is perhaps worthy of consideration in some cases.

J P CALVERT

University Hospital of Wales,
Cardiff CF9 4XW¹ Karp, L E, *et al*, *Obstetrics and Gynecology*, 1979, 53, 88.

Eyes run on light

SIR,—Mr M J Gilkes (23 June, p 1681) quoted numerous references pointing out a lacuna in the articles recently published under my name. He could understandably have been on the brink of using the term "black hole," which is as I understand it a disaster area in the universe that is inexplicable in normal scientific terms. He also referred to the fact that common sense is so often ignored because it is self-evident. Self-evidence can also be a black hole in all our judgments on visual impairment.

Common sense tells some people that weak eyes are strengthened by glasses or protected by them from becoming weaker, but others take the contrary common sense view that eyes are weakened by glasses and should be made to work without support from such crutches.

Common sense tells us that we have to conserve our eyesight like savings and that it can be used up like a bank account: that it can be preserved if we don't use artificial light, the wrong glasses, or perhaps any glasses, and don't read because of the strain. Common sense also tells us that use of a skill improves it, like sewing, skiing, or sailing. Common sense, which we all run on, is a primitive thread in all social, intellectual, and scientific patterns. Neither science in such a dilemma about black holes, based on analysis of fact, nor common sense confused by its reliance on contradictory self-evidence, is an ultimate authority.

The conclusion is that because neither science nor common sense can answer our questions consistently people might well resort to revelation.

Mr Gilkes has revealed a simple truth which should cast a light even in the confused and tortuous warrens of our general and primitive ignorance about basic visual processes.

P A GARDINER

Guy's Hospital,
London SE1 9RT

Rate of change of haemoglobin A₁ in diabetic patients

SIR,—We read with interest the reports of the change in concentration of glycosylated haemoglobin (HbA₁) occurring in newly diagnosed diabetic patients following treatment (14 April, p 979, and 7 July, p 19). Our own observations support the view of Leslie and others that "conclusions about diabetic control drawn from a single HbA₁ estimation should be treated with caution," and that "a single HbA₁ estimation may relate to a shorter period of hyperglycaemia than has been suggested."

The figure shows the concentration of

Changes in serum concentrations of glycosylated HbA₁ with treatment in eight newly diagnosed diabetics.

HbA₁ (measured by Fluckiger's method¹ modified to read absolute values against fructose standards as mg fructose/g Hb) in eight newly diagnosed diabetic patients (five men and three women aged 49 to 74 years; two requiring treatment with insulin, three chlorpropamide, and three diet alone) before and at weekly intervals after treatment. Control of the diabetes was assessed by measuring the 24-hour urinary glucose excretion and the blood glucose concentration two hours postprandially. Like Fraser's group and others,² we found a significant fall in HbA₁ four weeks after starting treatment, by which time the 24-hour urinary glucose had fallen to an insignificant amount (less than 10 mmols) in all patients. Thereafter the concentration of HbA₁ continued to fall more slowly, to either within or just above the upper limit of the normal range (mean value for normal subjects ± 2 SDs = 1.9 ± 0.68) after approximately eight weeks' treatment.

Consideration of these monthly measurements alone, however, obscures much individual variation. Thus, although the concentration of HbA₁ fell in the first week of treatment in seven patients, it rose in one, and in three others it rose again two, four, and five weeks after starting treatment, despite a continuing overall fall in glycosuria and blood glucose concentration. We are confident that these fluctuations are real and not due to methodological errors, since duplicate estimations performed by column chromatography (Biorad Kits) showed similar changes. Moreover, other more detailed studies involving daily estimation of HbA₁ in inpatients (to be published elsewhere) confirm and extend these findings.

A P BROOKS
JOYCE BAIRD
ISOBEL M NAIRNMetabolic Unit,
Western General Hospital,
Edinburgh EH4 2XU¹ Fluckiger, R, and Winterhalter, K H, *FEBS Letter*, 1976, 71, 356.
² Ditzel, J, and Kjaergaard, J-J, *British Medical Journal*, 1978, 1, 741.

Handbook of Medical Ethics

SIR,—I welcome both the Central Ethical Committee of the BMA's initiative in producing a new draft edition of the *Handbook of Medical Ethics* and the resolution of the Representative Body to instruct the Council to review the handbook on an annual basis. The profession would do well, however, to pursue Dr D L Gullick's point (7 July, p 71) and ask the Council of the BMA to establish the purpose and the meaning of the document

for the benefit of its members and non-members alike.

To me, it would be acceptable for the handbook to be ascribed the status of a code of practice to which the BMA asks but does not oblige its members to conform and which its representatives, but not necessarily individual members, uphold in negotiations with other bodies. It would be unacceptable for the GMC or any other body to use the document as standard for disciplinary measures.

PATRICIA A BRADLEY

Greater Glasgow Health Board,
Glasgow G2 3HT

Half-way houses

SIR,—Under the heading of Any Questions? (5 May, p 1190) your correspondent refers to the use of half-way houses in geriatrics. The original concept of these was put forward by me in a public lecture during May 1949, and the idea was taken up by King Edward's Hospital Fund during the following year.

At that time many elderly so called chronic sick had been kept in hospital for very long periods, sometimes several years. When they were rehabilitated in geriatric units they proved difficult to discharge. Many of them were frightened of the world outside and very averse to leaving the wards which had been their refuge for such a long time. We needed some place where they could continue their physical and mental rehabilitation under skilled supervision, while the social workers sought a suitable destination for them. It was desirable to have a building away from hospital premises in which they could learn to live normal lives once more. Hence the scheme for a half-way house, under the umbrella of the parent geriatric unit and its staff, to train the old folk for independent existence.

In my original half-way house Westmoor (associated with St John's Hospital, Battersea) out of the first group of admissions, two patients had been in hospital for ten years, one five years, and another three years before transfer. Later we had a woman who had stayed 15 years as a patient. All of these were discharged in the end usually to part 3 accommodation. Eventually our annual turnover rose from 25 beds to a level of 200, thus freeing beds at the parent hospital at an unprecedented rate. The services of physiotherapists and occupational therapists enabled us to overcome the dread of discharge so common in our patients, while the co-operation of municipal welfare authorities facilitated transfer to suitable hostels or sheltered accommodation.

It is essential to use a half-way house for rehabilitation and convalescence only. If any patient fails to respond, he should be returned to the parent unit and replaced by another. Otherwise the place merely becomes a dump for the chronic sick. Diseases which have been dealt with at Westmoor include osteoarthritis, rheumatoid arthritis, various fractures, cerebral infarcts, ischaemic heart disease, varicose ulcers, chronic bronchitis, malnutrition, and mild grades of senile dementia, among others. So it will be seen that a half-way house is not a substitute for simple discharge of a fully recovered patient, but a specialised unit for those whose return to the community would otherwise be doubtful.

TREVOR H HOWELL

Queen's Hospital, Croydon