will fail as a result of a bureaucratic approach. This will leave them with a statutory duty to train GPs, but no one prepared to carry it out. There are many who have the ability and interest to train young doctors for general practice and who have high standards, but have enough common sense not to be bound by inflexible rules.

D P BROWN

Mid-Staffordshire Postgraduate
Medical Centre,
Staffordshire General Infirmary,
Stafford ST16 2PA

Doubtful epilepsy in childhood

SIR,—An important point not covered in your leading article (7 July, p 1) is the question of anticonvulsant therapy in doubtful cases of epilepsy.

Many clinicians face the dilemma of whether to institute a trial of anticonvulsants for a doubtful case of epilepsy. It is tempting to do this when the patient continues to return with unabated attacks. Although there are exceptions to every rule, it is probably best to advise against a trial of anticonvulsant therapy. Time tends to reveal the nature of attacks and I have never yet been convinced of any benefit to be derived from prescribing anticonvulsants on a trial basis when the diagnosis of epilepsy is uncertain. Having seen a number of adolescents who have had the diagnosis of epilepsy incorrectly applied and received anticonvulsants for several years, I would make a plea for avoiding this particular line of management.

MICHAEL SAUNDERS

Swainby, Northallerton,
North Yorkshire DL6 3DG

Drug-induced oesophageal injury

SIR,—May I add compound codeine, aspirin, and phenacetin tablets to the list of drugs causing oesophageal injury, as reviewed by Mr F J Collins and others (23 June, p 1673)?

A 54-year-old woman was admitted to hospital following a haematemesis. She had been well until the evening of admission, when she had suffered an acute headache for which she swallowed two compound codeine, aspirin, and phenacetin tablets. She took the tablets broken in half with a glass of water. She admitted subsequently that she felt the tablets stick in the upper gullet. Half an hour later she vomited, initially food but later blood. Physical examination was unremarkable, but early endoscopy revealed an annular irregularity of the oesophageal mucosa extending from 22 to 27 cm from the teeth. The lesion was not bleeding, but an adherent clot was noted. Biopsies from the edge of this lesion showed inflammatory changes only. The rest of the oesophagus, stomach, and duodenum were normal. She was treated conservatively and a repeat endoscopy 10 days later showed marked healing with epithelialisation but friable mucosa and easy contact bleeding. Two months later the lesion had completely healed.

J G WILLIAMS

RN Hospital,
Plymouth, Devon PL1 3JY

Photocoagulation and diabetic retinopathy

SIR,—We fully agree with Mr B L Hercules and Dr I I Gayed (23 June, p 1710) and Dr

G P Walsh (7 July, p 48) that treatment of proliferative diabetic retinopathy when new vessels arise from the optic disc is urgent. Treatment is easiest, least time consuming, and most successful when it is given in the early stages.

Dr Walsh's suggestion that patients with eye complaints should go to their GP with dilated pupils is not, however, a satisfactory solution. Firstly, how are the patients to know they will need to dilate their pupils, and who will give them dilating drops prior to attending surgery? Secondly, recognition of early disc vessels is by no means always easy. Doctors working in diabetic clinics should be trained to recognise sight-threatening lesions. Thirdly, and most important of all, patients with proliferative retinopathy have commonly normal vision and no symptoms until their first vitreous haemorrhage. By this time treatment is often difficult, time consuming, and not always successful.

Patients at risk of developing new vessels should be screened regularly by adequately trained people so that treatment can be early.

EVA M KOHNER
P LEAVER

Hammersmith Hospital,
London W12 0HS

Seat-belt Bill before Parliament

SIR,—Mr Neil Carmichael's private member's Bill that proposes to make the wearing of seat belts compulsory will be coming before the houses of Parliament in the near future. This is the single most practical measure to reduce the number of deaths and serious injuries on our roads. It has been estimated that the present death rate of about 8000 a year could be cut by about 1000.

We hope that all doctors will write to and lobby their MPs and do all they can to make sure that this Bill becomes law.

K C EASTON
Chairman
D C RAWLINS
Honorary secretary
H M BAIRD
Vice-chairman
R L HERBERT
Honorary treasurer

British Association of Immediate
Care Schemes,
London SW7 1PY

SIR,—May I make a plea through your columns to all doctors to write to their MP, with a copy to Mr Neil Carmichael MP, urging support for his Bill to make the wearing of car seat belts compulsory? I no doubt must be left about the views of the medical profession on this issue.

NEIL DAVIDSON

Leith Hospital,
Edinburgh EH6 6TH

¹ *Lancet*, 1979, **1**, 1415.

Diagnosis of allergy to insect stings

SIR,—In Any Questions (19 May, p 1334) it is stated that specific IgE tests (the radio allerge sorbent test (RAST)) are not performed in routine laboratories.

We have been supplying Phadebas RAST reagents to routine laboratories for five years and they are currently in use in at least 25

NHS hospital laboratories, one supraregional assay service laboratory, and various research labs which also perform routine service to some extent. In addition, we operate a routine testing service in collaboration with the RAST allergy unit at Benenden Chest Hospital in Kent.

RAST for both wasp and bee stings plus 60 other allergens is thus routinely available.

JOHN SPEIGHT

Pharmacia,
Hounslow, Middx TW3 1NE

Unfit for holiday

SIR,—May I use your columns to appeal, as a GP in a holiday resort, to my colleagues in the "donor" areas to take greater care in the advice they give their patients when asked, "Am I fit to go on holiday?"

In the past even days I have seen:

(1) A 37-week primigravida who presented in the second stage of labour giving a history of being discharged from a consultant obstetric unit three days before, having been an inpatient for 10 days with a diagnosis of hypertension.

(2) A 12-week-pregnant multipara with a history of two normal confinements and three spontaneous abortions who had been in a professorial obstetric unit as an inpatient because of bleeding in the present pregnancy, who presented with heavy blood loss and products of conception in the vagina six days after discharge from hospital.

(3) An 82-year-old man presenting in left ventricular failure who had no idea why he should have continued to take "the water tablets" even while on holiday.

All three told me that they had been specifically told by their doctors that they were fit to go on holiday, which involved a 90 mile journey, two travelling by private car and one by coach. I think further comment is unnecessary.

A J E POLLOCK

Skegness, Lincs PE25 2AQ

Disposable syringes for insulin injection

SIR,—The article by Dr A Greenough and others (2 June, p 1467) is of great importance in the management of diabetics in the developing countries, as most diabetics find it financially difficult to purchase glass syringes for insulin injections.

I have gone some steps further than the authors in that nearly all diabetics requiring insulin in the diabetic clinic of this hospital are taught to use the ordinary 2 ml or 2½ ml disposable syringes with 10 marks in 1 ml for their injections. These plastic syringes with needles, which cost 12 kobo (about 0.09p), besides being a lot cheaper than the British standard 1619 plastic syringes with needles, which cost 21 kobo (about 0.16p), are more readily available. The slight disadvantage is that finer adjustments in insulin doses may be difficult with the higher strengths of insulin. However, since we mainly use the U-40 insulins, a mark on the syringe is therefore equal to 4 units of insulin, and half the mark is 2 units. I am therefore not looking forward to any proposed plan to change all insulin strengths to U-100 insulins.

I have found that the diabetics use on average a plastic syringe a month, changing