printed name and number (for example, 40 or 80) on the pack and label, ignoring colour as an indication of either. It would be advisable to keep an empty pack or vial to check the identity of repeat prescriptions.

HARRY KEEN
Chairman, Medical
Advisory Committee

PETER WATKINS
Secretary, Medical
Advisory Committee

ARNOLD BLOOM
Chairman, Medical and
Scientific Section

CHRISTOPHER HARDWICK
Chairman, Executive
Council

British Diabetic Association,
London W1M 0BD

Photocoagulation and diabetic retinopathy

SIR,—I was fortunate enough to be invited to a Guy's lecture on photocoagulation and diabetic retinopathy and heard Professor H Keen—like Mr B L Hercules and Dr I I Gayed (23 June, p 1710)—refer to the urgency of the need for photocoagulation in proliferative eye disease.

I understand that most such cases come from eye rather than diabetic departments and that roughly 3% in eye departments will have the condition. This is a serious matter because, whereas appointments for diabetic clinics are more or less immediate, for the eye departments six or seven months is a standard waiting time unless urgency is recognised. This is about the time for all to be over and done with so far as saving the sight is concerned.

A possible solution could spring from the patient with an eye complaint placing homatropine in the eye before attending a GP's surgery. The recognition of the condition is not difficult.

G P WALSH

Blackburn, Lancs

X-ray examination of acute ankle injuries

SIR,—Drs Gerald de Lacey and Shirley Bradbrooke suggest that if there is no swelling over the malleoli it is not necessary to x-ray a twisted ankle (16 June, p 1597).

Even for excluding a major fracture their advice is not entirely foolproof, as their figures show. However, x-rays should be taken in such patients' ankles not merely to exclude fractures but also to exclude a torn lateral ligament of the ankle. It is the localisation of tenderness as well as the presence or absence of swelling which provides the solution to this problem. If there is neither swelling nor tenderness over the malleoli there will be no major fracture. And if there is tenderness over the lateral ligament, forced inversion pictures should be taken.

While the absence of swelling and tenderness excludes serious pathology, their presence does not prove its presence. But exclusion of the possibility is then impossible without x-rays.

WILLIAM RUTHERFORD

Accident and Emergency Department,
Royal Victoria Hospital,
Belfast BT12 6BA

SIR,—Drs Gerald de Lacey and Shirley Bradbrooke (16 June, p 1597) merit our thanks for

reminding us of the direct association between fractures round the ankle and soft tissue swelling. I have found this association useful for many years and indeed it was standard teaching at St Thomas's 25 years ago. Careful radiography will show the base of the fifth metatarsal in ankle films, and this is the only likely bony involvement with rotation strains in the ankle.

Incidentally, has anyone reviewed the damage suffered by patients we have failed to x-ray?

R W WILKINSON

Royal Berkshire Hospital,
Reading, Berks RG1 5AN

Injuries to boys who scramble

SIR,—I would like to thank Drs R H Jackson and A W Craft for their comments on this topic (16 June, p 1625) and for pointing out an apparent error in my computation. In the explanation of my figures I did not make clear that the figure of 24 races includes the initial five practice races, but these were not included in the rider races calculation and there was an average number of 19 races per event.

I can only agree with the observation that children of this age are not suited physically or mentally to take part in such sports. In many cases I feel that the parents are encouraging the child not for his enjoyment but for the feelings of displaced pleasure that they experience. Injury to their children does not diminish their enthusiasm, and I must say I was expecting a much greater frequency of serious injury when we embarked on the study.

The implication of Drs Jackson and Craft's letter is that a medical presence at such events not only condones them but positively encourages the competitions. Sadly, the experience in other areas is that events take place irrespective of the medical cover provided, since there are no safety rules by which they must be governed. This organisation attends many functions in the interest of offering help to the "sick, wounded, disabled, or suffering" and unfortunately our personal attitude to them is not a consideration; our first concern is for those in pain. We hope our continued presence at such events will help to reduce the suffering of those injured or unwell, and that in our small way we can influence the organisers to pursue their interest in the safest possible way. Scrambling is increasing in popularity nationally, and regulations on the conducting of meetings are urgently required.

M PLACE

St John Ambulance,
Gateshead Borough Division,
Gateshead, Tyne and Wear

MRC treatment trial for mild hypertension

SIR,—In a paper given on 13 June at the sixth scientific meeting of the International Society of Hypertension, the management committee of the Australian National Blood Pressure Study reported results which have led them to stop their randomised controlled trial for the treatment of mild hypertension. Some of your readers may want to know how this will affect the somewhat similar MRC trial in Britain.

The entry criteria in the two trials differ in an important way. The Australian trial included men and women aged 30-69 with

screening pressures within the 95-109 mm Hg diastolic (V) range. Randomisation for active or placebo tablets was determined after blood pressure measurements had been recorded at a subsequent entry examination, when all with pressures of 95 mm Hg or over were eligible. At that stage 16% had pressures of 110 or over, and these were included. The Medical Research Council's trial includes men and women aged 35-64 with diastolic (V) pressures found at screening to be within the 90-109 mm Hg range but only if their pressures are confirmed *within that range* at the entry examination.

In the Australian study pharmacological treatment conferred benefit only in those whose pressures at the entry examination were 100 mm Hg or greater; in this group treatment significantly reduced the incidence of fatal terminating events ($P < 0.05$) and of all terminating events ($P < 0.01$) compared with control subjects receiving placebo tablets. But among the 2056 subjects with pressures above 100 mm Hg at entry were 530 (25.8%) with diastolic pressures of 110 and over who would not have been eligible by the MRC trial criteria.

The Australian trial therefore deals with a higher blood pressure range than the MRC study, and it was within the upper part of the range that significant benefit from treatment was demonstrated. Nevertheless, on hearing of the Australian results the MRC trial monitoring committee examined the data in a similar way and submitted them to the chairman of the trial's ethical committee, who advised continuation of our trial, urging us to state publicly that "the data to hand are too few to justify any comment."

W S PEART

Chairman,
MRC Working Party on
Mild-to-Moderate Hypertension

Medical Unit,
St Mary's Hospital,
London W2 1NY

W E MIALI

Secretary,
MRC Working Party on
Mild-to-Moderate Hypertension

MRC/DHSS Epidemiology and
Medical Care Unit,
Northwick Park Hospital,
Harrow, Middx HA1 3UJ

Depot neuroleptics in a community mental health service

SIR,—I was most interested to read Dr Ernest Bennie's letter (7 April, p 958) commenting on his preliminary experience with cisclopenthixol decanoate (Clopixol). I would like also to give my impressions of the use of this drug in a catchment area of 85 000 people, as compared with the more familiar depot neuroleptics fluphenazine decanoate (Modectate) and flupenthixol decanoate (Depixol).

My experience of the use of depot neuroleptics began with the introduction of fluphenazine enanthate in 1964 and the subsequent development of fluphenazine decanoate in 1969. Fluphenazine was an excellent drug for controlling many of the symptoms of the major psychoses and was largely responsible for the increased numbers of patients who were able to live their lives outside psychiatric hospitals.

Other depot neuroleptics were introduced, but were less satisfactory for the sustained control of psychotic illness. Flupenthixol decanoate was the one exception, and in patients with simple schizophrenia who were

withdrawn it produced better results than did fluphenazine decanoate.

Before the introduction of clopenthixol in 1978 there were 400 patients with psychotic illness receiving either fluphenazine (80%) or flupenthixol (20%). Many of these patients had been treated for years and although they have, in most instances, remained moderately well with care and supportive treatment from the community mental health service some have complained of akinesia and lack of drive. Reduction of dosage did not lessen their feelings of tiredness. When clopenthixol was introduced in 1978 we decided to try it in schizophrenic patients who failed to respond satisfactorily to the other depot neuroleptics. Sixty such patients have been treated with clopenthixol and there has been a marked improvement in their alertness and rapport. Symptoms of agitation and aggression have been better controlled with clopenthixol than with either fluphenazine or flupenthixol. This was particularly obvious when the drug was used in the treatment of severely ill or moderately handicapped patients.

The dosage of clopenthixol used has ranged from 50 mg to 400 mg at intervals of two to four weeks. It is necessary to assess each patient carefully and start with a low dose as extrapyramidal symptoms and marked akathisia can be produced. Provided a start is made with a low dose, the side effects are minimal.

My impression after treating 90 patients with clopenthixol is that it is a most useful neuroleptic, particularly for the control of symptoms such as agitation and aggression that occur in the major psychotic illnesses. There is also an indication for its use in the hyperkinetic, destructive patient with mental handicap, in whom doses of 300 or 400 mg every two to three weeks have produced satisfactory control of symptoms. While I do not envisage clopenthixol supplanting fluphenazine and flupenthixol, I believe that psychiatric treatment will be enhanced by its use.

B BLAKE

St Dymphna's Psychiatric Hospital,
Carlow, Ireland

Psychiatric symptoms and hallucinogenic compounds

SIR,—I was interested to read the case report describing persisting panic attacks that developed after eating psilocybin mushrooms by Mr C Benjamin (19 May, p 1319). I have seen a number of such cases in which similar episodes of acute anxiety were experienced following the use of a variety of hallucinogenic compounds, including cannabis, mescaline, LSD, and psilocybin. However, it is rarely possible to be certain of the nature and purity of the substance they had taken.

These phenomena more frequently occur after a relatively heavy dose, after a period of heavy exposure, or after a "trip" that was particularly unpleasant. The anxiety which patients experience is much as described in the article—sudden episodes of acute anxiety coming out of the blue, usually associated with a variety of perceptual phenomena, including depersonalisation, derealisation, déjà vu, visual changes, etc; and in my opinion it is probably related to the "flashback" phenomenon. While it may at times be difficult to distinguish such post-drug anxiety symptoms from a severe social phobia, the episodes in question

tend to occur anywhere, and are usually not associated with specific phobic stimuli.

Because of the episodic nature of these experiences I had wondered whether they could be related to induced changes in temporal lobe functioning. EEG recordings in subjects complaining of such attacks showed a high frequency of minor temporal lobe abnormalities (for example, the presence of slow waves and occasional sharp waves). The abnormalities tended to disappear as the symptoms settled. The significance of these findings is not clear. The EEG recordings could not be compared with a prestate record and often attendance for follow-up was unreliable. The possibility that the symptoms represent temporal lobe dysrhythmic abnormality, however, is supported by the fact that the symptoms are generally alleviated by a minor tranquilliser while remaining unaffected or even exacerbated by a major tranquilliser, as the case report describes. The latter drugs, of course, are likely to aggravate an epileptic abnormality. The possible epileptogenic activity of these compounds has been raised in several reports but the issue is as yet unresolved.^{1 2 3} It has been my experience that the anxiety attacks generally settle slowly, provided that the patient avoids further contact with the relevant substances.

Academic Unit of Human
Psychopharmacology,
Medical Colleges of St Bartholomew's and
The London Hospitals,
German Hospital,
London E8 1DF

¹ Feeney, D M, *Journal of the American Medical Association*, 1976, **235**, 1105.

² Perez-Reyes, M, and Wingfield, M, *Journal of the American Medical Association*, 1974, **230**, 1635.

³ Mandell, A J, and Geyer, M A, *Comprehensive Textbook of Psychiatry*, ed A M Freedman, H I Kaplan, and B J Sadock, 2nd edn, p 445. Baltimore, Williams and Wilkins, 1975.

The dilated upper urinary tract

SIR,—Your leading article on this topic (26 May, p 1382) compresses a complex problem into a few succinct paragraphs. These comments are intended to clarify the discussion of our work.

Our starting point is a dilated renal pelvis shown on intravenous urography. This may be due to an effect of outflow obstruction on the pelvis—an obstructive uropathy—or it may be due to dilatation of an unobstructed floppy-walled pelvis. Our thesis is that if there is an obstruction sufficient to affect renal function—an obstructive nephropathy—then such obstruction is worth operating on to prevent deterioration of renal function, whatever the other reasons for operating, such as pain, infection, etc. Since the usual measurement of individual renal function¹ is not sensitive enough to be used for this purpose we measured the renal parenchymal transit time (PTT).² This is a more sensitive index of renal function in this context, because the first renal effect of an outflow obstruction is an increased intratubular pressure resulting in greater salt and water reabsorption^{3 4} and hence a prolongation of the intratubular transit time of a non-reabsorbable solute such as is used for renal radiotracer studies.

You ask whether this technique "will provide a reliable answer in all cases of equivocal obstruction." Our published experience is so far limited to pelviureteric

junction obstruction² and we distinguish two types of "equivocal obstruction": that affecting renal PTT and that not affecting renal PTT. Since, however, the response of a kidney to an outflow obstruction is a graded one, there will also be a small group not separable into one or other of these two categories. Serial studies are then used to show the trend of the renal response.

K E BRITTON
C NIMMON

Department of Nuclear Medicine
and Radioisotopes

H N WHITFIELD

Department of Surgery,
St Bartholomew's Hospital,
London EC1A 7BE

¹ Britton, K E, in *Dynamic Studies with Radioisotopes in Medicine*, vol 1, p 215. Vienna, International Atomic Energy Agency, 1975.

² Britton, K E, *et al*, *Lancet*, 1979, **1**, 905.

³ Rector, F C, Brunner, F P, and Seldin, D W, *Journal of Clinical Investigation*, 1966, **45**, 590.

⁴ Brenner, B M, Bennett, C M, and Berliner, R W, *Journal of Clinical Investigation*, 1968, **47**, 1358.

Treatment of tetanus

SIR,—Dr R S Edmondson and Mr M W Flowers (26 May, p 1401) generously referred to our results¹ with intrathecal antitetanus serum (ATS) from rural India as being remarkable but not confirmed by the results of Vakil *et al*² in Bombay. We would point out, however, that Vakil's treatment was not strictly comparable to that which we adopted.

In the first place, the Bombay team gave a relatively bulky dose of 250 U human tetanus immunoglobulin into the cisterna magna. Our study indicated that the best results are obtained from very small amounts of ATS intrathecally. With us, the intrathecal injection (either lumbar or cisternal but preferably the former) of 1 ml with 1500 U antitetanus (horse) serum resulted in a mortality of 11% (8/74), while with 0.15 ml or 200 U ATS (horse) the mortality was 4.5% (5/110); there was an overall mortality among all patients given intrathecal ATS of 7% as compared with a mortality of 14% in the control series ($P < 0.01$).

Secondly, Vakil *et al* did not give routine systemic betamethasone. Our experience³ strongly suggests that tetanus intoxication should be regarded as a polysystemic condition requiring polysystemic therapy. Such therapy should include active measures against tetanus toxin involvement of heart, lungs, adrenals, the parasympathetic nervous system, and gut absorption as well as muscles and the central nervous system. In such a polysystemic regimen we have found high-dosage parenteral betamethasone to play a very important part.

We would strongly endorse two among many of Dr Edmondson and Mr Flowers's conclusions from their series of 100 patients, of whom 10 died. Firstly, only small amounts of therapeutic ATS are needed wherever given. The question today is not how much ATS should be used but how little, when, and where (but never none). Secondly, competent nursing care is vital. The Leeds team insists that a trained nurse to care for each patient must be available at all times. However, dispensing with excellent but sophisticated equipment means that one does not need such highly specialised and technically competent nursing staff, although their devotion is essential. We too would credit much of the success in north India to the conscientious caring of the nursing staff, who for the most part were students and