CORRESPONDENCE

Management of self-poisoned patients J A T Dyer, MRCPsych; E J Salter, MRCPsych 205	Hypnosis G P Maher-Loughnan, DM; D Zimmerman, MB 208	Sodium in peritoneal dialysis solutions A M W Forbes, MRCP, and H J Goldsmith, FRCP 211
Vitamin D deficiency in Asian immigrants O G Brooke, MD, and others; T E Oppé, FRCP 206	Who should remove the endotracheal tube? A Boba, MD 209	Legal ignorance T J David, MRCP, and C H H Butcher, BA 211
Age and death in breast cancer J M Gowdy, MD, and J D Solomon, MD; D G Belasco, MB 206	Artificial ventilation and survival P J Tomlin, FFARCS; J F Nunn, FFARCS, and J S Milledge, FRCPED 209	Pressure in cuffed tubes T H S Burns, FFARCS 212
Costs of unnecessary tests D A Haslam, MRCP 207	A "specialty" for second-class doctors? W J Bassett, MRCP; R Parkes, MB 209	Emergencies and the laboratory A R Henderson, MRCPATH 212
Seat-belt legislation A W Fowler, FRCS 207	An open-access morbid pathology facility for GPs? P J Hoyte, MRCS; E W Walton, FRCPath 210	Sweet medicines and teeth J S Kroll 212
Diazepam and traffic accidents A A Landauer, PhD 207	Coronary artery spasm C Rosendorff, MRCP 210	Disposable syringes for insulin injection M J E Ely 212
Whooping cough after stopping pertussis immunisation D Jenkinson, MB; R K Ditchburn, MD 207	Saving asthmatics Elizabeth A Hills, FRCP; P F Meyer, MD 210	Direct debit payments J E Miller, MRCP 212
Mental handicap and the BBC Margaret M Sealey, FFARCS 208	Long-acting nitrates in the prophylaxis of angina pectoris D Weiler-Ravell, MD, and M M Bassan, MD 210	Functional budgeting J N Todd, MB, and P M Worsley, ACMA 212
Lung cancer and coal workers' pneumoconiosis A Seaton, FRCP; M Jacobsen, PhD 208	Drugs and breast-feeding B Moore, MRCP, and J Collier, MRCP 211	Future of clinical medical officers M Murchison, MFCM 213
	Cetiprin and oesophageal ulceration A D Lauder, MB 211	Community medicine C A Birt, MFCM 213
	Tory health R H Lloyd-Mostyn, MRCP 211	Inertia at the Elephant H B Wright, FRCS 213
		Haemochromatosis and the sicca syndrome Sir Richard Bayliss, FRCP 213

We may return unduly long letters to the author for shortening so that we can offer readers as wide a selection as possible. We receive so many letters each week that we have to omit some of them. Letters must be signed personally by all their authors. We cannot acknowledge their receipt unless a stamped addressed envelope or an international reply coupon is enclosed.

Management of self-poisoned patients

SIR,—Dr D R Blake and Dr M G Bramble (30 June, p 1763) seem to think that the DHSS should alter its policy¹ on the management of self-poisoned patients on the basis of their work and that of Gardner and others². But the issues involved are more complex than Blake and Bramble would suggest.

There are really two questions which require answering in this context and Blake and Bramble confuse them. These are, "Does it matter who does the psychiatric assessment?" and "Does it matter whether, or not the patient is admitted to hospital?" These issues are, of course, linked in that it is easier to provide the facilities for a thorough assessment if the patient is admitted, but the issues must also be viewed separately.

Gardner and others showed that, in the circumstances of their study, it did not seem to matter in terms of repetition of self-poisoning whether the initial assessment was done by psychiatrists or by specially trained medical teams. This referred to patients already admitted to medical wards. Blake and Bramble have now shown that there was fairly high agreement between what psychiatrists decided to do and what junior doctors said they would do—also with admitted patients. In particular, junior doctors correctly identified 81% of the cases in which psychiatrists recommended inpatient care. (It would be valuable though to

have details of the 19% "missed," as this is clearly a most important group that could contain cases of severe depressive illness with relatively high short-term suicide risk.)

From this information, Blake and Bramble see fit to argue that the admissions policy could be changed. If this is their argument, they should at least have had all their junior doctor assessments made by casualty officers in the working conditions of the casualty department. They ignore the possibility that admission in itself could have a beneficial effect on the patient. No reference is made to Kennedy's Edinburgh study,³ which showed that repetition in all cases of self-poisoning known to GPs was only a third as common in those cases admitted to the regional poisoning treatment centre as in those not admitted, and that in those not admitted subsequent referrals for psychiatric assessment made no difference to repetition rates. This was not a randomised prospective study (none seems to have been done) but selection factors were controlled as far as possible. The cautious conclusion was that something about the admission experience appeared to make repetition less likely. What the effective agent was remains open to investigation: perhaps it was hospitalisation itself, which makes the whole event more public and important, or the fact that the psychiatric assessment was very prompt.

Finally, I was struck by the high rate of psychiatric aftercare for the patients in Blake and Bramble's study—70% of admissions receiving psychiatric inpatient or outpatient recommendations. In Edinburgh, where the regional poisoning treatment centre has a special psychiatric service attached, only about a third of admissions are currently offered psychiatric aftercare.⁴ Is it really required for all the Newcastle patients who get it? And, if so, does not the high prevalence of psychiatric pathology in these patients argue in favour of the assessment being done by psychiatrists?

We need far more exploration of these and other issues before insisting on changes in DHSS policy.

JAMES A T DYER

MRC Unit for Epidemiological Studies in Psychiatry, Edinburgh EH10 5HF

¹ *Attempted Suicide*. Circular HM (61) 94. London, Ministry of Health, 1961.

² Gardner, R, et al, *British Medical Journal*, 1978, **2**, 1392.

³ Kennedy, P, *British Medical Journal*, 1972, **4**, 255.

⁴ Dyer, J A T, Duffy, J C, and Kreitman, N, *Parasuicide in Edinburgh*, 1977, unpublished report.

SIR,—The paper by Dr D R Blake and Dr M G Bramble (30 June, p 1763) gives interesting figures on the percentage agreement between decisions reached by psychiatrists and junior medical staff on the appropriate management for self-poisoned patients.

There is a growing feeling that the "fail-safe" policy of admitting to a medical unit all cases of self-poisoning is no longer feasible in an over-stretched, under-funded NHS, and this feeling is reflected in the authors' report that 32 out of 236 cases of self-poisoning studied were discharged from casualty without