definite spontaneous activity but poor relaxation and severe contraction fasciculation. Twitching was associated with a polyphasic unit of long duration firing about five times a second. Units were prolonged and polyphasic up to 3 mV. The interference pattern was reduced. We concluded that the quadriiceps weakness was neurogenic and that the pattern of electrophysiological abnormality was consistent with motor neurone disease.

Left radical pneumonectomy was performed with good clearance of lymph nodes from the mediastinum. The neoplasm was a large-cell carcinoma with a suspicion of adenocarcinomatous tubules in places. All fasciculation immediately resolved and pain in the knees disappeared.

Six months later there was much less wasting of quadriceps and no fasciculation. Right quadriceps and hamstrings were slightly weak but the left leg was normal. Reflexes in the legs were normal and plantar responses flexor. Clubbing was reduced, and the gynecomastia had gone. Electromyography of the right vastus medialis and rectus femoris showed no fibrillation or fasciculation and, though the interference pattern was mildly reduced, the motor units looked normal. Some polyphasic units were seen in the right tibialis anterior but they were not prolonged or increased in amplitude. There was no evidence of a lower motor neurone lesion.

Comment

Carcinomatous neuromyopathies occur in 16% of men with bronchial carcinoma.³ Brain *et al* reported on 11 patients with a syndrome indistinguishable from motor neurone disease—though running a more benign course—associated with malignancy.¹ Five had bronchial carcinoma, one of whom partially improved after resection. They concluded that motor neurone disease may be a manifestation of carcinomatous neuromyopathy. Our patient achieved complete remission, which we believe suggests a causal relation between the carcinoma and the neurological syndrome.

We thank Dr C J Earl and Dr F H Scadding for their help and advice in preparing this case report.

¹ Brain, Lord, Croft, P B, and Wilkinson, M, *Brain*, 1965, **88**, 479.

² Jokelainen, M, *Journal of the Neurological Sciences*, 1976, **29**, 55.

³ Croft, P B, and Wilkinson, M, *Lancet*, 1963, **1**, 184.

(Accepted 21 May 1979)

Department of Thoracic Medicine, Brompton Hospital, London SW3 6HP

D M MITCHELL, MB, MRCP, registrar

S A OLCZAK, MB, MRCP, senior house officer (present appointment: lecturer, department of medicine, Addenbrooke's Hospital, Cambridge CB2 2QQ)

Acute tubular necrosis after exposure to diesel oil

Exposure to diesel oil has been implicated in two previous reports of acute oliguric renal failure.^{1,2} Our case represents the first such report from Britain and provides further evidence that diesel oil should be considered a nephrotoxin.

Case report

A 47-year-old man presented with a month's history of epigastric and loin pains, thirst, nocturia, nausea, anorexia, and severe exhaustion, and a week's history of scrotal swelling. This unusual presentation prompted an inquiry into possible toxic exposure, and he admitted to cleaning his hands and arms with diesel oil over several weeks while building a home extension. On examination he looked pale and ill and had bilateral loin tenderness and scrotal oedema. Initial investigations showed: haemoglobin 11.9 g/dl, white cell count $6.7 \times 10^9/l$, erythrocyte sedimentation rate (ESR) 76 mm in the first hour, blood urea 6.5 mmol/l (39 mg/100 ml); plasma sodium 141 mmol(mEq)/l, plasma potassium 4.1 mmol(mEq)/l, plasma bicarbonate 26 mmol(mEq)/l, total protein 77 g/l, and serum albumin 43 g/l; urine analysis showed no protein, cells, casts, or growth.

Two weeks later he developed pitting ankle oedema and further investigation showed: haemoglobin 9.4 g/dl, reticulocytes 0.2%; blood urea 17.0 mmol/l (102 mg/100 ml), plasma sodium 139 mmol/l, plasma potassium 5.0 mmol/l, total protein 66 g/l, serum albumin 35 g/l, serum creatinine 910 μ mol/l (10.3 mg/100 ml); urine analysis normal, 24-hour urine protein 0.19 g, and creatinine clearance 14.2 ml/min. Intravenous urography

showed a faint nephrogram, normal-size kidneys, and no evidence of obstruction.

His urine output fell to 100 ml in 24 hours but he responded to frusemide with diuresis of 1500-2500 ml per 24 hours with disappearance of his oedema. His blood urea peaked at 28 mmol/l (169 mg/100 ml) during the diuretic phase. Renal biopsy showed the classic findings of nephrotoxic acute tubular necrosis—namely, patchy degeneration and necrosis of proximal and distal tubular epithelium with preservation of basement membranes.³ There were 14 normal glomeruli and some interstitial oedema. He subsequently made a good recovery with a haemoglobin concentration of 12.6 g/dl, ESR 27 mm in the first hour, blood urea 6.5 mmol/l (39 mg/100 ml), and serum creatinine only slightly raised at 140 μ mol/l (1.6 mg/100 ml) three months later.

Comment

The intermittent but frequent use of diesel oil before the onset of this patient's illness may explain the subacute development of renal failure. Nevertheless, the patient had histologically proved acute tubular necrosis. Detailed questioning revealed no history of exposure to any other nephrotoxin. Reidenberg *et al*¹ described a patient with acute oliguric renal failure following 10 days' exposure to diesel oil vapour in the cab of a lorry. Abdominal cramps and epigastric and loin tenderness, and oedema were prominent features, as in our patient. Strong evidence for absorption of diesel oil through the skin was offered by the patient described by Barrientos *et al*.² He developed acute oliguric failure on the same day as washing his hair with diesel oil, but renal biopsy—performed much earlier than in our patient, on the second oliguric day—showed only tubular dilatation with casts and a "light proliferation" of mesangial cells in the glomeruli. Considering the widespread use of diesel oil, toxic effects are extremely rare, although contact dermatitis is well recognised.⁴ Nevertheless, diesel oil by topical exposure or vapour inhalation should be accepted as a rare cause of acute renal failure.

Requests for reprints should be sent to Dr B I Hoffbrand.

¹ Reidenberg, M M, *et al*, *American Journal of the Medical Sciences*, 1964, **247**, 25.

² Barrientos, A, *et al*, *Archives of Internal Medicine*, 1977, **137**, 1217.

³ Jones, D B, in *Pathology*, ed W A D Anderson and J M Kissane, p 959. Philadelphia, Mosby, 1977.

⁴ Borrie, P, *Roxburgh's Common Skin Diseases*, p 169. London, Lewis, 1975.

(Accepted 26 May 1979)

Whittington Hospital, London N19

A J CRISP, MA, MRCP, medical registrar

A K BHALLA, MB, CHB, senior house officer

B I HOFFBRAND, DM, FRCP, consultant physician

Insulin oedema

Oedema occurring soon after starting insulin treatment is occasionally mentioned in textbooks on diabetes mellitus,¹ but we have found few descriptions of cases. We here report one such case.

Case report

A 31-year-old man was admitted as an emergency with a 5-day history of vomiting, diarrhoea, and increasing confusion. He had been fit, and there was no family history of diabetes. He was disorientated, dehydrated, and smelled of acetone. Urine tests showed large quantities of sugar and ketones, and his blood glucose concentration was 52.8 mmol/l (951 mg/100 ml). He was given intramuscular injections of 5 units of soluble insulin every hour and appropriate intravenous fluids. He responded well, and after eight hours his blood glucose concentration had fallen to 13 mmol/l (234 mg/100 ml). He was eventually stabilised on 50 units of Monotard insulin daily.

After seven days he developed ankle oedema, which became generalised over the next 24 hours and was particularly noticeable on the face; he resembled a patient with the nephrotic syndrome. His weight increased from 63 kg on the second day, when he had already been rehydrated, to 67 kg on the eighth day. His blood pressure and electrocardiograph were normal, and he felt well. Twenty-four-hour urinary protein output was 2.43 g/l and no erythrocytes or casts were seen. Serum total protein concentration was 60

g/l (6 g/100 ml) and serum albumin concentration 37 g/l (3.7 g/100 ml). Urinary fibrinogen degradation products were more than 10 µg/l (normal <2 µg/l). Creatinine clearance and serum urea and electrolyte concentrations were normal. The oedema persisted for a week but disappeared rapidly after treatment with bumetanide 1 mg daily. Four weeks later the diabetes was well controlled on 50 units Monotard insulin daily, the oedema had not returned despite stopping diuretics, and he had a normal urinary protein concentration.

Comment

The insulin oedema syndrome was first described by Leifer² and few other reports exist, although Shaper³ reported 17 cases presenting in Kampala between 1950 and 1961, representing 3.5% of registered diabetics for that period. In his patients oedema varied from mild facial swelling to anasarca. The patients were negroes and were said to have a considerable tendency to hypoglycaemia and recurrent oedema; they did not receive monocomponent insulin. Sudden retention of tissue fluid owing to glycogen deposition, accumulation of lactic and pyruvic acids in the tissues with a relative deficiency of thiamine, and increased glucocorticoid secretion were regarded as possible causes. Insulin may play an indirect part in regulating vascular permeability⁴ or may modify some of the inflammatory responses in delayed hypersensitivity.⁵ Some disorder of this nature would best fit the findings in our patient. The diabetes had started recently with oedema and proteinuria, although the latter was not severe enough to account for the considerable degree of oedema. A transient phase of increased vessel permeability seems the most likely explanation, although the picture was not that of angioneurotic oedema. We were puzzled by the rapid development of oedema. We have talked with colleagues who treat patients with diabetes and, although some were aware of the possibility of oedema, clearly it must be a rare phenomenon in Britain; its temporary nature is reassuring.

¹ Jackson, W P W, and Vinik, A L, *Diabetes Mellitus*. London, Edward Arnold, 1977.

² Leifer, A, *Journal of the American Medical Association*, 1928, **90**, 610.

³ Shaper, A G, *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 1966, **60**, 519.

⁴ Garcia-Leme, J, *et al*, *European Journal of Pharmacology*, 1974, **29**, 218.

⁵ Koltai, M, *et al*, *International Archives of Allergy and Applied Immunology*, 1975, **49**, 358.

(Accepted 16 May 1979)

St Chad's Hospital, Birmingham B16 9RQ

N R BLEACH, medical student

P J DUNN, MB, CHB, house physician

M E KHALAFALLA, MB, MRCP, registrar

B MCCONKEY, DM, FRCP, consultant physician

Acute myeloblastic leukaemia complicated by non-Hodgkin's lymphoma

Various types of leukaemia complicating Hodgkin's disease have been documented, though we could find only one report of a patient with acute myeloid leukaemia subsequently developing a lymphoma. This patient¹ was an 8-year-old girl with acute myeloid leukaemia treated by irradiation, who was found at necropsy to have Hodgkin's disease. We describe a patient with acute myeloid leukaemia who subsequently developed a non-Hodgkin's lymphoma.

Case report

A 69-year-old housewife presented in March 1977 with a short history of general malaise and lethargy. On examination she had generalised lymphadenopathy and hepatosplenomegaly. The peripheral blood smear contained 32% blast cells, and a bone-marrow aspirate 60% pleomorphic blast cells, establishing the diagnosis of acute myeloid leukaemia. She was treated with seven intermittent courses of cytarabine (cytosine arabinoside), 200 mg by intravenous infusion daily for three days, and doxorubicin, 60 mg intravenously on day 3. By July 1977 she was in complete remission with no lymphadenopathy or hepatosplenomegaly. She then received maintenance

treatment with cytarabine and doxorubicin and immunotherapy in the form of irradiated blast cells.²

She remained well until August 1978, when she complained of a two-week history of enlarged glands in the right groin and swelling of the right leg. She was found to have bilateral cervical, inguinal, and axillary lymphadenopathy, pitting oedema of the right leg extending from ankle to groin, and hepatosplenomegaly. No blast cells were seen in the peripheral blood, and a bone-marrow aspirate confirmed that she was still in remission. A bone-marrow trephine showed no tumour infiltration. Lymph-node biopsy was performed, and the histology was as follows. The node architecture was almost completely destroyed and replaced by a malignant lymphoma showing capsular infiltration. The lymphoma was composed largely of small lymphocytes of uniform size, which often showed angulated (cleaved) nuclei, best seen in a resin-embedded 1-µ section (fig). An extremely small proportion of less-well-differentiated lymphoid cells was present, as were several reactive histiocytes often containing nuclear debris. Occasional mitoses were seen. The tumour was regarded as a non-Hodgkin's lymphoma of small follicular-centre cells, classified as diffuse centrocytic (Kiel)³ or intermediate lymphocytic (WHO).⁴

Lymphoma showing lymphocytes with cleaved or angulated nuclei and occasional blast cells (arrow); 1-µ section, methacrylate embedded. Haematoxylin and phloxine × 400 (original magnification).

Comment

Various types of leukaemia have been described as a complication in Hodgkin's disease, and they are usually assumed to be related to cytotoxic chemotherapy or radiation, both of which are associated with leukaemia in patients with other diseases. In a review of 10 000 cases of Hodgkin's disease Crosby⁵ noted that the incidence of acute myeloid leukaemia was 10 times that in the general population. While the development of lymphomas is not associated specifically with cytotoxic chemotherapy, they sometimes occur in patients receiving long-term immunosuppression (for example, renal transplant recipients), and their development in a patient with acute myeloblastic leukaemia could be related to immunosuppression. Our patient was treated with doxorubicin and cytarabine, drugs not usually associated with powerful immunosuppression. In addition, these drugs were given intermittently with long drug-free periods between doses. This must raise the possibility of a direct link between acute myeloid leukaemia and non-Hodgkin's lymphoma. Any link between lymphomas and acute myeloid leukaemia might have escaped attention before because of the short survival of patients with these disorders. Owing to improved methods of treatment survival for both diseases is lengthening, and clinicians should be aware of a possible relation between them.

¹ Skworzoff, M A, *Frankfurter Zeitschrift für Pathologie*, 1930, **40**, 81.

² Powles, R H, *et al*, *British Journal of Cancer*, 1973, **28**, 365.

³ Gerard-Marchant, R, *et al*, *Lancet*, 1974, **2**, 406.

⁴ Mathé, G, *et al*, *WHO International Histological Classification of Tumours*, 1976, **14**, 37.

⁵ Crosby, W H, *Clinical Research*, 1969, **17**, 463.

(Accepted 8 June 1979)

University Hospital of Wales and Welsh National School of Medicine, Cardiff CF4 4XW

S WILLIAMS, MB, CHB, senior house officer, department of haematology

J A WHITTAKER, MD, MRCP, senior lecturer and consultant haematologist

J GOUGH, MD, MRCPATH, senior lecturer and consultant pathologist