its case. I do not think any of the letters that I have read following his original letter have tried to prove anything, but are merely suggesting that possibly it is worth considering as a way of helping people on their way to recovery.

Dr Kinnell also says that not everybody will use hypnosis with the same care and expertise as Dr Waxman (19 May, p 1354). It is precisely for this reason that the British Society of Medical and Dental Hypnosis has given training courses to numerous doctors and dentists in all parts of the country and does stress these factors, especially to the novice learning how to use it responsibly. In London, in the metropolitan branch, I personally have been involved in running seminars at regular intervals to which the novice can come and discuss any problems he may be having in the use of hypnosis in his practice; at the same time all those present learn something in the discussion which follows. Would Dr Kinnell suggest that because a surgeon amputates the wrong limb surgery should be condemned? However, this is what he is more or less suggesting with the use of hypnosis—that because some doctors use it irresponsibly the whole subject should be dismissed as a part of fringe medicine with no real value.

What I would like to suggest to Dr Kinnell is that he opens up his mind a bit more and starts running some of the trials which he condemns, to see whether there is more in the hypnotic technique than he so far likes to believe, and at the same time encourage other doctors in the Health Service to run such trials. Might I suggest, however, that it is no good having people who have been using hypnosis for a few months engaged in carrying out such trials, as it is only after many years of use that one is competent to use it most successfully.

D ZIMMERMAN

London WIN 1DA

¹ Sacerdote, P, *American Journal of Clinical Hypnosis*, 1966, 9, 100.

² Clawson, T A, jun, and Swade, R H, *American Journal of Clinical Hypnosis*, 1975, 17, 160.

³ Willard, R D, *American Journal of Clinical Hypnosis*, 1977, 19, 195.

⁴ Friedman, H, and Taub, H A, *American Journal of Clinical Hypnosis*, 1978, 20, 184.

⁵ Frankel, F H, and Zamansky, H S (editors), *Hypnosis at its Bicentennial: Selected Papers*. New York, Plenum Press, 1978.

⁶ Scott, D, *Proceedings of the British Society of Medical and Dental Hypnosis*, 1975, 1, 10.

⁷ Ewin, D M, in *Hypnosis at its Bicentennial: Selected Papers*, ed F H Frankel and H S Zamansky, p 155. New York, Plenum Press, 1978.

Who should remove the endotracheal tube?

SIR,—Dr B H Goodrich (21 April, p 1079) and Dr J M Cundy (5 May, p 1213) have written about avoiding vomiting and aspiration following extubation after anaesthesia.

I would like to submit that a great deal of sorrow and unhappiness could be avoided if the anaesthetist could accept the fact that no patient should be extubated—in the sense that the endotracheal tube should be left in place until the patient is awake and alert and has recovered his strength and co-ordination to such an extent that, on a verbal command such as “Reach up with your hand and pull that tube out,” he or she can then carry out that instruction. This has been my practice for years, and I have been quite satisfied that if a patient meets the above mentioned criteria he is most likely to be able to handle regurgitation.

This approach does not avoid the need for vigilance by the recovery room staff, nor does it preclude the use of the head-down or lateral position; but, unfortunately, it does require significant readjustment in thinking. Firstly, the anaesthetist should accept the fact that there are certain things that the patient can do better for himself; and, secondly, the recovery room staff should accept that their activities should encompass more than pleasant baby-sitting for the patient.

A BOBA

Department of Anesthesia,
Kern Medical Center,
University of California School of
Medicine at Los Angeles,
Bakersfield, California 93305, USA

Artificial ventilation and survival

SIR,—Dr J F Nunn and others in their recent article (9 June, p 1525) state that there is little information on mortality in wards after transfer from an intensive therapy unit. Perhaps their attention should be drawn to a report¹ of a series of 1719 consecutive patients admitted to an intensive therapy unit in whom well over 80% were ventilated for more than eight hours to give an in-unit mortality of 13.5% and a post-unit mortality of 7.1%. This differs from the 33% in-unit mortality and 30% (20 out of 67 discharges) post-unit mortality reported from Northwick Park. Their paper reports that of the 100 patients ventilated 20 died in the ward, yet their text says that this mortality was 17%. These figures are a little mystifying.

The fact of artificial ventilation of itself, however, is meaningless unless the criteria for artificial ventilation are stated. Thus in our unit the prophylactic use of artificial ventilation following major cardiac surgery has significantly reduced mortality and morbidity, similarly following reconstructive surgery of the lower aorta prophylactic use of artificial ventilation has reduced morbidity. Undoubtedly some of these patients would have survived without artificial ventilation but we have frequently regretted it when in individual cases we have attempted to predict that artificial ventilation would be unnecessary. A more recent development of our prophylactic use of artificial ventilation is following major neurosurgery with the objective of monitoring a lower P_{aCO_2} and so a lowered intracranial pressure. In such patients there is no question of respiratory failure. It is a matter of regret that the publication from Northwick Park does not indicate whether all the patients had established respiratory failure, which would otherwise have given a more meaningful interpretation to the follow-up data presented.

P J TOMLIN

Queen Elizabeth Hospital,
Birmingham B15 2TH

¹ Tomlin, P J, *Anaesthesia*, 1978, 33, 710.

* * * We sent a copy of this letter to the authors, whose reply is printed below.—ED, *BMJ*.

SIR,—We are grateful to Dr P J Tomlin for drawing our attention to his review of experience from the Queen Elizabeth Hospital intensive therapy unit in Birmingham. It is difficult to compare our experience with his, however, because:

(1) The majority of patients in the Birmingham unit were recovering from cardiothoracic surgery, for which the mortality is quite

different from general surgical and medical cases.

(2) It is not possible, from Dr Tomlin's paper, to derive mortality in patients who required ventilation as opposed to those who did not.

If we take only those patients in his paper characterised as general surgery and general medicine (some of whom were ventilated and some not) we find that the average mortality in patients in the intensive therapy unit for the four years of his study was 30% and 29% respectively. This compares with our unit mortality of 33% for ventilated cases and 16% for all cases, but obviously the statistics are very dependent on the type of patient admitted and the readiness with which ventilation and other measures are adopted, so that comparisons of this sort should be treated with caution. Our main interest was not so much the mortality in the unit, or even on the ward, but in the more long-term follow-up.

We are also grateful to Dr Tomlin for pointing out the error in our text in the discussion section, where the mortality in the ward following discharge from the intensive therapy unit should read not 17% but 20%, as is clear from the relevant table and text in the results section. We apologise for this error.

JOHN NUNN
J S MILLEDGE

Clinical Research Centre,
Harrow HA1 3UJ

A “specialty” for second-class doctors?

SIR,—Dr M P Roseveare's letter (30 June, p 1797) prompts a reply. While welcoming advances in general practice and in particular training requirements for practice, I am a little concerned that mature and experienced hospital doctors might be discouraged from entering general practice. The division of British medicine into narrow and protected bands of interest—call them what you will—can only frustrate many young doctors, and benefit little the health service. By coincidence, perhaps, the same issue carries a letter about “disillusioned registrars” (30 June, p 1798). Surely the answer to both letters lies in the expansion of a system of general practice that allows well-trained principals opportunities to exploit their special skills and expertise, as indeed some have done for over 10 years in Livingston.¹

Rigidity of thinking that bars the entrance to general practice of doctors from any source or that frustrates and wastes their skills is to be deplored. This in no way denies the need for vocational training but is simply a plea for an open-minded attitude to recruitment.

Dr Roseveare and Dr Taylor can be assured that general practice does, if there is the will and determination, offer scope for doctors to exercise their special skills at the same time as enjoying a wider experience of medicine. Certainly there is no need to think of being a second-class doctor.

WILLIAM J BASSETT

Craigshill Health Centre,
Livingston, West Lothian EH54 5DY

¹ *The Livingston Project—The First Five Years*. Scottish Health Services Studies No 29 (SHHD 1973).

SIR,—It cheered me to read Dr M P Roseveare's letter (30 June, p 1797) and I hasten to rally to him in his views about the