present. There should be some evidence of upper motor neurone damage such as large amplitude stable "F-responses," or the presence of "H-reflexes" in muscle groups where these are not usually seen. (2) Normal sensory action potentials. (3) Muscle biopsy demonstrating the features of denervation. (4) Biopsy of a sensory nerve (for example, sural nerve) which shows no abnormalities. (5) Normal biochemical and haematological findings—for example, normal B_{12} , folate, and creatinine phosphokinase values.

The second difficulty, as Mitchell and Olczak point out, is to show a definite association between treatment of the carcinoma and improvement in the neurological illness. In only three other cases in the literature was such improvement demonstrated.¹⁻³

Our patient is a 64-year-old man (we and V Gallai described this case to the May 1979 meeting of the Italian Society of Neuropathology). He presented in October 1978 with a history of cough, dyspnoea, and progressive shoulder girdle weakness. He was found to have weakness, wasting, fasciculation, and pathologically brisk reflexes in the upper limbs, with weakness and brisk reflexes in the lower limbs. Electromyographic studies of the affected muscles demonstrated lower motor neurone denervation, the surviving motor units being of long duration with amplitudes up to 8000 μ V. There were unusually prominent F-responses in the hand muscles and an H-reflex was recorded in (R) abductor hallucis. Both the sensory nerve fibre action potentials and the motor conduction velocities were normal. Biopsy of an affected muscle (right deltoid) showed only the changes of denervation. Biopsy of the right sural nerve gave normal results. Chest x-ray showed a mass in the lingula of the left lung, which on biopsy proved to be a squamous cell carcinoma of the bronchus. Subsequent treatment of the carcinoma with radiotherapy resulted in simultaneous shrinkage of the tumour and improvement in the neurological illness—an improvement which has so far been maintained.

The association of motor neurone disease with carcinoma is an important one to make, but we feel that the documentation must be very careful in order to exclude other conditions such as polymyositis, peripheral neuropathy, and myelopathy, which are already well documented as occasional complications of carcinoma.⁴

ANDREW PEACOCK
KEITH DAWKINS
GEOFFREY RUSHWORTH

Department of Neurology,
Churchill Hospital,
Headington, Oxford

¹ Lord Brain, Croft, P B, and Wilkinson, M, *Brain*, 1965, 88, 479.

² Stephens, T W, Rousgas, A, and Ghose, M K, *British Journal of Diseases of the Chest*, 1966, 60, 107.

³ Croft, P B, and Wilkinson, M, *Brain*, 1969, 92, 1.

⁴ Anderson, G, *Paramalignant Syndromes in Lung Cancer*. London, Heinemann Medical Books, 1973.

Psychiatric symptoms and hallucinogenic compounds

SIR,—I would like to reply to matters raised by Dr P Petch (9 June, p 1568) and Dr D Jacobs (7 July, p 49) following my report on persistent psychiatric symptoms after eating psilocybin mushrooms (19 May, p 1319).

Only a dozen or so of the numerous species of fungi native to the British Isles are hallucinogenic. The commonest of these are *Psilocybe semilanceata* and *Amanita muscaria*.¹ Hallucinogenic fungi can be divided into two groups according to the psychoactive drugs present in them. Fungi in the first group contain psilocin and psilocybin. Examples

include *Psilocybe semilanceata*, *P montana*, and *P coprophila*. Three other genera are included in this group—namely, *Panaeolus*, *Stropharia*, and *Hyphaloma* (*cyaneus*).

The distinguishing characteristics of *P semilanceata* are its cap, which has a diameter of between about 5 and 25 mm and is one and a half times as tall. It is of conical or hemispherical shape and it has a prominent nipple, and the margin of the cap is curved inwards. The stem is 2.5 mm thick, up to 76 mm tall, and cream coloured, paler than the cap. The edges of the gills are white, and are attached in an adnate-to-adnexed fashion with little spacing between them.

P semilanceata contains from 0.1 to 0.4% psilocybin by weight of the dried mushroom.¹ Hence 30-40 g of fresh or 5 g of dried mushroom contain from 3 to 12 mg of psilocybin. The dose of psilocybin required to produce hallucinations in humans is from 3 to 6 mg or 0.1 mg per kg body weight. In practical terms this is achieved by eating about 30 small mushrooms. I am satisfied that my patient described to me the features of *P semilanceata*, and that he ate a sufficient quantity to suffer ill effects. The effect of the mushroom is augmented by drinking alcohol. An electroencephalogram was done on this patient, but it proved to be normal. I am unable to add any comments to the hypothesis of Dr D Jacobs (7 July, p 49) that these symptoms represent temporal lobe dysrhythmic abnormality.

C BENJAMIN

University of Manchester Medical
School,
Manchester M13 9PT

¹ Cooper, R, *A Guide to British Psilocybin Mushrooms*, ed C Render, 2nd edn. London, Hassle Free Press, 1978.

Tuberculous paraplegia

SIR,—The old adage quoted in your leading article on tuberculous paraplegia (2 June, p 1442) might be rewritten to read "The treatment of tuberculosis changes with time and place." Certainly the treatment of Pott's paraplegia is changing now as many surgeons become disenchanted with the difficulties, complications, results, and expense of radical surgical procedures. Our experience is no exception.¹

Between 1965 and 1975, in the orthopaedic centre at Tunis, we treated 63 patients with Pott's paraplegia, 37 by bed rest and chemotherapy (streptomycin for three months, isoniazid and ethionamide or rifampicin or aminosalicylic acid for 18 months) and 26 by anterior decompression as well, the selection of treatment being arbitrary. We drew the following conclusions. The more proximal the vertebrae affected and the more marked the kyphosis, the greater was the incidence of paraplegia. Chemotherapy alone was more effective than combined medical and surgical treatment. Sixty-eight per cent of the medical group had complete neurological recovery but only 35% of the surgical group. Medical treatment was safer than surgery, having a mortality rate of 6% compared to 23%.

Even when paraplegia was complete, or had been present for more than a year before starting treatment, recovery was still possible. Complete neurological recovery was unlikely, however, if the patient had total paraplegia and anaesthesia with urinary retention. Patients in either group who had no neuro-

logical improvement within six months of the beginning of treatment were unlikely to recover.

JOHN V FOWLES

Care-Medico Afghanistan,
Kabul, Afghanistan.

¹ Sliman, N, *et al*, Les Paraplégies Pottiques. Congrès France-Italien des Groupes d'Etude de la Scoliose. Rome, 1977.

Indications for Keflex

SIR,—Once more out of my *BMJ* fell an advertising slip asking the question "How can you free the family from throat infections?" The answer to this apparently is supposed to be with Keflex, 500 mg twice daily. The text inside indicates that this is an inexpensive routine treatment for acute throat infections, by which it means streptococcal pharyngitis.

Treatment with penicillin is considerably cheaper and surely at least as effective. The only reason for not using penicillin for the treatment of streptococcal pharyngitis would be if the patient were known to be allergic to penicillin; and in this situation the cephalosporins would also be contraindicated because of the possible cross-sensitivity, and erythromycin would be the better choice. I think that the *BMJ* should not allow the inclusion of this advertising handout.

KENNETH HARDEN

Bearsden,
Glasgow G61 2DN

*.*We sent a copy of this letter to the manufacturers, whose reply is printed below.—Ed, *BMJ*.

SIR,—Thank you for the opportunity to comment on the letter from Dr K A Harden, who raises some important and interesting issues.

I do not accept that cephalosporins are contraindicated in patients who are allergic to penicillin. Probably no more than 10% of patients who are allergic to penicillin will be allergic to cephalosporins, and this may make them a very suitable choice in this particular group. With the much better blood levels achieved they are probably a better choice than erythromycin.

One of the main areas for concern in therapeutics is that of compliance, and it is likely that patients will be more compliant if they have fewer daily doses to remember. It could well be that Keflex would be particularly suitable if this were a major concern in an individual situation, as it can be used in a twice-daily dosage for throat infections.

Finally, on the question of cost, it is true that Keflex is considerably more expensive than the cheapest generic penicillin. However, most physicians prefer the medicines they prescribe to come from companies who have a consistent record of high quality products. In this situation Keflex is not a particularly expensive product.

BRIAN GENNERY

Lilly Industries Limited,
Basingstoke, Hants RG21 2XA

Gutter treatment for ingrowing toenails

SIR,—The authors of the article "Gutter treatment for ingrowing toenails" (21 July, p 168) are to be congratulated in achieving a 56% cure rate with a non-surgical method of