CORRESPONDENCE

Maternal phenylketonuria Isabel Smith, MRCP, and O H Wolff, FRCP... 127	"Diuretic escape" and "rebound oedema" L E Ramsay, MRCP; M K Chan, MRCP, and others... 131	Extradural haematoma: effect of delayed treatment A D Mendelow, FRCS, and F J Gillingham, FRCS... 134
Preventing postoperative thromboembolism V V Kakkar, FRCS; C R M Prentice, FRCP, and others; A J W Hilson, MRCP, and J G Ayres, MRCP; C A Wells, FRCS; R H Fell, FFARCS... 127	Prostaglandins and impotence P M S Rao, MB... 132	Tap water instead of electrode jelly for electrocardiographic recording O Dehlin, MD, and others... 134
Malaria prophylaxis Honor E Munro-Faure, BM, and D Donaldson, MRCPATH... 129	Mental handicap and the BBC R B Jones, MRCP... 132	Fifty years of penicillin A G Clarke... 134
Rubella vaccination S C Rowlands, MB... 129	Drugs for epilepsy J C Mucklow, and D F Smith, MB, MRCP... 132	Care of low-birth-weight babies M H Yardley, BM... 134
Value of cytology for detecting endometrial abnormalities Dulcie V Coleman, MD, and others... 130	Drug-induced oesophageal injury R G Hughes, FRCS; R E Cowan, MB, and others... 132	Review Body's award and consultant contract D E Bolt, FRCS... 135
Minor operations in general practice S M Lord, FRCSGLAS; J S Blackburne, FRCS 130	Sulphinpyrazone and warfarin after myocardial infarction J A Tulloch, FRCP, and T C K Marr, FRCP... 133	New consultant contract: the shelf or the bin? D N S Kerr, FRCP... 135
Saving asthmatics J W Paulley, FRCP... 130	Oral choline in cerebellar ataxia N J Legg, MRCP... 133	Emergency recall fees and basic salary increase N F Harley, FFARCS, and others... 135
Pathogenesis of pelvic inflammatory disease F M L Charnock, FRCS... 130	Prurigo and pets Susannah J Eykyn, MRCPATH... 133	A body to review not criticise? I B Cookson, MRCPSYCH... 135
What is to be done with the XYY fetus? Wendy Farrant, PHD, and Maj Hulten, MD; S R Wilkinson, MRCPSYCH... 131	Withdrawal of alclofenac (Prinalgin) R D Mann, MD... 133	Ethics, strikes, and the GMC M C Grayson, FRCSed... 135
Pneumoperitoneum associated with artificial ventilation J K Scott, FRCPed... 131	Bedside library for medical students D A P Cooke, FRCGP... 133	Manpower and GMC elections N G Graham, FRCS... 136
	Coronary care I W B Grant, FRCPed... 133	Pay-beds J M Shennan, FRCS... 136
	How to use an overhead projector T S Murray, MRCP... 134	Sir Thomas Browne: appeal for St Peter Mancroft, Norwich Reverend D Sharp, MA, and others... 136

We may return unduly long letters to the author for shortening so that we can offer readers as wide a selection as possible. We receive so many letters each week that we have to omit some of them. Letters must be signed personally by all their authors. We cannot acknowledge their receipt unless a stamped addressed envelope or an international reply coupon is enclosed.

Maternal phenylketonuria

SIR,—The paper by Dr G M Komrower and others (26 May, p 1383) based on their personal experience provides a valuable guide to the management of pregnancy in women with phenylketonuria; but, as the authors say, it is not yet possible to give confident advice and there is therefore need for a prospective study.

The steering committee of the MRC-DHSS phenylketonuria register, of which Dr Komrower is a member, has a valuable guide to the management of pregnancy in women with phenylketonuria; but, as the authors say, it is not yet possible to give confident advice and there is therefore need for a prospective study. The approach will be an epidemiological one, the aim being to follow up a comprehensive national sample of women (through contact with their medical advisers) in order to document any pregnancies and assess outcome in relation to management. Full details of the register procedures are being sent to clinicians already known to have female patients aged 14 and over under their care, and the register staff will gladly communicate with anyone else involved in the medical care of such patients.

The potential value of such a study can be illustrated by considering the birthweight figures of the Manchester and other treated cases quoted by the authors from the literature. The mean birthweight of 2781 g is below the mean of 3287 g for the UK population in 1970¹ and the distribution of the birthweights shows

a marked shift downwards, as has already been observed in untreated cases.² The infants in whom treatment was started during the first trimester had a higher mean birthweight (2960 g) than the infants in whom treatment was started later in pregnancy (2560 g). The finding of a higher incidence of congenital malformations in the earlier treated group (three out of six) than in the later treated group (one out of five) adds significance to the difference in birthweight between the two groups because of the recognised association between congenital malformations and low birthweight.

It is not surprising that dietary treatment started at some time after the first missed menstrual period may fail to prevent the gross malformations found in infants of mothers with phenylketonuria, because they have their origin very early in the pregnancy during the period of organogenesis. The first report of dietary treatment being started before conception has just been published by Nielsen *et al.*³ The pregnancy resulted in the birth of a healthy infant whose birthweight of 3500 g is greater than that recorded in any of the infants born to mothers in whom treatment was started after conception, and greater than that of any of the infants born to the mothers who received no treatment reported by Frankenburg *et al.*² These data taken together suggest that the greatest benefit will be

obtained by starting treatment prior to conception, but, failing that, the earlier in pregnancy the diet is started the greater the benefit to the fetus is likely to be. A larger sample of cases and consideration of other factors known to affect birthweight will provide further evidence in support of, or against, these speculations and should make it possible to give soundly based advice on the management of these pregnancies.

ISABEL SMITH
O H WOLFF

Institute of Child Health,
Guilford Street,
London WC1N 1EH

¹ *British Births 1970*, ed R Chamberlain, p 50. London, Heinemann Medical, 1975.

² Frankenburg, W K, *et al*, *Journal of Paediatrics*, 1968, **73**, 560.

³ Nielsen, K B, *et al*, *Lancet*, 1979, **1**, 1245.

Preventing postoperative thromboembolism

SIR,—I believe the conclusions of Professor E S Immelman and his colleagues of the Groote Schuur Hospital Thromboembolus Study Group (2 June, p 1447) are not supported by their data.

Perfusion lung scanning, is not specific for diagnosis of pulmonary embolism (PE). Forty-three patients developed new post-operative perfusion scan defects; in 28 of these, the score for PE was between 1/6 and 3/6. If one uses the criteria as defined by the authors (score 1/6 to 3/6 "PE possible but unlikely"), it is *unlikely* that 65% of scan abnormalities detected were caused by pulmonary emboli. It is amazing to read that these patients were still included in the

analysis. Furthermore, one can argue that even in 40% of patients the scan abnormalities considered by the authors to represent "definite" pulmonary emboli were due to some other pulmonary pathology since regular monitoring failed to demonstrate the presence of venous thrombosis—the precursor of pulmonary emboli. If all the patients with unlikely and probable PE are excluded, then one is dealing with only 6 and not 43 patients who developed so-called "non-fatal pulmonary embolism." Perfusion lung scanning alone to diagnose PE has now been abandoned in most institutions all over the world. However, with the more accurate combined perfusion-ventilation scanning 23.4% of patients in a control group developed mismatched ventilation perfusion scans compared to 7.8% of those who received 5000 units of heparin every 12 hours after major abdominal surgery.¹

The other conclusion of the Groote Schuur paper—that low-dose heparin does not reduce the incidence of proximal segment thrombosis—is also not valid owing to the small numbers of patients. The true extent of venous thrombosis can be determined only at necropsy. In the multicentre trial² thrombi were found at necropsy in the popliteal, femoral, or iliac veins in 18 patients in the control and three in the heparin group ($P < 0.001$). The authors also claim that low-dose heparin prophylaxis "fails to prevent significant thromboembolic complications" yet they provide no data at all to support such a claim. If one defines the significant thromboembolic complication to be one which requires treatment (rather than the perfusion defect), then there is no doubt that low-dose heparin is effective, as indicated by the results of the multi-centre trial²; 146 patients in the control and 41 in the heparin group required treatment for deep-vein thrombosis or pulmonary embolism or both, the difference being highly significant ($P < 0.005$).

Nor is it true that "It is unlikely that a more definitive multicentre trial with fatal PE as the end-point will be mounted." The findings of the multicentre trial² have now been confirmed by a large-scale Swiss-Scandinavian multicentre trial involving 4352 patients (see table),³ the detailed results of which will be published shortly.

During the last four years 35 clinical trials on low-dose heparin prophylaxis have been reported from a variety of patient populations. The results of these trials indicate that low-dose heparin is highly effective not only in reducing the incidence of significant thromboembolic complications but also in preventing deaths due to postoperative massive embolism. To call the perfusion scan defects, which may represent minor pulmonary emboli, non-fatal pulmonary embolism is not only inaccurate but, to say the least, misleading. Professor Immelman and his colleagues have suggested that the recommendation by the American Heart Association that "low-dose heparin

prophylaxis should be used in all haemostatically competent patients over the age of 40 who undergo abdominothoracic surgery" is premature. I am sure that the authorities concerned will be able to judge for themselves whether the data in their paper would justify withdrawing this recommendation. Equally disappointing were the comments of Professor J R A Mitchell (9 June, p 1523). Fourteen lives could possibly have been saved if low-dose heparin prophylaxis was used in 2076 patients undergoing major surgery in the control group.² Patients die from several other causes during the postoperative period, whose frequency is not modified by low-dose heparin. We would submit that it is totally unjustifiable to expect that a trial designed to assess the specific cause-related mortality should also provide an answer relating to total mortality. In many centres low-dose heparin prophylaxis is now being used as a routine prophylaxis for all high-risk patients who are likely to develop postoperative thromboembolic complications. It is the experience of such centres which will demonstrate in future whether one should continue the use of this form of prophylaxis.

V V KAKKAR

Thrombosis Research Unit,
King's College Hospital Medical
School,
London SE5 8RX

¹ Butterman, J, *et al*, *Medizinische Klinik*, 1977, **72**, 1624.

² International Multicentre Trial, *Lancet*, 1975, **2**, 45.

³ Saldeen, T, personal communication.

SIR,—We enjoyed the article by Mr E J Immelman and others (2 June, p 1447) and the comments by Professor J R A Mitchell (9 June, p 1523), but feel that the conclusions of Mr Immelman and colleagues, albeit preliminary, are not supported by their data.

These authors set out to study the natural history of postoperative deep-vein thrombosis (DVT) and its modification by low-dose heparin in a controlled trial. The numbers of patients intended to be enrolled in the trial, 200 in each group, were calculated so that if successful treatment halved the expected number of total DVT the trial result would have an 80% probability of detecting the difference at a 5% level. However, the authors have now given us a preliminary look at the data after only 100 patients in each group, so that the statistical pitfalls of having multiple looks at the data are already present.¹

The reason for the preliminary publication was the implication that iliofemoral DVT (shown in table III) and non-fatal pulmonary embolism (table IV) are not affected by heparin therapy. But the data do not warrant this conclusion. The end-point of iliofemoral DVT occurred in only 5% of control patients, and similarly "definite" pulmonary embolism detected by scanning was also present in 5% of controls. We have calculated that in order

to reduce an event occurring in 5% of control patients to half that number, with an 80% probability of detecting a difference at the 5% level of significance, at least 700 patients in each group would be required. We suggest that Mr Immelman and his colleagues might consider carrying out a large but valuable study of this type rather than proceeding with a second small study using a different type of heparin. Only when that number of patients has been studied can a confident statement be made that heparin does not affect major thrombotic end-points (as opposed to ¹²⁵I-fibrinogen-positive scans). The dangers of carrying out studies too small to ignore the beta-effect have recently been emphasised.^{2,3} Perhaps we should have a moratorium on the publication of preliminary results on the clinical assessment of antithrombotic agents, as only studies with an adequate number of patients will provide the statistically acceptable end-points to advance our knowledge.

We think that Mr Immelman and his colleagues are unduly pessimistic when they state that it is unlikely that a definitive multi-centre trial, with total mortality as the end-point, will be mounted. Surely this is exactly the trial which is required. The implication, in terms of the costs of drugs, labour, and possible haemorrhagic side-effects, is too great to be ignored. Large-scale clinical trials, though laborious and difficult to organise, are still the only means of reaching the correct conclusion in a situation where meaningful clinical end-points, such as total mortality, are few. It is unlikely that a single trial in this area, however well conducted, will provide a permanent answer to the question of which treatment, if any, is best for the prophylaxis of post-operative DVT. Only when the findings of a multicentre trial are independently confirmed will they be accepted into everyday clinical practice. Perhaps we should take a lesson from Wimbledon. It takes at least the best of three tennis sets to find the winner.

COLIN PRENTICE
GORDON LOWE
CHARLES D FORBES

University Department of Medicine,
Royal Infirmary,
Glasgow G4 0SF

¹ Federal Drug Administration, *Drug Bulletin*, 1978, **8**, 19.

² Freiman, J E, *et al*, *New England Journal of Medicine*, 1978, **299**, 690.

³ Rennie, D, *New England Journal of Medicine*, 1978, **299**, 828.

SIR,—We must take issue with the statement in the letter by Dr R S Stubbs (23 June, p 1707) that the work of Bell and Simon¹ calls into serious question the accuracy of diagnosing pulmonary embolism by means of perfusion lung scans.

Even allowing for the statement by Bell and Simon that "in this report, some of the difficulty in interpretation may have resulted from poor technical scan quality (p 705)", the correlation coefficients among the five lung scan panels ranged from 0.77 to 0.97,² compared with correlation coefficients among the three angiographic panels of 0.78 to 0.86. Furthermore, the lung scans in the urokinase pulmonary embolism trial were all performed during 1970-3, using in all cases a rectilinear scanner and in 91% of cases ¹³¹I-labelled macroaggregated albumin (MAA), and only two views were required.² Even in 1975, when Bell and Simon wrote their paper, this was not the optimum

Some results of two multicentre trials of heparin prophylaxis

	Multicentre Trial ² (1975)	Swiss-Scandinavian Trial ³ (1979)
Patients who received heparin prophylaxis ..	2045	2193
Patients who died	80	37
Autopsy rate	66%	86%
Fatal PE	2*	3
Contributory PE	3	3

*One of the 74 patients who received heparin but were excluded from the analysis because randomisation was not followed also died from pulmonary embolism.