some burden and the site of recurrent infarctions. Apart from the few cases of myelosclerosis with splenic haemopoiesis most big spleens are associated with many harmful effects of the "big spleen" syndrome notwithstanding the basic disease state. Provided that splenectomy is not delayed until it becomes a last-ditch procedure it is likely to improve the quality if not the quantity of life in a chronically ill patient.

There is also a very real place for splenectomy in the management of the patient with undiagnosed splenomegaly. Most large series of splenectomies contain a group of patients in whom the cause for the splenic enlargement was uncertain before operation. Usually histological examination of the excised spleen will reveal the nature of the basic disease but in some patients meticulous histological examination will still fail to provide a diagnosis, and long-term follow-up over a period of years may be necessary to detect the frequent development of haematological disease, usually lymphoma. With gross splenomegaly and an uncertain diagnosis many advantages may accrue from splenectomy, even considering the appreciable risks of the operation itself.

P J TOGHILL

Department of Medicine,
University Hospital,
Nottingham NG7 2UH

Suicide and attempted suicide

SIR,—The excellent article on "Suicide and attempted suicide" by Professor W H Trethowan (4 August, p 319) reminds us of the very great changes which have occurred not only in the law and the public's attitude to self-destruction but in the agents used in these forms of "communication between a person in distress and his environment," to use Professor Trethowan's own words. Few of these gestures, as he points out, prove fatal; but as the pharmaceutical scene has changed over the last 20 years so have the agents used in these gestures of despair.

We¹ reported in 1964 from St Stephen's Hospital, Fulham, on 126 patients (100 female, 26 male) admitted under our care into our general medical beds with suicidal gestures between 1955 and 1960. None died. One-third were considered straightforward and not suffering from psychiatric abnormalities; the other two-thirds were seen by the psychiatrist but only a small minority were transferred for psychiatric treatment and care. The agents most commonly used were barbiturates, as these were then the drugs most often present in the home drug cupboards; but now the more popular agents are the non-barbiturate sedatives and anxiolytics and the more popular analgesics. At the time of our study physical measures were very rarely used and gas fires in only 15%, mostly in the older age groups. Where such suicidal gestures were repeated the same agents, usually barbiturates, were often used again. M W Johns² has noted that hospital admissions due to acute barbiturate poisoning have decreased in England and Wales since 1965, roughly in parallel with barbiturate prescribing in the NHS, their place being taken by the less dangerous benzodiazepines and other agents.

It would seem that the most popular therapeutic agents of the day—that is, the ones in the family drug cupboard—will be the ones most likely to be used in these gestures

in the future as they have been in the past; but the fact that an agent, effective and non-toxic when used therapeutically in the correct dosage, can prove lethal when used for suicidal purposes in excessively high dosage, often with large amounts of alcohol, can hardly be set against it in terms of everyday therapeutics. What might be called pharmacological parasuicides, emotional gestures using whatever is in the home drug cupboard, occasionally prove fatal because of unanticipated side effects—for example, liver damage after ingestion of excessive amounts of paracetamol—but this risk will continue so long as drugs have side-effects and so long as the sad and distressed make these gestures of despair.

F DUDLEY HART

London W1

¹ Taylor, D J E, *et al*, *Practitioner*, 1964, **192**, 251.

² Johns, M W, *British Medical Journal*, 1977, **1**, 1128.

Wernicke's encephalopathy

SIR,—Your leading article (4 August, p 291) suggests that Wernicke's encephalopathy should be both treated early and ideally prevented. It is a rare complication of chronic alcoholism, only occurring in those patients who have a genetic abnormality of transketolase, which binds thiamine pyrophosphate less avidly than normal.¹ If all alcoholics were recognised and tested for this abnormal enzyme the development of the condition could be prevented in those vulnerable to it by regular supplements of thiamine, if necessary by the parenteral route, in much the same way as many schizophrenics receive depot neuroleptics.

You also suggest fortifying foodstuffs with thiamine. Vitamin depletion in alcoholics has been studied by Thomson,² who found many contributory mechanisms, including malnutrition, malabsorption, increased metabolic demands, and impaired utilisation. The only logical way round this would be to incorporate high doses of thiamine in alcoholic beverages before consumption, say 5 mg in a pint of beer, and 50 mg in a bottle of spirits. This could be done during manufacture or during inspection for duty purposes. After all, we have long accepted vitamin fortification of our breakfast cereal and margarine.

GARETH JONES

Whitchurch Hospital,
Cardiff CF4 7XB

¹ Blass, J P, and Gibson, G E, *New England Journal of Medicine*, 1977, **297**, 1367.

² Thomson, A D, *Clinics in Endocrinology and Metabolism*, 1978, **7**, 405.

SIR,—Your leading article (4 August, p 291) rightly draws attention to the frequency with which this eminently treatable but potentially fatal condition is missed. We are preparing a report of the results of a recent 24-month study of all patients referred to a single medical service screened for the presence of an "alcohol problem." In this period 73 such patients were identified, an incidence of 3%. Of these no fewer than nine had the Wernicke-Korsakoff syndrome, and none had been diagnosed before admission. Polyneuropathy was present in 10, tremor in 14, and cerebellar degeneration in three, with response to vitamin B complex in many instances.

A full clinical nutritional examination revealed virtually no other evidence of

deficiency. Liver involvement was frequent. A seven-day dietary recall showed a deficient thiamine intake in about 90% of these patients, a higher incidence of deficiency than for any other nutrient.

The close association with alcoholism, with its rising rate and especially high incidence in Scotland, gives cause for concern. Recognition by general practitioners of early thiamine deficiency in alcoholics might be life saving. You call for preventive measures—control of alcoholism and fortification of foodstuffs. The former is difficult to achieve but the latter in the form of increased fortification of bread, from which nearly one-third of our thiamine comes, would be simple, cheap, and safe.

D S McLAREN

Department of Physiology,
University of Edinburgh EH8 9AG

DANIEL H A BOYD

Leith Hospital,
Edinburgh EH6 6TH

Herpetic proctitis and sacral radiculomyelopathy in homosexual men

SIR,—Mr P L Samarasinghe and others (11 August, p 365) have drawn attention to the importance of herpetic proctitis in homosexual men. Severe anorectal herpes infections may be the commonest cause of acute urinary retention in young men. We in fact described two such cases in 1975.¹

The appearance of the rectal mucosa in the cases described by Mr Samarasinghe ranged from "hyperaemia and oedema to frank ulceration." *Chlamydia trachomatis* can cause exactly this picture of inflammatory rectal change² but, of course, not (to my knowledge) the associated neurological syndrome. In my series of homosexual men with herpetic proctitis, of which I hope to publish fuller details soon, ulceration was present on the anal mucosa up to the anorectal junction. Above this point—that is, on the rectal mucosa itself—no ulceration was present, but rather a hyperaemia that ranged from mild to severe, with areas of haemorrhage. Two asymptomatic excretors of virus (Mr Samarasinghe's "silent shedders") both had a normal anorectal mucosa as seen by the naked eye, but showed obvious hyperaemic change when the rectum was viewed and magnified through an operating microscope. The absence of ulceration on non-squamous-celled epithelium, however, is not a new observation: Coutts³ observed hyperaemia and haemorrhage only on the urethral mucosa in patients with herpetic urethritis as long ago as 1948.

DAVID GOLDMEIER

Department of Genitourinary Medicine,
University College Hospital,
London WC1E 6AU

¹ Goldmeier, D, Bateman, J R M, and Rodin, P, *British Medical Journal*, 1975, **1**, 425.

² Goldmeier, D, *A Study of Non-specific Proctitis in Homosexual Men*. MD thesis, University of London.

³ Coutts, W E, *British Journal of Venereal Diseases*, 1948, **28**, 205.

Relation between herpes simplex virus and carcinoma

SIR,—The paper on herpetic proctitis by Mr P L Samarasinghe and others (11 August, p 365), reporting urinary dysfunction as a complication, was of considerable interest and served also as a reminder of interrelation