In our survey² we found in the five types of schools where a cold buffet was offered that the average food per plate provided 10-25% less than the hot meal in the same school both for energy and for protein. At that time we were considering whether or not the target was being achieved and this greater shortfall with the cold buffet indicated that it was not getting as near the target as was the hot meal. However, since children vary so much in their needs perhaps, when offered a choice, they select what they need rather than what the nutritionist thinks that, on average, they ought to have.

On this basis there might be an actual saving in the cost of food if the children were offered a choice instead of the traditional approach. Certainly sandwiches, beans on toast, and similar snacks which are usually looked down on can be as nutritious as a traditional hot meal.

A E BENDER

Department of Food Science and Nutrition,
Queen Elizabeth College,
London W8 7AH

¹ Department of Education and Science, *Nutrition in Schools*, London, HMSO, 1975.

² Bender, A E, Harris, M C, and Getreuer, A, *British Medical Journal*, 1977, **1**, 757.

³ Bender, A E, Magee, P, and Nash, A H, *British Medical Journal*, 1972, **2**, 383.

⁴ Essex-Cater, A, and Robert-Sargeant, S, *Health and Social Service Journal*, 1975, **85**, 758.

⁵ Paul, A A, and Southgate, D A T, *McCance and Widdowson's The Composition of Foods*, 4th edn, p 266. London, HMSO, 1978.

Care of children in general practice

SIR,—Dr Stuart Carne's paper (21 July, p 190) is certainly an interesting account of one man's practice of child care in general practice. There are, however, some points that he raised that I would take issue with.

As I understand him, he criticises the hospital paediatric specialist for offering only episodic illness care. Yet, if I read Dr Carne correctly, by offering an illness care service on to which seems to be grafted a method of surveillance contriving to assess overall development as well as vision and hearing carried out on sick or recovering children, he seems guilty of offering just such a service, based as it is on the episodic illness care service of conventional general practice. I also concur with Dr E M Davies (18 August, p 443) in her doubts the wisdom of carrying out a surveillance examination on a sick or recovering child.

It also seems that Dr Carne is unclear about the real aims of offering a surveillance service for under-5s. Those whose hobby horse it is—and I'm proud to be numbered among them—never forget why such a service was set up in the first place in this country. In the developing world the lessons are there for all to see. When more than 80% of the under-5-year children and their parents in any country are making regular visits to the clinics of such a programme with a more than 80% immunisation rate combined with sound nutritional advice, an open-door service for illness care, and a developmental surveillance service, changes occur. The benefits are that mortality and morbidity decline, the growth of the individual child improves, and the demands on the curative services fall.

In this country today there are very few of us who can claim to have achieved these objectives. The general practitioner paediatrician as proposed in the Court Report is certainly an excellent starting point from

which to evolve such a nationwide programme. It is already a good sign that a few GPs are running such programmes and many more are actively exploring the possibility of doing so, despite the rejection of the concept by our leaders.

Finally, I would agree with the other correspondents that there is a need for further education. All three services that care for children—the community health services, the hospital services, and the general practitioner services—have a lot to learn from each other. Mutual respect is built on trust, and perhaps it is because we do not trust the quality of each other's skills that some of us believe that when consultants learn something about this aspect of medical care all will then be well: I disagree. My opinion is that we can and must all learn from each other. Then and only then can we start improving child care in this country.

GRAHAM CURTIS JENKINS

Ashford, Middx TW15 2TU

Referral of mothers and infants for intensive care

SIR,—In their excellent paper on the referral of mothers and infants for intensive care (18 August, p 414), Anthea M Blake and her colleagues seriously underestimate the extent of the antenatal referral service that would have to be provided to cause a reduction in regional perinatal mortality rates. In a region with 40 000 annual births, 120 lives would have to be saved to reduce the rate by three points. It is impossible to be sure of the outcome in a baby of a referred mother had the birth taken place in the district general hospital instead of the perinatal centre. It is fair to assume that babies who develop incipient respiratory failure would die if they were nursed in a district general hospital without facilities for mechanical ventilation. In the authors' analysis of babies of referred mothers, for each baby who required mechanical ventilation, a further two did not. Assuming an optimistic 80% survival rate for ventilated babies I calculate that the saving of 120 lives would entail the referral of at least 450 mothers annually (not allowing for multiple pregnancy). Yet the University College Hospital unit only received a quarter of this figure during a three-year period.

There is insufficient evidence that the availability of procedures other than mechanical ventilation (such as parenteral nutrition, continuous positive airways pressure, and continuous blood gas analysis) has such an ameliorating effect on perinatal mortality rates that transfer in utero is justified for all babies who might require such care. If that were so, many perinatal centres would be required for each region. We can also only speculate on the magnitude of any beneficial effect of neonatal care on the incidence of long-term handicap.

It is not difficult to predict the birth of a low-birthweight baby. It is harder to predict antenatally the occurrence of incipient respiratory failure in such a baby. Those pregnancies most likely to culminate that way are often the ones most likely to give rise to an obstetric emergency during transfer (for example, antepartum haemorrhage, severe pre-eclamptic toxæmia). It would be a retrograde step if we sacrificed maternal wellbeing in our efforts to secure neonatal health.

When comparing the neonatal mortality rate in babies of referred mothers with the national figures one should remember that the latter are likely to be over-represented by socially deprived mothers who received virtually no antenatal care. Before a system of antenatal referral widely develops I hope all the implications will be carefully debated—not the least being the effect on the education of midwives, obstetricians, and paediatricians working in district general hospitals. Surely our priorities ought to be: firstly, an improvement of neonatal staffing and facilities in district general hospitals; secondly, a network of neonatal flying squads to transfer babies with respiratory failure to a regional perinatal centre; and, thirdly, a very selective antenatal referral system for certain women in whom elective caesarian section is necessary before term and fetal lung maturation is uncertain or inadequate.

MALCOLM L CHISWICK

St Mary's Hospital,
Manchester M13 0JH

Breast or bottle

SIR,—The recent correspondence (25 August, p 492; 8 September, p 609) in which your readers gave a first-hand account of their success or failure in breast-feeding has a number of messages for those engaged in care of the newborn.

Although claims have been made in the past that successful breast-feeding may be established by virtually every mother I doubt whether this is true in our society in 1979. On the Isle of Wight we have recently studied all infants born during a 12-month period and are attempting to relate problems in the first year of life to the mode of feeding. Approximately 1000 infants were included in the study and most of the mothers kept a diary of problems they attributed to feeding. Reading these diaries has been a salutary experience, for, although 60% of Isle of Wight mothers are breast-feeding at the end of the first week after the birth of the infant and 33% after three months, such success is not easily achieved. The health visitors recorded "inadequate lactation" as the commonest cause for abandoning breast-feeding; and, although this may sometimes be the most convenient excuse for the mother giving up something she did not really wish to do, my study of the diaries convinces me this often is the true explanation.

I suspect that the pressures and demands of modern living, even on this peaceful island, make breast-feeding very difficult for many mothers. The Isle of Wight study confirms that breast-feeding protects against certain allergic diseases developing in the first year of life and the increased prevalence of infection in the bottle-fed baby has been well established.

The question arises whether, if breast-feeding is impossible, cows' milk is the best alternative. Perhaps we should consider the wider use of hypoallergenic feeds for the substantial numbers of mothers unable to breast-feed.

DAVID W HIDE

Shorwell,
Isle of Wight PO30 3JG

SIR,—I was amused by Dr Paula H Bolton-Maggs's account in Personal Paper (11 August, p 371) of her experiences in establishing

breast-feeding in an atmosphere of apparent disinterest or mismanagement, both by GPs and by hospital clinics despite the well-publicised advantages of breast-feeding. Some of her experiences seemed only too familiar.

Attitudes encouraging breast-feeding could be more positive. Continuity of care in the approach to breast-feeding with perhaps a personally supportive contact throughout and after pregnancy might go a long way towards correcting present deficiencies. Who better to do this than the health visitors? Success established by the National Childbirth Trust in its approach to breast-feeding may be through its personal sponsorship scheme whereby a recently lactating member of the trust adopts a local antenatal member and offers support in the perinatal and postnatal period. Unfortunately, the National Childbirth Trust has a rather middle-class appeal.

In this group practice responsibility for establishing contact lies with our practice-attached health visitors. All expectant mothers, whether booked for hospital or for GP antenatal care, are visited by them at home in the seventh month of pregnancy to discuss and, hopefully, influence the choice of infant feeding. Following delivery the same health visitor continues supportive care at home and in the practice's well-baby clinics. We recently analysed the feeding patterns of all 107 babies in the practice who arrived in the period August 1977 to August 1978 and whose mothers were given this support. Sixty per cent were breast-fed at birth, falling to 59% at six weeks and 36% at seven months. Seventy per cent of these babies were born into social classes III and IV. These results are not as depressing as those quoted by Dr Bolton-Maggs for a national survey. As was to be expected, bottle-feeding was the norm in social classes IV and V.

In an attempt to redress the balance in favour of the lower socioeconomic groups, our health visitors suggested more intensive domiciliary visiting to women in classes III-V. These mothers not only are less determined to succeed with breast-feeding but also have a lower attendance rate at antenatal classes. Encouragement to attend such classes could be given in the first trimester, perhaps even when the result of the pregnancy test is given, and some antenatal classes arranged during the first four months of pregnancy in the evenings, at a time when the NHS has little antenatal care to offer.

An integrated care scheme within a general practice setting can prove more supportive than the often impersonal approach of hospital clinics.

HELEN GRAHAM

London SE22

SIR,—Following Dr Paula H Bolton-Maggs's paper on breast-feeding (11 August, p 371) and Anne Chase's reply (25 August, p 492), I feel that I must redress the balance by describing my own experience of feeding my four-month-old firstborn son. At the age of 36 (Mrs Chase's midwife, please note!), not only did I have no problem establishing breast-feeding but I found it one of the most pleasurable and rewarding experiences of my life.

Much of the credit for this must go to St Mary's Hospital, Manchester, where mothers were actively encouraged to breast-feed. During the parentcraft classes not only did

we see films on breast-feeding, but a young mother also gave us a practical demonstration. Pregnant women wishing to feed were examined and counselled by a specialist midwife, and after the delivery were given free access to their babies; members of the nursing staff were only too willing to help with the first feeds. Problems of perineal soreness were overcome by the provision of a rubber air ring; but it was also possible to feed the baby lying on one's side, "face to face," a position that I still adopt for night feeds.

I do agree with Dr Bolton-Maggs that the positive aspects of breast-feeding should be more publicised—for instance, the delightful sensation of the baby sucking and his small hands on one's skin, accelerated weight loss (a fact I can confirm), and the convenience of carrying a food supply around with one. To those mothers who are really embarrassed about feeding in public to the extent of this being a deterrent I would suggest the purchase of a Kaneson breast pump¹—milk can be expressed at home and given to the baby at leisure. (The pump is designed like a syringe, the outer part doubling as a feeding bottle; and is very easy to use.) However, I have found that any loose blouse or smock enables one to feed quite discreetly; and nursing bras with a zip under each cup, as designed by Balance Maternities, enable one to manoeuvre without fumbling.

I hope this encourages would-be breast-feeders to take heart—it really can be a trouble-free and marvellous experience.

CAROLYN JONES

Manchester M21 2GQ

¹ Supplied by Kimal Scientific Products Ltd, Kimal House, Uxbridge Road, Hillingdon Heath, Uxbridge, Middx UB10 0PW.

*.*This correspondence is now closed.—ED, *BMJ*.

Fetal malnutrition—the price of upright posture?

SIR,—Professor John Dobbing (25 August, p 492), in his criticism of my paper (4 August, p 317), is perfectly right when he says that a longitudinal measurement cannot be assimilated to a curve of weight which is related to volume. However, instead of modifying biparietal diameter curves by a fourth-degree polynomial or considering a plot of necropsy data as he suggests, I think it is preferable to compare growth of the same parameter prenatally and postnatally.

In this respect measurements of head circumference are very appropriate and confirm my assertion that growth falters slightly before term. Campbell¹ showed that its velocity decreased from 11.5 mm per week at the beginning of the third trimester of pregnancy to 1.8 mm per week towards term, which is indisputably lower than the average of 4 mm per week deduced from the standards of head circumference between birth and 3 months.² This makes Professor Dobbing's statement that brain growth accelerates at the end of intrauterine life rather surprising, even though the relation between head circumference and brain weight is not linear. One becomes even more confused if by "accelerating" he means "increasing velocity" as generally accepted: in that case his assertion

is in conflict with the data of an article of his to which he referred.³

Professor Dobbing may, however, have other data supporting this new point of view. Before accepting them, I would like to have some reassurance: is he sure that his samples are not biased by an excess of newborn babies with relatively big heads towards term when mechanical problems of delivery become more complicated? Has he taken into account the opposite possibility, that he may have an excess of small-for-dates babies with a reduced head volume for shorter gestational period? Does he believe that necropsy data are more representative of what happens in utero to a normal fetus than ultrasound studies or even cross-sectional surveys of liveborn babies?

Finally, I wonder if Professor Dobbing really considers that his data prove that biparietal diameter is not an indicator of head growth. This could be established only by demonstrating that head volume and biparietal diameter can vary independently, which he has not done. At all events, this would imply that the head grows more like a cylinder of increasing diameter than like a sphere. Personally, I believe that this is not supported by everyday observation, and Professor Dobbing should agree on this point at least: his mathematical model of head growth is based on the assumption that its shape is roughly spherical.³

Can I ask him to solve the contradictions of his different interpretations of head growth before further questioning the conclusions of the ultrasonographers? This should help him to find the solution of his dilemma about fetal growth faltering in late pregnancy.

A BRIEND

ORSTOM,
Section Nutrition,
Dakar, Senegal

¹ Campbell, S, in *Fetal Physiology and Medicine*, ed R W Beard and P W Nathanielsz, p 271. London, W B Saunders, 1976.

² Vaughan, W C, and McKay, R J, *Nelson Textbook of Pediatrics*, 10th edn. London, W B Saunders, 1975.

³ Dobbing, J, and Sands, J, *Early Human Development*, 1978, 2, 81.

Hats for the newborn infant

SIR,—We were interested to read the paper by Dr Chaput de Saintonge and others (8 September, p 570) on the effect of Gamgee-lined hats in reducing heat loss in infants immediately after birth. Their results are the opposite of those that we obtained in a similar study of the stockinette hats which are more commonly used in this country.

We studied the effects of these hats and of overhead heating lamps in preventing a fall in body temperature in the first hour of life. We used a randomised study of factorial design with 20 full-term newborn infants allocated to one of four groups: hat and lamp, hat but no lamp, lamp but no hat and neither hat nor lamp. Initial (1-minute) rectal temperature and birth weight were similar in the different groups. The average fall in temperature in the first hour is given in the table. The stockinette hats did not reduce the fall in temperature, there being a small effect in the opposite direction. The effect of the overhead heating lamp in reducing cooling was, however, statistically significant ($P < 0.05$).

It seems therefore that the stockinette hats