diseases. Also, in the light of past investigations, I agree that all patients on psychotropic drugs should be warned that their driving might be adversely affected by their drugs and that this effect can be aggravated by alcohol. If all medical practitioners give this warning and all patients heed it, we are going to find it even more difficult to carry out the randomised control trial that is so badly needed.

F A WHITLOCK

University of Queensland,
Department of Psychiatry,
Royal Brisbane Hospital,
Brisbane, Australia

¹ Clayton, A B, *Human Factors*, 1976, **18**, 241.

Benzodiazepines and traffic accidents

SIR,—Dr A Landauer (21 July, p 207) raises several points relating to the role played by diazepam in road traffic accidents.

All 1,4-benzodiazepines have been shown to possess sedative activity, which can severely impair the regulation and performance of the sensory-motor tasks undertaken by patients during the course of their everyday behaviour. It is not just impairment of car driving ability attributable to diazepam which should give rise to concern but the effect of 1,4-benzodiazepines in general on the performance of the psychomotor tasks associated with work, home, and leisure pursuits.

Laboratory assessments of performance do relate to the real life situation if appropriate measures are used. Reaction time to visual stimuli has been shown¹ to be an appropriate analogue of real life performance where co-ordination of eye, hand, and brain is an important feature. The 1,4-benzodiazepine derivatives in general have been found to reduce the critical flicker fusion threshold, giving objective credence to the often-reported subjective drowsiness experienced by patients taking such drugs. We have been able to show consistent depression of critical flicker fusion thresholds and impairment of reaction time tasks following both acute and chronic administration of a range of 1,4-benzodiazepine derivatives.²⁻⁴

Laboratory information has been augmented by concurrently obtained data from tests of car driving ability administered independently by Advanced Driving Institutes. The impairment of performance shown in the laboratory tests following the administration of a 1,4-derivative, lorazepam, has been mirrored in the reduced performance on actual car driving tests of brake reaction, steering, width estimation, parking, and garaging undertaken by the same subjects in a placebo-controlled study.

Such findings lead us to believe⁵ that the administration of 1,4-benzodiazepines, not just diazepam, produces an increased risk of accident in situations where the integrity of the sensory and motor systems is an essential prerequisite for the safe performance of the task. The pathologically anxious or insomniac patient might be a better performer following treatment with a benzodiazepine but a consideration of the vast number of these derivatives prescribed would suggest that many non-pathological individuals are receiving these drugs and suffering performance decrements.

Others have concluded⁶ that the administration of a benzodiazepine derivative does not necessarily cause a death or road traffic

accident. However, this is missing the salient point of the recently published epidemiological survey by Dr D C G Skegg and others (7 April, p 917), which is that there is sufficient cause for concern that the administration of a 1,4-benzodiazepine derivative increases the risk of accident in any situation where safety is dependent on unimpaired psychomotor performance.

IAN HINDMARCH

University Department of Psychology,
Leeds LS2 9JT

¹ Linnoila, M, and Mattila, M J, *Pharmacopsychiatry*, 1973, **6**, 128.

² Hindmarch, I, *Arzneimittel-Forschung*, 1975, **25**, 1836.

³ Hindmarch, I, *Arzneimittel-Forschung*, 1976, **26**, 2113.

⁴ Hindmarch, I, *British Journal of Clinical Pharmacology*, 1979, **7**, suppl p 77.

⁵ Snaith, R P, *et al*, *British Medical Journal*, 1977, **2**, 263.

⁶ Finkle, B S, *Journal of the American Medical Association*, 1979, **242**, 429.

Rupture of the uterus during prostaglandin-induced abortion

SIR,—We read with interest the letter by Mr A I Traub and Mr J W K Ritchie (25 August, p 496), in which they quite rightly point out the hazards of combining oxytocin with prostaglandin in the induction of a mid-trimester abortion; and we agree with their advice to delay the infusion of intravenous oxytocin. We think that it should be stressed, however, that the patient whom we reported (7 July, p 51) ruptured her uterus following the insertion of intra-amniotic prostaglandin and hypertonic saline. At no stage was oxytocin used.

G J JARVIS
D A N JOHNSON
SIMON EMERY

Jessop Hospital for Women,
Sheffield S3 7RE

Induction of labour

SIR,—Following the article by Mr J H Shepherd and others (14 July, p 108) advocating the use of prostaglandin pessaries for induction of labour, we would like to illustrate the points raised in the letters of Mr A W Banks and Mr C J Hutchins (4 August, p 332) with two case histories and comment on our experience of the use of prostaglandin E₂ pessaries in a busy district general hospital.

The first case illustrates the dangers of unexpected delivery of an at-risk fetus in an inappropriate environment. A 24-year-old secundigravida (gestation confirmed by ultrasound scan at 17 weeks) was noted to be clinically small for dates from 27 weeks and so was monitored with plasma oestriols, serial ultrasound, and cardiotocography

(CTG). These were normal, but she developed mild pregnancy hypertension with proteinuria at 39 weeks. It was decided to induce labour, but as the Bishop score of the cervix was only 3, a 2-mg Prostin pessary was administered at 1300. She complained of irregular contractions soon after, and a CTG showed frequent but apparently small contractions. She did not require analgesia. She was judged not to be in labour, and monitoring was discontinued. At 1830 she developed the desire to bear down, and unexpectedly delivered on the antenatal ward. The baby weighed 2080 g but was fortunately in excellent condition, and needed no resuscitation.

The second case illustrates the real need for continuous CTG monitoring after use of prostaglandin pessaries in any at-risk pregnancy. A 27-year-old primigravida was clinically small for dates from 34 weeks. Plasma oestriol and human placental lactogen levels were always low, and from 37 weeks fell to below the 10th percentile. Antenatal CTG was normal, but because of the placental function tests, it was decided to induce labour at term. The Bishop score of the cervix was only 4, so a 2 mg prostaglandin pessary was inserted at 1120. Continuous CTG monitoring was performed (see figure). Because of large variable decelerations, which appeared over the next one-and-a-half hours despite keeping the patient on her left side, abnormal uterine action, and a cervix still only 3 cm dilated, it was decided to perform a caesarean section. A 3080-g infant was delivered in good condition.

We feel there are limitations to the use of prostaglandin pessaries for the routine induction of labour in a busy unit because: (a) the onset of labour cannot be predicted accurately; (b) patients may need cardiotocography for an indeterminate length of time; (c) when the cervix is favourable prostaglandins do not have any definite advantages over the tried and "midwife trusted" methods of amniotomy and oxytocin infusion; (d) they are expensive; (e) the potential advantage of mobility is frustrated by the need to monitor the patient and provide analgesia.

Prostaglandins do, however, have a valuable role in ripening the cervix and rendering it more favourable for induction by amniotomy and oxytocin infusion.

We are grateful to Messrs D Fairbairn and D P L May for permission to report on patients under their care.

M SUTTON
PHILIP STEER

Kingston Hospital,
Surrey

SIR,—I was pleased to see your leading article (18 August, p 407) stating that interest in methods of induction of labour is passing increasingly to non-invasive techniques which are more comfortable for the patients, as this statement is undoubtedly true. The simplest of these techniques is the intravaginal appli-

Cardiotocographic tracing in case 2.