CORRESPONDENCE

Treatment of biliary atresia A P Mowat, FRCP..... 605	Breast or bottle Isobel D Walker, MRCPATH; Hannah Gaston, FRCS; Janet F Gray, MRCP..... 609	Choline in a hereditary ataxia D V Philcox, FCP(SA), and B M Kies, FCP(SA) 613
Dangerous antihypertensive treatment D W Barritt, FRCP, and A J Marshall, MD.. 606	Enuresis G N Flaherty, FRACGP..... 610	Association between raw milk and human Salmonella dublin infection R G Small, FFCM, and J C M Sharp, MFCM 613
Rebound effect after stopping beta-blockers M J Lewis, MB, and others..... 606	Respiratory obstruction by epiglottitis J R Davies, FFARCS..... 610	Psychosocial stress in pregnancy J Dearlove, FRCP(C)..... 613
Short-term vocational medical work overseas A R Turner, FRCS, and Jean G C Turner, FRCS..... 606	Travel and health risks E J Trimmer, MB..... 610	Support for AETCEIITIP? D H Judson, FRCP..... 613
Mission hospital medicine P F Green, MB, and R K M Sanders, MD; E Ll Lloyd, FFARCS; P R Valentine, MB.... 607	Measles and vaccine protection L J Fish, MB, and P B Stones, FRCPATH.... 610	Costs of unnecessary tests G Sandler, FRCP..... 613
Clioquinol and related drugs L Berggren, MD, and others..... 607	Unusual manifestation of hypocalcaemia R J Schen, MB..... 610	Not an ethical dilemma J C Darnley, MRCS; J R B Cooper, MB 614
Metronidazole in the aetiology of pseudomembranous colitis M R B Keighley, FRCS, and D W Burdon, MRCPATH..... 607	Chronic ambulatory peritoneal dialysis D G Oreopoulos, FRCP(C)..... 610	Anatomists' and physiologists' pay D R Bowsher, MD..... 614
Iatrogenic collapse P K Schutte, MB; B F Hemphill, MB..... 608	Money and medicine R J Marcus, FRCS..... 611	Vocational training C A Stubbings, MRCP..... 614
Prevention of pulmonary embolism G R Osborn, FRCP..... 608	Doubtful epilepsy in childhood C E C Wells, FRCP..... 611	Hospital practitioner and clinical assistant grades J S Martin, FRCS..... 614
Domestic obstetrics W G Mills, FRCOG..... 608	Celtic and non-Celtic nostalgia J D Spillane, FRCP..... 611	Royal Commission report Moreen C Willis, MB..... 615
Induction of labour using prostaglandin E₂ pessaries J H Shepherd, FRCS..... 608	Reduction of catheter-associated sepsis in parenteral nutrition using heparin Margaret Wood, FFARCS, and A J J Wood, MRCP..... 611	Cuts in the health service D Adler, MRCS..... 615
Postpartum haemolytic uraemic syndrome J M Stark, FRCPATH..... 609	The Budd-Chiari syndrome G M Edington, FRCPATH..... 612	Private practice and nurse training K J Dennis, FRCOG..... 615
Neonatal care T H Hughes-Davies, FRCP..... 609	Medicine and scientific method R T Frankford, MB..... 612	Points Non-specific genital infection (E A Knappett); VAT on emergency equipment (I W W Fingland, Margaret J Fingland); Do- it-yourself waller for patients' notes (J R Scott); What's in a name? (A N Bamji)..... 615
	Rickets in Asian immigrants Juliet E Compston, MD..... 612	Correction: Umbilical hernia in child swimmers (Skidmore)..... 615
	Reversible renal failure during treatment with captopril P Collste, MD, and others..... 612	

We may return unduly long letters to the author for shortening so that we can offer readers as wide a selection as possible. We receive so many letters each week that we have to omit some of them. Letters must be signed personally by all their authors. We cannot acknowledge their receipt unless a stamped addressed envelope or an international reply coupon is enclosed.

Treatment of biliary atresia

SIR,—The recent leading article on the treatment of biliary atresia (30 June, p 1745) highlights the dramatic improvement in the prognosis of biliary atresia subsequent to the introduction of the operation portoenterostomy, and some of the outstanding problems. Dogmatic statements in the last paragraph of the article may militate against the best interests of infants with suspected biliary atresia and should be questioned. I agree that the current evidence suggests that surgical correction of biliary atresia should be performed as soon as the diagnosis is established. The rate of progression of hepatic disease in biliary atresia is very variable, however. The statement that the maximum age at which portoenterostomy can be carried out to give adequate bile drainage and arrest of the disease process is 60 days may deprive infants recognised late of the possible benefits of surgery.

Of 29 infants with biliary atresia seen in the last five years, portoenterostomy performed by my colleague Mr E R Howard has produced good bile drainage with normal serum bilirubin and stable or improving liver function

tests in 13 infants, all of whom are growing and developing normally. Operations by 12 weeks of age were successful in eight (47%), and after 12 weeks in five (41%). Although the period of follow-up is brief, currently we have no evidence to indicate that the prognosis for the latter group will ultimately be worse than for those operated on earlier.¹ Similar experience has been reported from France.²

I strongly disagree with the recommendation that liver biopsy (at laparotomy presumably) and operative cholangiography should be carried out at the beginning of the investigation of patients with suspected biliary atresia. Patients with biliary atresia present with features which are frequently indistinguishable, both clinically and using standard tests of liver function, from those with various forms of hepatitis.³ In south-east England such cases are three times as frequent as those of biliary atresia.⁴ They may frequently be distinguished from those requiring surgery without recourse to laparotomy.⁵

Of over 200 infants with hepatobiliary disorders admitted under my care in the last nine years, a correct preoperative diagnosis of

extrahepatic biliary atresia was made in 63, and of choledochal cyst in seven. One-hundred-and-eighty-six infants with hepatocellular disease were managed without laparotomy. An incorrect diagnosis of biliary atresia was made in two instances. In one, α -1-antitrypsin deficiency was erroneously excluded by the finding of an α -1-antitrypsin concentration within the normal range by an immunological technique. It was subsequently shown that this child had α -1-antitrypsin deficiency (phenotype ZZ) using an acid-starch gel electrophoretic technique.⁶ This mistake should be avoided in future. The second case was an infant with severe intrauterine growth retardation, marked failure to thrive, and complete cholestasis without striking changes in the hepatic parenchyma or in the portal tracts. At laparotomy this child was found to have an intact extrahepatic duct. Intractable convulsions developed within 24 hours and persisted until the child's death on the third postoperative day.

During the same period I have observed postoperatively a further seven infants with hepatitis who had been submitted to laparotomy by other clinicians. Three died within one to three months of surgery. In all who died the clinical condition had been stable before laparotomy but deteriorated following surgery. None had evidence of intrauterine infection, chromosomal or congenital abnormalities, or metabolic disorders such as α -1-antitrypsin deficiency which might have been expected to cause such early demise.

With cryptogenic disease it is impossible to be certain of the prognosis for these infants if they had not had surgery, but such deaths are exceptional and some component of the laparotomy or cholangiography may have contributed. These infants avoided a more immediate hazard of laparotomy—namely, in those with severe liver disease and markedly reduced bile flow, the removal of narrow but patent bile ducts because of a mistaken diagnosis of biliary atresia.⁷

Further, Thaler and Gellis⁸ have shown that infants with hepatitis submitted to laparotomy and operative cholangiography are more likely to develop cirrhosis than those with a condition of similar severity who are spared these investigations. I therefore submit that laparotomy and operative cholangiography are not simple procedures in infants with hepatobiliary disease and can safely be avoided in the vast majority. It would be a pity if the improved survival with a good quality of life for some infants with extrahepatic biliary atresia were to be counteracted by unnecessary morbidity and mortality from laparotomies in infants with hepatitis.

While the lure of "new and exotic liver function tests" may contribute to the delay in surgical treatment, more important factors appear to be the failure of parents, health visitors, and some family doctors to appreciate that hyperbilirubinaemia with dark urine is always pathological and requires investigation, and persisting and understandable doubts among some paediatricians about the long-term prognosis in biliary atresia even if satisfactory bile drainage is achieved following portoenterostomy.

ALEX MOWAT

Department of Child Health,
King's College Hospital Medical School,
London SE5 8RX

¹ Psacharopoulos, H T, and Howard, E R, submitted for publication.

² Odievre, M, *World Journal of Surgery*, 1978, **2**, 589.

³ Mowat, A P, Psacharopoulos, H T, and Williams, R, *Archives of Disease in Childhood*, 1976, **51**, 763.

⁴ Psacharopoulos, H T, and Mowat, A P, *Proceedings of an International Workshop on Neonatal Hepatitis and Biliary Atresia*, Washington DC, 1977, in press.

⁵ Howard, E R, and Mowat, A P, *Archives of Disease in Childhood*, 1977, **52**, 825.

⁶ Porter, C A, et al, *British Medical Journal*, 1972, **3**, 435.

⁷ Witzelben, C L, *Proceedings of an International Workshop on Neonatal Hepatitis and Biliary Atresia*, Washington DC, 1977, in press.

⁸ Thaler, M M, and Gellis, S S, *American Journal of Disease in Childhood*, 1968, **116**, 257.

Dangerous antihypertensive treatment

SIR,—We feel that a note of caution should be sounded about the introduction of combined tablets of beta-blocking agents with a diuretic in the treatment of hypertension. Both beta-blocking agents and diuretics given alone can be potent hypotensive agents. Oxprenolol and other similar agents may lower the systolic pressure by as much as 60 mm Hg¹ and thiazide diuretics are capable occasionally of producing similar large falls. It seems clear that when the two agents are used together their effect is simply additive.

Your recent leading article (28 July, p 228) on dangerous hypotensive therapy has correctly drawn attention to the dangers of lowering blood pressure too rapidly. One of the patients reported by Dr D H Cove *et al* (p 245) lost vision when treated by mouth with methyldopa, to which propranolol and later a diuretic were added. Their blood pressure chart shows the systolic pressure falling from

almost 300 mm Hg down to 150 mm Hg in 12 hours and to 100 mm Hg on the second day. It was at this time that the patient was found to be blind. Our own experience supports the conclusion that rapid reduction of raised blood pressure, particularly by the intravenous route, may be dangerous and we have certainly witnessed cerebral infarction occurring within a few days after beginning hypotensive therapy by mouth. This has led us to speculate over the relationship between pressure reduction and cerebral events.

In our view therefore it is correct to use either a beta-blocking agent or a diuretic in the first place and if the second agent needs to be added then it is eminently sensible to give the two agents in one tablet. This minimises the number of tablets to be taken. The tablets can be presented in calendar packs to aid the patient's compliance and the majority of hypertensives will probably not then need more than one tablet each day. Both beta-blocking agents and diuretics have a very flat dose-response curve^{2,3} and in most combined preparations the doses contained in the tablet are at the top of this curve. Apart from the risk of inducing complications from rapid lowering of blood pressure, the use of two agents when one might suffice doubles the chance of producing unwanted side effects. With beta-blockade either airways obstruction or cold extremities may be unnecessarily produced. If beta-blockade alone suffices to lower pressure then it is bad medicine to introduce the further risk of producing hypokalaemia, gout, or diabetes with a diuretic.

Promotion of the combined tablets should be preceded by the advice that either a beta-blocking agent or a diuretic should be used alone in the first place and the combination is only indicated if a second agent is really necessary.

D W BARRITT
A J MARSHALL

Bristol Royal Infirmary,
Bristol BS2 8HW

¹ Marshall, A J, Barritt, D W, and Harry, J D, *Postgraduate Medical Journal*, 1977, **53**, suppl No 3, p 168.

² Cranston, W I, et al, *Lancet*, 1963, **2**, 966.

³ Barritt, D W, and Marshall, A J, *British Heart Journal*, 1977, **39**, 825.

Rebound effect after stopping beta-blockers

SIR,—Dr T J B Maling and Professor C T Dollery (11 August, p 366), describing four patients followed for four days after stopping propranolol, observed no rebound increase in supine or erect blood pressure or heart rate, so adding more apparently reassuring evidence to counter the anecdotal alarms about the dangers of propranolol withdrawal. We feel we should draw attention to data^{1,2} which clearly show that a rebound increase in adrenergic activity does occur after stopping propranolol. Our more recent findings (unpublished) indicate that the phenomenon occurs also after treatment with other beta-blockers, particularly those that lack intrinsic sympathomimetic activity. As might be expected, the manifestations of this presumed increase in beta-receptor "sensitivity"³ become more pronounced under conditions of increased sympathetic drive.¹ We suggest that this is why the phenomenon has been missed in other formal studies of beta-blocker withdrawal and, equally but more importantly, why it may be dangerous when sympathetic

activity is high enough to override residual beta-blockade. Vasodilators, for example, will increase sympathetic activity, and the sympathetic stress of coronary artery surgery must be overwhelming. Catecholamines are a well-known experimental means of inducing myocardial necrosis indistinguishable from the changes of ischaemia.⁴

M J LEWIS
P J ROSS
A H HENDERSON

Departments of Pharmacology and
Cardiology,
Welsh National School of Medicine,
Cardiff CF4 4XN

¹ Ross, P J, et al, *Lancet*, 1979, **1**, 875.

² Nattel, S, et al, *Circulation*, 1979, **59**, 1158.

³ Glaubiger, G, and Lefkowitz, R J, *Biochemical and Biophysical Research Communications*, 1977, **78**, 720.

⁴ Rona, G, et al, *Archives of Pathology*, 1958, **67**, 443.

Short-term vocational medical work overseas

SIR,—Doctors who consider short-term work overseas (14 July, p 111) fall into two broad groups: those in hospital specialties, and those in general practice, public health, or the uncommitted stages of hospital practice. The major deterrent for most of the former is the near impossibility of regaining on return a place on any rung of their specialty ladder comparable to that which they left.

Most developing countries which ask for hospital doctors now require trained specialists; and, apart from the legal requirements of seniority, the unsupervised nature of the work and the extent and variety of clinical experience which many such posts present would place an intolerable burden on inexperienced shoulders. This means that most hospital doctors entering specialist work overseas will leave this country at consultant, senior registrar, or late registrar level. Consultants may be allowed prolonged leave of absence; senior registrar posts cannot usually be "held" for more than a year while the incumbent is abroad; and registrar posts must often be resigned completely.

Work overseas for a year or less at this level is of limited usefulness because of the problem of language and general familiarisation with the environment and the spectrum of disease. But those who resign their posts in these grades to serve for longer than a year face extreme difficulty in returning to their grades in the NHS, especially in major specialties, presumably because of doubts about the training content of largely unsupervised work.

The following suggestions might be considered in an attempt to resolve this impasse. (1) A small number of posts, decided on an individual basis, could be kept open for a longer period—say two years. (2) Each candidate should ensure, before signing any contract, that he acquires as much reliable information as possible about job content: newly created posts merit particularly close scrutiny. There should be much more collation of information from all potential overseas employers, together with "feedback" from past employees for circulation to postgraduate deans and others who may be called on to advise doctors—an "Overseas Yearbook" perhaps? (3) The establishment of a close link by the overseas doctor with a senior specialist in the UK in an advisory capacity would do much to maintain standards and morale. We envisage the advice being related to methods of collecting and recording clinical information and preparing material for possible publication; provision of references would also be an asset. (4) As a condition of retaining his UK post, the candidate should be required to produce regular and reasonably detailed reports on his work, perhaps to the joint higher training committee of his specialty. (5) It might be possible to arrange occasional inspection visits by, for example, UK