consultants if they were in that part of the world anyway. It amazed us to find how many "travelling experts" came within a few hundred miles of our remote mission hospital in Kurdistan. (6) For the few who for any reason (such as revolution) return after a longer period, the authorities might consider creating personal part-time supernumerary posts of limited duration. This would be infinitely preferable to a succession of locum posts, or prolonged period of unemployment, with no eligibility for state benefits because of previous absence from the UK, often on a salary too low to save capital or maintain UK National Insurance contributions.

Doctors who are more junior would be well advised to look for supervised posts—for example, in the universities of developing countries or in larger Christian hospitals with experienced senior staff, so that responsible reports of their work would be provided. At this stage a shorter period of a year or so could be useful to the overseas community.

Finally, those individuals who work in developing countries may have something worth while to offer the NHS on return. They value highly the privileges of working in our NHS. In addition to their extensive clinical experience, they have experience in decision making and responsibility under difficult circumstances which may give them a more mature viewpoint in consultant practice; there should be scope for those who have gained a broader experience of medicine from contact with different spectra of diseases and alternative schemes of health care. Since no "ideal" training for consultant practice has yet been devised or identified, these points provide a valid argument at least for considering such work for retrospective recognition by joint higher training committees.

Those who choose to work in developing countries are likely to have a high sense of vocation in their careers, and, although expecting no preferential treatment, they should not be penalised. They have in fact much to contribute to the NHS, which was after all conceived as a service to the community.

ALLAN R TURNER
JEAN TURNER

Limekilns,
Nr Dunfermline, Fife

Mission hospital medicine

SIR,—Following Dr Anne Savage's article (14 July, p 111) there have been a number of letters expressing interest in medical missions and the desire for an office of information of such opportunities abroad (4 August, p 335 and 11 August, p 388). Both the Medical Missionary Association and the Christian Medical Fellowship have for years provided such advice, the former through their quarterly magazine *Saving Health*, which includes an analysis of advertised needs of the many British missionary agencies. The CMF maintains individual correspondence and liaison with Christian medical associations of the Third World through which (and including missionary and government agencies) over 300 members of the CMF are currently serving. Also, Dr Peter Green of the MMA is the official BMA representative for British missionaries.

We would be happy to advise any who want to help overseas missions. Varied opportunities and needs abound. However, it should be noted that most mission hospitals request expatriates who are at least in sympathy with the ideals of serving in the name and spirit of Jesus Christ, the significance of which is not readily apparent in privileged Britain. We would gently, modestly, and in all sincerity plead for expatriates to "give up" (you won't,

you'll gain!) a few years or make an open-ended commitment to serve abroad, and to go not primarily seeking "interest" or "experience" but to understand the real needs of the situation and adapt oneself to meeting those needs. Besides, there is a growing resentment in the Third World against those whose contribution is so obviously geared to personal ambition or specialty. As one national superintendent put it, "Let them make such mistakes on their own people before they practise on us."

P F GREEN

Medical Missionary Association,
London N1 2NJ

R K M SANDERS

Christian Medical Fellowship,
London SE1 8XN

SIR,—My experience of mission hospital medicine was in a hospital specialising in eye surgery in the Bihar area of North India and my lasting impression is of the tremendous amount of high quality work that was done with the minimum of staff and equipment. The experience has also made me, like Dr C J Burns-Cox (11 August, p 388), critical of the waste and pomposity prevalent in the NHS and I am sure that a spell doing clinical medicine or surgery in a poor country would be of value for anybody intending to work in clinical medicine, though it might be of even greater value for potential community medicine specialists.

In contrast to the BTU (been to an underdeveloped area), the BTA (been to North America) has always been considered to be an advantage to a career in the NHS. Nevertheless, the fear of losing a place on the NHS career ladder meant that no senior registrar from my teaching hospital was interested in following me for a year as staff anaesthetist in a Toronto teaching hospital. Dr C J Chandler (p 389) should therefore not be too surprised at the lack of interest in the BTAf.

E LL LLOYD

Department of Anaesthetics,
Royal Infirmary,
Edinburgh, EH3 9YW

SIR,—The two recent letters by Drs C J Burns-Cox and C J Chandler (11 August, p 388) on mission hospital medicine reinforced the call by Dr Anne Savage (14 July, p 111) for a central register both to refill vacated Third-world posts and to find the returning doctor a rung on the ladder back home.

Neither mentions one other very important function of such a registry. It must have up-to-date information on the political and economic climate of the country involved. Although few doctors are likely to go to Uganda at the present time, there was a time when Makerere University was a hive of research for adventurous registrars, and let us hope it will be again. About three years ago I asked a BMA spokesman about the advisability of going to a particular Third-world country. Although it was a well-known area, he could neither tell me anything positive nor put me in touch with any doctor who had recently been there. All he could say was, "Australia and Canada are always a safe bet." Now, only three years later, both those countries are overdoctored.

It is a sad fact that there was never a time when Schweitzerian philosophy and medicine were more apt than now, and never a time

when they were less likely to be carried out. The "wind of change" has become a tornado, and let no doctor think he is immune. Such a registry therefore must have realistic information about the country or State to which it is recommending the enthusiastic doctor. If it does not, rather than graduate BTAf, he may join the even more distinguished band of NHS (never heard of since) doctors.

P R VALENTINE

Bristol

Clioquinol and related drugs

SIR,—All of us were foreign (that is, non-Japanese) participants in the International Conference Against Drug-Induced Sufferings, held in Kyoto 13-18 April 1979. We find the evidence overwhelming that medicines containing certain hydroxyquinoline compounds, such as clioquinol, can cause serious and irreversible injury to the nervous system. We urge those manufacturers who are still selling these products either to provide clear evidence that there are benefits which justify the risks or to withdraw them. In our opinion, the reassessment of these drugs is an urgent task for national drug regulatory authorities and international organisations.

LENNART BERGGREN (Sweden)	ÅKE LILJESTRAND (Sweden)
LARS ERIK BÖTTIGER (Sweden)	N D W LIONEL (Sri Lanka)
IWAN DARMANSJAH (Indonesia)	P K M LUNDE (Norway)
MAURICE N G DUKES (The Netherlands)	MIA LYDECKER (USA)
PAUL HANGARTNER (Switzerland)	M L MASHFORD (Australia)
OLLE HANSSON (Sweden)	MOHD ZAINI BIN ABDUL RAHMAN (Malaysia)
FREYA HERMANN (USA)	MILTON SILVERMAN (USA)
ANDREW HERXHEIMER (UK)	PECK CHUAN TEOH (Singapore)
TSU-PEI HUNG (Taiwan)	GIANNI TOGNONI (Italy)
SHIN HYUN-DUK (Korea)	MICHAEL TUVESON (Sweden)
KARL H KIMBEL (Federal Republic of Germany)	ALEXANDER M WALKER (USA)
BIRGITTA LANNEK (Sweden)	BARBRO WESTERHOLM (Sweden)
WIDUKIND LENZ (Federal Republic of Germany)	

Metronidazole in the aetiology of pseudomembranous colitis

SIR,—It has been suggested that metronidazole alone may be implicated in the aetiology of antibiotic-associated pseudomembranous colitis following the report by Dr A Kappas and others (18 March 1978, p 675) that a single patient developed pseudomembranous colitis after having metronidazole. As metronidazole alone has little influence on faecal microflora we questioned whether this patient received any other antibacterial agents. A detailed study of the notes revealed that the patient was given lincomycin in addition. The prescription sheet was missing from the notes and the information that this person received lincomycin became evident only as a result of careful scrutiny of the nursing records.

We apologise for this error and we would emphasise that the paper in question was concerned with the diagnosis of pseudomembranous colitis and not with a detailed investigation of its aetiology. We feel that it is most important to point out this error since we know of no patient reported so far in the