present infection rate is one episode every nine patient months.

Finally, I would like to add that one of the major advantages of this technique is that it is a dialysis treatment which does not require a machine. This has been appreciated especially by those patients who have been converted from intermittent peritoneal dialysis or hemodialysis to CAPD.

D G OREOPOULOS

Toronto Western Hospital,
Toronto, Ontario M5T 2S8

¹ Oreopoulos, D G, et al, *Transactions of the American Society for Artificial Internal Organs*, 1978, **24**, 484.

Money and medicine

SIR,—Professor Rudolf Klein's commentary (21 July, p 220) is an interesting contribution to the annual pay struggle that exercises successive ministers of health and the medical profession in spite of the interposition of the Review Body. While I would welcome his facts and not argue with them, his views adhere more to the "mediaeval theologians" approach to a "just wage" than to the climate of today.

Society has regrettably become besotted with unsocial hours, a concept not introduced by doctors or nurses. My wife works Sundays as a midwife on time-and-a-half, but the domestic working with her is on double pay. Does Professor Klein really justify this because my wife reputedly likes her work and the domestic is in servitude? Junior doctors work many unsocial hours but their overtime pay is well below their standard rate—a state of affairs surely incomprehensible to most union members. Consultants remain "on-call" as part of their total commitment, and recall fees, agreed on four years ago, remain unfunded. When the Royal Commission on Doctors' and Dentists' Remuneration laid down its guidelines in 1960 the distortions created by the unsocial hours philosophy hardly existed, and the comparison of the medical profession with other professions, working few, if any, unsocial hours, urgently needs revision.

We still live in a free society and doctors have to pay the going rate for other people's services and commodities. In the real world of private enterprise a doctor has to pay a minimum call-out fee of about £10 for somebody to fix his television, yet most general practitioners charge about £5 for a private visit because they consider this more advantageous than losing the patient to the NHS. Another cause of anguish to doctors must be their total inability to obtain the quality of service from others that society demands from them. If the skills displayed by garages had been perpetrated by the medical profession we should all have been sued into oblivion by now.

Professor Klein appears to favour financial enslavement of a profession because there is a glut of those wishing to enter it. I think many students would willingly have gone into the other sciences had there not been a world recession and no market for their skills. Architects have suffered during recent times, but senior accountants, actuaries, barristers, solicitors, architects, surveyors, and engineers can all delegate work and derive personal income from it and when business is good they prosper greatly. Surely the selection, training, responsibility, and skills required of the professions must be taken into account and gain preference over job satisfaction, which is subjective and nebulous.

If Professor Klein seeks to regulate medical pay by supply and demand then he must admit that UK doctors in the NHS today are paid well below the free enterprise rate (the same issue of the journal carries an advertisement for an anaesthetist in Holland offering £25 000 a year plus single accommodation).

Differentials in society at large are changing and if the present trend continues the country will get the mediocre doctors and university teachers it deserves and is willing to pay for. Professor Klein's philosophy did not do much good for industry under Labour, and the new tax cuts are an attempt to get quality and not quantity back instead of an overdraft, frustration, and the search for emigration.

RICHARD MARCUS

Stratford-on-Avon CV37 7BA

Doubtful epilepsy in childhood

SIR,—Your editorial of 7 July (p 1) was well named and, with few changes, would cover this difficult ground for patients of all ages. Neurologists, like paediatricians, would agree that the distinction "between epilepsy and non-epilepsy" is sometimes impossible. No matter: the best of all investigators, time, brings the inexorable answer. How strange that you lost patience at the very last sentence but one and surrendered your clinical sense to a machine: "Thus headaches, vertigo, and abdominal pain may occasionally remain unelucidated until an EEG is performed."

Even the references betray you. Swaiman and Frank¹ described six children with paroxysmal headache, none of whom had had any episode of impaired, let alone loss of, consciousness although four of the six had spikes or spikes and waves in their electroencephalograms. To argue that such an EEG diagnoses epilepsy in these children because a similar pattern occurs in the EEG of another child with undoubted epilepsy is to fall into the logician's trap of the fallacy of the undistributed middle.² Nor should epilepsy be diagnosed because their headaches went away when the doctor prescribed phenytoin. Does anyone diagnose epilepsy because trigeminal neuralgia, the tonic spasms of multiple sclerosis, or toothache are relieved by an anticonvulsant?

Eviatar and Eviatar³ found that half of a group of 50 children with bouts of vertigo were epileptic. This was not altogether unexpected: gyrate epilepsy is well known; some say that it caused Julius Caesar's falling sickness. Eleven of their patients had unquestionably major fits, attacks with vertiginous prodromata "culminating at times in a tonic or tonic-clonic generalised seizure." Fourteen others had giddy spells with headache and vomiting which might progress to "loss of postural control and loss of consciousness with or without hypertonicity." Despite the spike and wave patterns in their EEGs, despite even the minor twitchings and drooling and loss of speech sometimes seen, some doubt must remain that these were truly epileptic attacks and not just syncopal episodes after vertigo—of whatever cause—accompanied by the occasional myoclonic jerk.

Papathoeophilou et al⁴ studied 50 children with recurrent abdominal pain and re-examined the 14 respondents from an equally sized follow-up group in the hope of "clarifying some of the above conflicting findings in the literature." The low yield of cases with epilepsy

—one from each group and two others with photosensitive epilepsy—was a warning against twisting any human frailty into the tidy categories of limbic-lobe and complex partial epilepsy. This is what they wrote: "It seems to us that only in a very small proportion of children is recurrent abdominal pain or 'periodic syndrome' an epileptic phenomenon and that the finding of 'abnormalities' in the EEG, other than spikes or spike and wave, should not be taken as evidence of epilepsy." Please note also that they call spikes and spike and wave evidence of epilepsy and do not equate the electrical sign with diagnosis.

CHARLES WELLS

University Hospital of Wales,
Cardiff CF4 4XW

¹ Swaiman, K F, and Frank, Y, *Developmental Medicine and Child Neurology*, 1978, **20**, 580.

² Fowler, H W, *A Dictionary of Modern English Usage*, 2nd edn, revised E Gowers, p 662. Oxford, Clarendon Press, 1965.

³ Eviatar, L, and Eviatar, A, *Pediatrics*, 1977, **59**, 833.

⁴ Papathoeophilou, R, Jeavons, P M, and Disney, M E, *Developmental Medicine and Child Neurology*, 1972, **14**, 31.

* * * To elucidate means "to throw light on." To say that the EEG may occasionally elucidate a clinical problem is not the same as saying that it proves a diagnosis of epilepsy. Dr Wells appears to have ignored the last sentence of the leading article: "Nevertheless the mainstay of diagnosis in the paroxysmal disorders remains the doctor's ability to take and analyse the patient's history and assess his symptoms in the context of his psychological make-up." We believe as firmly as does Dr Wells that the EEG, like all laboratory investigations, must be assessed critically by the clinician.—ED, *BMJ*.

Celtic and non-Celtic nostalgia

SIR,—Dr Dewi Davies's charming piece on nostalgia (18 August, p 433) includes mention of *The New Oxford Book of Light Verse*. Its editor, Kingsley Amis, chose a particularly good sample of Celtic nostalgia in a verse by Wynford Vaughan-Thomas, now with us here in Pembrokehire. It is called "*Hiraeth in N.W.3*" (*hiraeth* is Welsh for nostalgia). It begins:

"The sight of the English is getting me down,
Fly westward my heart from this festering town . . ."

and ends:

"Glorious welcome that's waiting for me,
Hymns on the harmonium and Welsh cakes for tea,
A lecture on Marx; his importance today,
All the raptures of love with a Bangor B.A."

When Dr Davies does come home to these parts, he will find that my next-door neighbour grew up with him on the next farm—also having spent a lifetime in south-east England.

So—Croeso.

J SPILLANE

Newport, Dyfed SA42 0NR

Reduction of catheter-associated sepsis in parenteral nutrition using heparin

SIR,—We read Mr M J Bailey's study (23 June, p 1671) with interest and were impressed by the reduction in the incidence of infection. However, we felt it important to draw attention to an effect of heparin, even at low dose, which he does not discuss. We have shown that through its ability to increase lipoprotein