CORRESPONDENCE

Too few necropsies J R Anderson, FRCPATH..... 932	The grumbling appendix D M A Francis, FRCS..... 936	University examinations and the medical student D S McLaren, MD..... 938
Reactions to whooping cough vaccine P E Brown, FRCPED; J B P Stephenson, FRCP 932	Susceptibility to primary biliary cirrhosis F Arriaga Chapper, MD, and others..... 936	Practical problems with the Clinitest B A Elliott, FRCPATH..... 938
Rubella vaccination M Gringras, MB..... 933	Serum bilirubin and hepatic enzyme induction R P H Thompson, FRCP..... 936	Choline chloride in the treatment of ataxia I R Livingstone, MB, and F L Mastaglia, FRCP..... 939
Circumcision and cruelty to children J E Oliver, MRCPsych..... 933	Spontaneous recovery from rapidly progressive glomerulonephritis D J Evans, MB..... 936	Pindolol and pulmonary fibrosis P Krupp, MD, and J M Crawford, MB..... 939
Risks of amniocentesis Helen M E Bantock, MRCP, and I Sutherland, DPHIL..... 933	Kidney transplants and long-term immunosuppression H Graf, MD, and others..... 936	Wanted: a new wound dressing R Bewick, FRCS..... 939
Induction of labour J Clinch, MRCOG..... 934	On-demand analgesia equipment W D White, FFARCS, and others..... 937	Royal Medical Benevolent Fund Christmas appeal 1979 Sir Thomas Holmes Sellors, FRCS..... 939
Fetal malnutrition—the price of upright posture? J Dobbing, FRCPATH..... 934	Unusual haemolytic-uraemic syndrome E N Wardle, MD..... 937	Review of composition of the General Optical Council N E Leigh..... 939
Serum free thyroxine in pregnancy O Djahanbakhch, MRCOG..... 934	Marital pathology and the choice of partner P R V Tomson, FRCGP..... 937	Unblocking beds L A Wilson, FRCP..... 939
Psychosocial stress in pregnancy F H Creed, MRCPsych..... 934	Mechanical aids to ventilation for use in the field R Greenbaum, FFARCS..... 937	Cost of treatment in the NHS J C Smith, FRCS..... 940
Samaritans versus suicide G H Day, MD..... 935	Diagnostic imaging R H Corbett, MB..... 937	Health centres apparent Lord Taylor, FRCP..... 940
High- and low-carbohydrate diets J I Mann, DM, and J Eaton, SRD..... 935	Cubital tunnel syndrome S Renfrew, FRCP..... 937	Points Prediction of fetal sex: a cautionary tale (J A Nunez); Familial appendicitis (M Segal); Easy-rider sling (Tessy K Hanid); Remedy for excessive salivation (Roberta Butler); Photographing slides during scientific presentations (R T Jung); Art and audacity (H G Easton)..... 940
Cancer of the ovary M V Williams, MRCP, and C L Harmer, FRCR; M H Wilkins, FRCSED..... 935	Recommended terminology of urinary tract infection N B Eastwood, FRCGP; H G Hanley, FRCS.. 938	
Soft-tissue sarcomas S Santavirta, MD, and others..... 935		

We may return unduly long letters to the author for shortening so that we can offer readers as wide a selection as possible. We receive so many letters each week that we have to omit some of them. Letters must be signed personally by all their authors. We cannot acknowledge their receipt unless a stamped addressed envelope or an international reply coupon is enclosed.

Too few necropsies

SIR,—In writing of his difficulty in elucidating the long-term behaviour of sarcoid heart disease (25 August, p 501), Dr H A Fleming criticises the inadequacy of some necropsies. He also complains of the falling rate of necropsies, and suggests that this may be attributable to the lack of help given to clinicians by inadequate postmortem examinations. I wish to make some personal observations on these matters and also to comment on behalf of the Royal College of Pathologists.

In my experience there is considerable variation in the interest of individual clinicians in the necropsy. Among clinical units served by the same department of histopathology, I know of one which achieves a necropsy rate of over 90%; the clinicians (including consultants) attend necropsies on their patients and give the pathologist a full clinical account of the patient's illness, stressing points of particular concern. In such cases a thorough necropsy is performed with extensive histological and often biochemical studies, to the mutual benefit of clinicians, pathologists, and students. By contrast, there are some clinical units in which the necropsy rate is low, the history supplied to the pathologist is quite inadequate, and the clinicians are reluctant to attend necropsies. In these instances examination is often inadequate because the pathologist is not fully aware of the clinical problems relating to the case. In general, the benefit of a necropsy to the clinician is related directly to the interest he takes in it. It is certainly difficult and disheartening to try to

maintain a high quality of service without clinical interest, and the educational value of a necropsy, which should be a clinicopathological conference, is greatly diminished when the clinical presentation is based on a few words scribbled on a request form.

I do not wish to suggest that pathologists are blameless for deficiencies in the necropsy service. This college has noted complaints of various clinical specialty groups, particularly in cases of perinatal, infant, and maternal deaths and also in deaths following anaesthesia. The college is emphasising the importance of adequately supervised necropsy work during training and has made obligatory the performance of a necropsy in the MRCPath final examination in histopathology.

Another reason for inadequate necropsies is, as Dr Fleming notes, the shortage of pathologists. Staffing levels are well below those recommended by the college¹ and inevitably the surgical pathology and cytology services take priority over the necropsy service. The college is making strenuous efforts to improve recruitment. It is perhaps worth noting also that the necropsy rate has not fallen in all hospitals and where it has that this may be due to the increased numbers of deaths in hospital and not necessarily to a decline in the number of necropsies performed.^{2,3} Post-mortem examinations requested by coroners present a particularly difficult problem, for they are often performed by a busy pathologist trying to provide a satisfactory hospital service; accordingly the examination is often

restricted to ascertaining the cause of death. This problem is also receiving the college's close attention.

J R ANDERSON
President,
Royal College of
Pathologists

London SW17 5AF

¹ Royal College of Pathologists, *Staffing and Workload*. London, Royal College of Pathologists, 1972.

² Cameron, H M, *et al*, *British Medical Journal*, 1977, **1**, 1577.

³ *British Medical Journal*, 1977, **1**, 1560.

Reactions to whooping cough vaccine

SIR,—In his letter on reactions to whooping cough vaccine Dr J S Robertson (22 September, p 735) rightly points out that the liability to brain damage cannot be assessed by statistical studies on the part played by coincidence in apparent time relationships between immunisation and convulsions. But the alternative he proposes does not go to the root of the problem.

Above all we need to establish whether whooping cough vaccine ever has caused brain damage, however rarely—and in some of the case reports of Werne and Garrow,¹ Byers and Moll,² and Brody³ coincidence is clearly out of the question. Next we have to determine the full range of the manifestations of this brain damage; the constituents of the vaccine giving rise to it; the likely pathological mechanisms, toxic or allergic; and why only a few children suffer while most escape. We

also have to balance the risks of the vaccine against the benefits of immunisation in the community, and this is the true role of epidemiology. The other problems are clinical or pathological. But in an attempt to adapt them to epidemiological investigation oversimplifications have been introduced which obscure the fundamental issues. The most important of these oversimplifications is the use of "convulsions within 72 hours of inoculation" as a convenient criterion of vaccine damage.

The confusion began with Miller and Stanton's description of the typical neurological reaction.⁴ "Clinically," they wrote, "the picture is a striking one, most commonly of convulsions occurring 20 minutes to 72 hours after injection, followed by coma and hemiplegia or both." Attentive reading of their citations, however, will show that the cases on which they based this description were selected *because* they presented with convulsions. All non-striking presentations, immediate and delayed, were consequently overlooked. In the course of repeated summaries and transcriptions the rule then became "pertussis vaccine encephalopathy begins with convulsions"; then "Brain damage is not attributable to the vaccine unless there were convulsions within 72 hours of inoculation"; and finally the coma and hemiplegia were split off and discarded and the syndrome became indistinguishable from the benign febrile convulsions of infancy. It could then be statistically proved that whooping cough vaccine was not a cause of febrile convulsions, when nobody claimed that it was.

It is still possible that any of the reported neurological consequences of the parenteral use of any foreign protein or any vaccine, or of the toxins which are active in whooping cough itself, may in time prove to occur after whooping cough vaccine. It is also possible that in children liable to convulsions brain damage from the vaccine manifests itself in the form of convulsions, whereas in children not so liable the onset is less striking. Until such questions and a host of others are settled it is too early to think of balancing the risks.

P E BROWN

Department of Community
Medicine,
University Medical School,
Sheffield S10 2RX

¹ Werne, J, and Garrow, I, *Journal of the American Medical Association*, 1946, **131**, 730.

² Byers, R K, and Moll, F C, *Pediatrics*, 1948, **1**, 437.

³ Brody, M, *Journal of the American Medical Association*, 1949, **140**, 555.

⁴ Miller, H G, and Stanton, J B, *Quarterly Journal of Medicine*, 1954, New Series **23**, 1.

SIR,—Dr J S Robertson (22 September, p 735) has hit the nail—or in this case the needle—on the head, in his criticisms of Professor Gordon Stewart's analysis of neurological ills from whooping cough vaccine.¹ Five years ago Prensky² with similar arguments showed the weakness of the case against pertussis immunisation. It is thus more than unfortunate that Stewart has continued to propagate his opinions without adding substantial new evidence.

I have had the opportunity to investigate the nature of pertussis immunisation convulsions,³ using the technique previously used to differentiate "anoxic" from "epileptic" febrile convulsions.⁴ Of 12 consecutive children with pertussis immunisation convulsions seen

over a two-year period, 11 had excessive asystole on ocular compression. This exaggerated vagal response to ocular compression in pertussis immunisation convulsions was indistinguishable from that found in "anoxic" febrile convulsions, but significantly different from that in "encephalopathic" febrile convulsions ($P < 0.01$, multiple linear regression analysis). Although pertussis-containing vaccines are undoubtedly reactogenic,⁵ my study indicates that pertussis immunisation convulsions resemble cardiogenic "anoxic" seizures (fainting fits)⁶ and are not evidence of any direct encephalopathic effect of the injected substances.

Since vagal anoxic seizures are preventable by atropine sulphate⁶ or atropine methonitrate (unpublished observations), prophylaxis of pertussis immunisation convulsions may be a practicable proposition and deserves study.

J B P STEPHENSON

Fraser of Allander Assessment Unit
and EEG Department,
Royal Hospital for Sick Children,
Glasgow G3 8SJ

¹ Stewart, G T, *Journal of Epidemiology and Community Health*, 1979, **33**, 150.

² Prensky, A L, *Developmental Medicine and Child Neurology*, 1974, **16**, 539.

³ Stephenson, J B P, *Lancet*, 1979, **2**, 416.

⁴ Stephenson, J B P, *British Medical Journal*, 1978, **2**, 726.

⁵ Barkin, R M, and Pichichero, M E, *Pediatrics*, 1979, **63**, 256.

⁶ Stephenson, J B P, *Archives of Disease in Childhood*, 1978, **53**, 193.

Rubella vaccination

SIR,—Dr Nicholas Black (22 September, p 735) advocates the serious consideration of immunisation against rubella without serotesting. In support he quotes figures by Mayon-White and Bull,¹ who found only 50% acceptance for serotesting.

It would have been fairer had he also quoted papers showing a much higher acceptance rate. For instance, the paper by myself and my colleagues² reports a study in which virtually all the patients approached agreed to be tested and where only two patients out of 55 did not return for vaccination. This would have produced cost figures very different from the £20 a conversion quoted by Dr Black.

Dr Black states "that immunising without serotesting is not without medicolegal difficulties." I feel we should be worrying more about the ethical considerations of producing terminations where we need not have done so had we ignored the cost.

MAX GRINGRAS

Poynton, Cheshire SK12 1RA

¹ Mayon-White, R T, and Bull, M J V, *Practitioner*, 1976, **216**, 317.

² Gringras, M, et al, *British Medical Journal*, 1977, **2**, 245.

Circumcision and cruelty to children

SIR,—Conscientious surgeons and paediatricians have listed the complications of neonatal circumcision both in this journal (Drs P M Fleiss and J Douglass (1 September, p 554)) and in other scientific papers. Without anaesthetic the operation in babies causes pain, intense and prolonged crying, air swallowing, vomiting sometimes followed by apnoea, and sometimes permanent local complications. With anaesthetic "the procedure should not be considered minor," because of both local and anaesthetic com-

plications. There is a small risk of death following circumcision, which means that there must be clear-cut immediate medical indications to justify that risk. Some arguments in favour of circumcision for long-term reasons smack of sophistry.

Drs Fleiss and Douglass dispassionately described a cruel practice. If circumcision without anaesthetic had not been sanctioned by authority, there would be grounds for criminal prosecution of the operator for inflicting unnecessary cruelty, and grounds (in the UK) for a care order removing the child from parental authority. Indeed, such action has been taken in the UK against parents who inflict bizarre (but less common) tribal mutilations on their young children. Doctors do not supervise or administer facial cicatrization (with the appropriate sterile precautions and antibiotics) to help the child identify with his ancestral tribe. They do not remove toes because of a long-term risk of bunions, poor hygiene, ingrowing toenails, or toe cancer.

If circumcision for non-medical reasons has died out in Scandinavia and is dying out in the UK, why has the practice persisted in parts of the UK and much of the United States? Do religious and social pressures perpetuate unnecessary circumcision (or any other unnecessary mutilation or operation)? It may be true that the stress to a child in a Moslem enclave in the UK would be great if he were excluded from being the same as his peers, and made a pariah from his group. If this is correct, cannot circumcision for religious (or social) reasons be delayed until 16 years, when the young man can take the positive decision? He may then be determined to accept risk and pain for the sake of his beliefs. It just seems wrong for authority to continue to encourage certain parents to subject young children to unnecessary risk, pain, and distress, and for young children to be denied any protection from the law just because a malpractice is rather common.

J E OLIVER

Burderop Hospital,
Swindon, Wilts SN4 0QA

Risks of amniocentesis

SIR,—The review article on amniocentesis by Barbara J Culliton and Wallace K Waterfall (22 September, p 723) is based on part of the report of the Consensus Development Conference on Antenatal Diagnosis¹ which took place at the US National Institutes of Health in March 1979. The article, like the report, refers briefly to the Medical Research Council's assessment of the risks of amniocentesis,² both references carrying the implication that the MRC findings can be ignored because of methodological shortcomings, and thus do not cast doubt on the general conclusion that "midtrimester amniocentesis is both safe and effective," as they say in their article.

We wish to make two points. Firstly, this was an internal conference with no representation from any country outside the United States, so that the "consensus" does not necessarily extend to other countries. Secondly, a number of methodological criticisms of the MRC study from the United States have been raised since the conference,^{3 4} and have been shown to be without foundation.^{5 6}

The MRC study of the risks of amniocentesis