CORRESPONDENCE

Prospects in pathology J R Anderson, PRCPATH.....	1217	Fetal malnutrition—the price of upright posture? A Briend, MD, and D G Altman, BSC.....	1221	Hong Kong dog and the Tokyo trots R S Thubron, MB.....	1225
Dangerous antihypertensive treatment F Jonasson, MD.....	1218	The premature breech M J Johnstone, MRCOG.....	1222	Primary anaerobic peritonitis J Totten, FRCSed.....	1225
Renal artery stenosis and hypertension P Sharpstone, FRCP, and others.....	1218	Stress and premature labour Judith M Lumley, MB, and R J Bell, MB... ..	1222	Treatment of hyperhidrosis B Bedford, MB, and E V McKinnon.....	1225
Burning and inadequate management of epilepsy W F Durward, MRCP, and M G Harrington, MRCP.....	1218	Reproductive mortality G V P Chamberlain, FRCOG.....	1222	Penis captivus W G Mills, FRCOG.....	1225
Poisoning with chlormethiazole G T McInnes, MRCP, and others; R W Morris, MB.....	1218	Abnormal smears in pregnancy W A Curtain, FRCS.....	1222	Medicine in the USSR C G Iwegbu, FRCS.....	1225
Neutropenia associated with metronidazole T Bergan, MD; R Windle, FRCS.....	1219	Pregnancy in patients presenting with hyperprolactinaemia W F Kelly, MRCP, and others.....	1222	Lack of flexibility in vocational training? C J R Hart, MRCP.....	1225
Withdrawal of cyanocobalamin N McD Davidson, MRCP; A H Goodspeed, MB.....	1219	Preventing postoperative thromboembolism E J Immelman, FRCS, and others.....	1223	The middle grade E G Herzog, MB.....	1226
Treatment of tumours in children J Martin, FRCP, and Jillian R Mann, MRCP.....	1220	Hyperprolactinaemia and cardiac disorders G Curtarelli, MD, and C Ferrari, MD.....	1223	Medically qualified preclinical academics C W Ogle, MB.....	1226
Clinical photographs of children D P Addy, MRCP.....	1220	Vasodilators in senile dementia J A Yesavage, MD.....	1223	Consensus management G G Savage, FFA.....	1226
Putting children first Jean M West, MB.....	1220	What are accident and emergency departments for? J V V Lewis, MRCP.....	1224	Medical examination for elderly drivers J R Greaves, MB.....	1226
Accidental injury in children and interrogation of families.....	1220	Disasters, flying squads, and immediate care K C Hines, MB.....	1224	Advertising for locums for one week or less G Pfarrer, MB.....	1227
Circumcision and cruelty to children R T Marcus, FRCS.....	1220	Appendicectomy during caesarean section in a developing community C H W Bullough, MB.....	1224	Saving money on the drug bill F Honigsbaum, PHD.....	1227
Pregnancy hazards and the child's charter G M Jones, FRCGP, and J J M Arter, LLB... ..	1221	Appendicectomy and family history J C Leedham-Green, FRCS.....	1224	The care of children in hospital J Wilson, FRCP.....	1227
No raw deal for breast-fed babies A Lucas, MRCP, and J D Baum, FRCP.....	1221	Fulminating meningococcal septicaemia D T Thomas, MFCM.....	1224	If I was forced to cut D N S Kerr, FRCP; J F Moorhead, FRCP; P R F Bell, FRCS; R J Jarrett, MD; P Davies, FRCP.....	1227
				Correction: Wanted: a new wound dressing (L E Hughes).....	1228

We may return unduly long letters to the author for shortening so that we can offer readers as wide a selection as possible. We receive so many letters each week that we have to omit some of them. Letters must be signed personally by all their authors. We cannot acknowledge their receipt unless a stamped addressed envelope or an international reply coupon is enclosed.

Prospects in pathology

SIR,—Your leading article "Prospects in pathology" (22 September, p 692) presents a picture which I believe to be unduly pessimistic.

Firstly, it must be emphasised that career prospects in the major pathological specialties (chemical pathology, haematology, histopathology, and medical microbiology) are excellent. Demand for consultants far exceeds supply and there is strong evidence that it will continue to do so. Secondly, it is not true that those interested in a career in a pathological specialty must make an early and irrevocable decision. Following the preregistration year, trainees are required to spend at least 18 months in recognised departments before taking the college's primary examination; this consists of a single multiple-choice question paper in the four specialties plus immunology and a practical and oral examination in any one specialty. Having passed the primary examination the trainee is requested to complete a total of five years, postregistration training before taking the final examination in any one specialty (which need not be the same as the specialty selected for the practical part of the primary). The final examination is regarded as the appropriate qualification for

appointment to a consultant post. In accordance with the increasing clinical bias of the pathological specialties, the MRCP, FRCS, and MRCOG each carries exemption from the college's primary examination, and up to one year of training in a relevant clinical post is accepted as part of the five years' training in pathology. Accordingly, a trainee who has obtained one of the above clinical qualifications can take the final examination after the four-year period usually regarded in most specialties as normal for higher professional training (this could be reduced if the trainee has held a post in pathology before taking the MRCP, etc). I suspect that these arrangements are not widely known, particularly by trainees in the clinical specialties, who may also not all be fully aware of the intensity of the competition for senior registrar posts in the major clinical specialties. It might be helpful to print in the *BMJ* the review of career prospects published annually by the Departments of Health.

As regards merit awards, it is not true to say that pathologists fare badly. As shown in the *BMJ* (18 August, p 456), they are above the average for all consultants, and indeed of the 15 specialties in which there are over 200

consultant posts in England and Wales pathology has a higher proportion of merit award holders than all except general medicine and general surgery.¹ The position is similar in Scotland. It is, however, true that training posts, at least in histopathology, usually carry fewer UMTs than many of the acute clinical specialties.

You mention arguments about who should manage laboratories. These have affected very few laboratories, and in the very great majority there is full co-operation and mutual respect between all grades of staff. Nor should doctors be deterred from taking up pathology because of the increasing number of non-medical scientists working in NHS laboratories. The consultant and top-grade scientist are a strong combination; their work is partly complementary and there is room for both. During 1966-1977 the number of consultant posts in pathology increased by 42%, which is above the average increase for all specialties. Our clinical colleagues appreciate the support of consultant pathology services and will continue to do so provided that consultant pathologists apply their combined clinical and laboratory training to diagnostic problems and the monitoring of patients' progress. This should

include frequent visits to wards to discuss problems relating to individual patients. As you point out, few doctors have opportunity to work with pathologists during their undergraduate course and the preregistration year, and in consequence they are usually unaware of how satisfying and rewarding a career in pathology can be. You refer to a paper of mine which was unfortunately published in incomplete form.² These matters are discussed in a subsequent complete version.³

Finally, the nature of the pathologist's work allows of some flexibility in its performance. It is well suited to part-time training and consultant practice, and the part-timer can usually make arrangements which are compatible with other demands. For this reason, and also because of the relative availability of senior registrar and consultant posts, pathology has particular advantages for doctors whose working hours and mobility are restricted by domestic duties.

JOHN ANDERSON
President, Royal College
of Pathologists

London SW1Y 5AF

¹ Institute for Manpower Studies, *The Determinants of Doctors' Career Decisions*, 1979.

² Anderson, J R, *BMA News Review*, 1979, 5 (4), 26.

³ Anderson, J R, *BMA News Review*, 1979, 5 (6), 72.

Dangerous antihypertensive treatment

SIR,—I refer to your leading article (28 July, p 228), the report by Dr D H Cove and others (p 245), and the following case report.

In July 1978 a 24-year-old housewife was admitted as an emergency to the National Hospital of Iceland. She had sinus tachycardia of 120 beats/min, blood pressure of 300/200 mm Hg, haematuria, and blurred sight of the left eye, but no cardiac enlargement or focal neurological signs. Fundoscopy showed grade IV hypertensive retinopathy. She was twice given diazoxide 300 mg intravenously, with a one-hour interval, together with frusemide; after this the blood pressure fell by one-third only to rise again. Sodium nitroprusside infusion¹ was then commenced, the drip rate being adjusted according to the blood pressure. This was discontinued eight hours later, the blood pressure being 120/80 mm Hg, and from then she received atenolol 100 mg twice daily and diuretics. Ten hours later, when her blood pressure was 110/75 mm Hg and she had sinus bradycardia, she complained of total visual loss. On examination, she did not perceive light, her pupils were bilaterally dilated and unresponsive to light, and fundoscopy showed grade IV hypertensive retinopathy with a macular star.

Assuming that the cause of her blindness was imbalance between the perfusion pressure to her eyes and the intraocular pressure, I immediately commenced a 25% mannitol infusion, and she received 300 ml within 15 minutes in order to reduce the intraocular pressure and so restore the balance between the two pressures. Within 20 minutes the pupils began to react to light and the patient reported that she was seeing again.

Three hours later the patient again lost her sight, her blood pressure being 90/60 mm Hg, when the same treatment was unsuccessfully repeated together with intravenous dexamethasone. After 24 hours' duration of blindness, by the time her blood pressure was 180/130 mm Hg the patient again reported that she was perceiving light and she continued to improve from then on. The underlying cause for her hypertensive crisis was not found, and she has been well controlled on antihypertensive treatment. Visual acuity is 6/6 in each eye, the visual fields are full, and eight months later hypertensive retinopathy was no longer present.

I would like to suggest that in conditions like this lowering of the intraocular pressure

might be considered, with either mannitol or acetazolamide sodium intravenously.

Visual loss in temporal arteritis is considered to be associated with a fall in the perfusion pressure of the extraocular arteries with subsequent imbalance between this and the intraocular pressure.² In that particular disease there are reports suggesting that regaining sight after visual loss of more than 24 hours' duration is unusual.^{3 4}

FRIDBERT JONASSON

Eye Department,
Landakot Hospital,
Reykjavik, Iceland

¹ Cohn, J N, and Franciosa, J A, *New England Journal of Medicine*, 1977, 297, 27.

² Hayreh, S S, *British Journal of Ophthalmology*, 1974, 58, 955.

³ Cullen, J F, *British Journal of Ophthalmology*, 1967, 51, 513.

⁴ Jonasson, F, Cullen, J F, and Elton, R A, *Scottish Medical Journal*, 1979, 24, 111.

Renal artery stenosis and hypertension

SIR,—Reporting the development of renal artery stenosis in two hypertensive patients with previously normal renal angiograms, Dr Mattias Aurell (5 May, p 1180) suggested that "the vascular disease caused the hypertension." A similar sequence of events in a patient of ours has led us to the opposite conclusion.

The patient presented at the age of 54 with a short history of symptoms of uraemia and left ventricular failure. Her blood pressure was 260/140 and her fundi showed bilateral papilloedema with haemorrhages and exudates. Her blood urea was 66 mmol/l and creatinine clearance 1.3 ml/min. The kidneys were normal in size and shape radiologically, so a kidney biopsy was carried out and the histology showed hypertensive nephrosclerosis. Because of clinical features suggesting the possibility of a pheochromocytoma, bilateral selective renal angiography was done. There was no evidence of a pheochromocytoma and the major renal arteries were normal, but there was an intrarenal arteriovenous fistula at the site of biopsy in the left kidney.

During three years of haemodialysis and treatment with antihypertensive drugs her hypertension was never well controlled and her blood pressure ranged from 160/100 to 210/210. Since the iatrogenic arteriovenous fistula could have been contributing towards her hypertension we considered surgical treatment and a second renal angiogram was carried out. The fistula had disappeared and the kidney had shrunk to 10 cm in length, but there was now a tight stricture of the left main renal artery 1.5 cm from its origin with poststenotic dilatation. On the right there were multiple stenoses of the right main and intrarenal arteries.

This patient had radiologically normal main renal arteries at the time of her presentation with hypertension, as did one of Aurell's patients. We do not know the pathology of the stenosis she later developed, but it seems reasonable to postulate that her continuing hypertension may have been the cause. Although many patients have been cured of hypertension by correction of their renal artery stenosis, we wonder whether some of the surgical failures may have been because their hypertension was the cause rather than the consequence of their renal artery stenosis.

PAUL SHARPSTONE
ANTHONY TRAFFORD
RICHARD MCGONIGLE

Royal East Sussex County
Hospital,
Brighton BN2 5BE

Burning and the inadequate management of epilepsy

SIR,—Burning is a potentially fatal hazard faced often by the poorly controlled epileptic. We have been impressed—and depressed—by the poor quality of previous management of epilepsy in patients admitted to the regional burns unit in this infirmary. Admission is sought for patients with especially large burns, burns involving important areas such as hands or face, skin grafting, etc. We studied the management of 10 consecutive patients with epilepsy admitted to the burns unit and offer a brief summary of two cases together with comment on the others.

A 51-year-old man had a three-year history of tonic-clonic seizures following skull fracture. Despite taking phenytoin sodium regularly in a dose of 100 mg thrice daily control remained poor. During a seizure he sustained an extensive full-thickness burn to his right foot from an electric fire. The serum phenytoin sodium level of admission was 10 µmol/l (therapeutic range 40–80 µmol/l) and he required increase in dose to 700 mg daily to obtain control.

A 26-year-old man had an eight-year history of tonic-clonic seizures of idiopathic type. Despite taking, or claiming to take, phenobarbitone 60 mg thrice daily control remained poor but specialist help was not sought. During a seizure he sustained a full-thickness burn to his left leg. The serum phenobarbitone levels were below the therapeutic range of 40–120 µmol/l.

Of the eight other patients, one was untreated, two were on inappropriate therapy, one failed to take all doses of the prescribed anticonvulsant, two had inadequate doses of an appropriate anticonvulsant (judged by continuing seizures in the presence of low serum levels), and two were taking several anticonvulsants at inadequate doses. Of the ten patients, only one had been seen by a neurologist and two had been admitted to general medical wards to seek better control.

The risks of anticonvulsant therapy have become increasingly apparent.^{1 2} The lack of benefit and hazards which often result from therapy with several anticonvulsants have been brought to our attention.² We agree with a policy of seeking control on minimum effective therapy, and with the reluctance to use more than one anticonvulsant until all appropriate drugs have been tried alone in adequate doses. However, undertreatment, including inappropriate treatment, also carries risks, of which severe burning is an especially dramatic example.

At the risk of being thought patronising and pompous, we wish to remind our colleagues that optimal control of correctly diagnosed epilepsy depends on the selection of an appropriate anticonvulsant which is then given in adequate dose. Specialist advice should be sought for refractory cases; we would arbitrarily define as such the failure to respond to a single anticonvulsant.

W F DURWARD
M G HARRINGTON

Royal Infirmary,
Glasgow G4 0SF

¹ Richens, A, in *Drug Treatment of Epilepsy*, ed A Robin. London, Henry Kimpton, 1976.

² Reynolds, E H, *Lancet*, 1978, 2, 721.

Poisoning with chlormethiazole

SIR,—We were interested in the report by Dr R N Illingworth and others (13 October, p 902) of severe poisoning with chlormethiazole. All their patients had taken oral overdoses, and while the previously noted effect of this