psychiatric morbidity looming large on the horizon.

A comprehensive psychiatric assessment and management at this point of crisis in a patient's life can have a very important preventive role, as well as providing for early detection of psychiatric illness in the making before the prognosis becomes unfavourable and unnecessary demands are made on psychiatric services, including long-term care.

R K BRAHMA

Whipps Cross Hospital,
London E11

¹ Kreitman, N (editor), *Parasuicide*. New York, Wiley, 1977.

² Kreitman, N, *British Journal of Psychiatry*, 1979, **135**, 275.

Carbon monoxide poisoning

SIR,—I was interested to read the timely warning by Drs A J Crisp and K M Sherry (1 December, p 1438) regarding poisoning by carbon monoxide. It should be remembered that carbon monoxide is produced by the incomplete combustion of any of the common fuels, and that such incomplete combustion is likely to be due, in the domestic scene, to insufficient ventilation.

May I add a rider to the warning from Drs Crisp and Sherry? I am concerned that cases of subacute or chronic carbon monoxide poisoning may be going undetected. The symptoms—"headache, nausea, want of energy, a tired feeling, lack of concentration, irritability, breathlessness, tachycardia, giddiness, gastrointestinal disturbances, and ataxia"¹—are so all-embracing that patients presenting with this type of picture are likely to be regarded as neurotic and receive short shrift at the hands of casualty officers and others.

I would recommend inquiry about fires and other heating appliances in such cases. The diagnosis is easily made today by the use of the Dräger respiratory carbon monoxide test tube to analyse the patient's exhaled air. This device will give an instant, accurate estimation of the carbon monoxide content of exhaled air, from which the percentage of carboxyhaemoglobin in the blood can be derived. It is accurate over the range 2-35%—and it is also remarkably helpful in educating smokers.

NEVILLE DAVIS

Brownlow Medical Centre,
London N11 2BD

¹ *British Encyclopaedia of Medical Practice*, 1953, **12**, 126.

Carbon monoxide yield of cigarettes

SIR,—Dr Keith Ball's (22 September, p 731) suggestion that the Government should publish carbon monoxide (CO) yields of individual brands of cigarettes is based on three conclusions of doubtful validity.

Firstly, in stating that the evidence on CO as an important health hazard for smokers is growing he ignores the recent important retraction by Astrup¹ of his earlier studies² of the effect of CO exposure on normolipidaemic rabbits. Though there is evidence that CO can increase the severity of coronary artery disease in cholesterol-fed pigeons, rabbits, and monkeys, there is now no evidence that CO exposure in animals or birds fed on normal diets has any effect.³

Secondly, Dr Ball's conclusion that CO

levels have increased owing to the switch to filter cigarettes, which occurred mainly in the 1960s,⁴ is based on comparisons of yields of filter and plain cigarettes made much more recently and is in fact likely to be incorrect. The CO yield of unventilated cigarettes is strongly related to the permeability of the cigarette paper⁵ (and also to the size of the cigarette), and cigarette paper used currently in either filter or plain cigarettes is much more permeable than that of the typical old plain cigarette of 20 or more years ago. Gori⁶ has recently concluded that CO yields in US brands appear to have declined and it seems likely that considerable reductions have occurred in UK brands also—a view strengthened by the observation that a brand manufactured in 1937 delivered over 43 mg CO per cigarette, more than twice that delivered by present-day cigarettes.⁶ Dr Ball's fears that the "higher" CO levels may have led to more deaths from coronary heart disease are in any case at variance with the epidemiological evidence, which shows a reduced risk of coronary heart disease in smokers of cigarettes with reduced tar and nicotine levels⁷ and in smokers of filter rather than plain cigarettes.⁸

Finally, Dr Ball's statement that the publication of the tar tables stimulated the industry to lower tar levels of cigarettes ignores the fact that by far the major part of the tar reduction occurred *before* the publication of the first tar table in 1973 on the industry's own initiative. Average tar levels were 31.4 mg in 1965, 18.7 mg in 1973, and 17.4 mg in 1978.

There seems no strong case for publication of CO yields.

PETER N LEE

Cheam, Surrey SM3 8PY

¹ Stender, S, *et al*, *Atherosclerosis*, 1977, **28**, 357.

² Astrup, P, *British Medical Journal*, 1972, **1**, 447.

³ Turner, D M, *et al*, *Atherosclerosis*, in press.

⁴ Lee, P N, *TRC Research Paper No 1*. London, Tobacco Research Council, 1976.

⁵ Gori, G, and Ellis, R L, *Preventive Medicine*, 1979, **8**, 358.

⁶ Spincer, D, and Evans, N, Imperial Tobacco Limited Report No R417, August 1976.

⁷ Hammond, E C, *et al*, *Environmental Research*, 1976, **12**, 263.

⁸ Dean, G, *et al*, *TRC Research Paper No 14*, part I. London, Tobacco Research Council, 1976.

Rectal dosage of metronidazole

SIR,—May I draw the attention of your readers to an error in the ninth edition of *A Paediatric Vade-mecum*¹?

The rectal dose of metronidazole is incorrectly given (on p 191) as 500 mg three times daily for ages 2 weeks to 7 years. This is a toxic dose at the lower age range and should not be given. The May and Baker data sheet states that under 1 year the dose should be 125 mg (a quarter of a suppository), from 1 to 5 years 250 mg (half a suppository), and from 5 to 12 years 500 mg (1 suppository) eight hourly.

The suppositories are difficult to subdivide accurately and there is probably a need for lower-dose suppositories to be made available.

I am very grateful to Dr J Goulton of Messrs May and Baker for drawing my attention to this error and apologise to readers for its occurrence.

BEN WOOD

Department of Child Health,
Southampton General Hospital,
Southampton SO9 4XY

¹ Wood, B (editor), *A Paediatric Vade-mecum*, 9th edn, p 191. London, Lloyd-Luke, 1977.

Addiction to smoking and drinking—our most serious preventable diseases

SIR,—Smoking and drinking give rise to poisoning of the organism by tobacco and alcohol. Poisoning of the organism constitutes a disease: a "morbid condition" of the body (Oxford English Dictionary). And because of their prevalence the addictions to tobacco and alcohol are by far the most serious avoidable diseases in the Western world—for both are preventable and curable. Yet we continue to evade taking effective action, on the misconception that they are merely habits and not addictive diseases. We accept unquestioningly as diseases their several complications, such as bronchial carcinoma and cirrhosis, but still fail to indict and control the primary addictive conditions.

LENNOX JOHNSTON

Lymington, Hants

JOHN ANDERSON

Taunton, Somerset

Trial of trimipramine for depression

SIR,—I never expected to see such an article as that by Dr J P R Young and others (24 November, p 1315) emanating from Dr William Sargent's old hospital.

It is surprising to find that anybody would nowadays take "a large, unselected group of depressed outpatients" for treatment without sorting them into recognised clinical categories.

There are many reports which indicate that the monoamine oxidase inhibitors (MAOIs) are the treatment of choice in so-called atypical depression. Does this investigation tell us whether trimipramine is as good as, better, or worse for the treatment of this group of patients? Patients are individuals and they come to us not to have their Hamilton or Beck or MRC scores lowered but to be made to feel better. Does this study show us how to do this?

The authors conclude that neither MAOIs nor MAOIs combined with trimipramine is the treatment of first choice in unselected patients with depression. Those of us who were treating patients in the early 1960s learned that. Since then we have learned something about the selection of patients.

W L JONES

Mapperley Hospital,
Nottingham NG3 6AA

Management of haemorrhoids

SIR,—The advocates of stretching, banding, and freezing of haemorrhoids are many, but rarely do they define exactly the severity of their patients' symptoms, nor do they compare their results with management by injection therapy and haemorrhoidectomy.

The latest paper by Mr M R B Keighley and others (20 October, p 967)¹ is an example of this. An attempt has been made to divide patients into two groups on the basis of pressure studies, but their definition of the two groups is rather curious. Their first or high-pressure group were mainly men (99/108) and pain and bleeding were the principal symptoms (92/108), but their second or low-pressure group—"usually women with prolapse as a principal symptom"—contained 55 men and 53 women, while 56 had bleeding as a