window seal. The fact that the authors did not find a perilymph fistula does not exclude this possibility as they can heal very rapidly, as after routine stapedectomy. I note that tinnitus and nausea had resolved by the time of surgery. Incidentally, experimental inoculation of mumps virus at the oval window does not lead to labyrinthitis in monkeys.³

A G GORDON

London SE5

¹ Overman, J R, *Archives of Internal Medicine*, 1958, **102**, 354.

² Foldi, M, *American Heart Journal*, 1977, **93**, 121.

³ Karmody, C S, *Annals of Otolaryngology and Rhinology*, 1975, **84**, 179.

Abdominal and thoracic pressures during defecation

SIR,—Following the article by Dr S S Fedail and others (13 January, p 91) entitled "Abdominal and thoracic pressures during defecation" I would like to comment on the method and interpretation of the results.

Firstly, the technique of locating the diaphragm by the point of respiratory reversal on the manometry tracing is somewhat erroneous. The point of respiratory reversal is normally at the upper end of the high-pressure zone (sphincter + diaphragm) above the lower oesophageal sphincter component. The diaphragmatic component of the high-pressure zone normally appears at the lower (distal) aspect. However, the proximal tube would undoubtedly have been in the oesophagus and the pressures they measured would have accurately reflected the difference between stomach and thorax. There is naturally a significant difference between these results, but the important factor is the gastro-oesophageal pressure difference. There was no significant difference between the squatters and the sitters. The authors fail to mention the important closing and antireflux mechanisms that occur at the gastro-oesophageal junction—for example, angle of implantation, mucosal flap, and right crura, all of which depend to a large extent on a positive intra-abdominal pressure.

In conclusion, I think that their idea is correct and our mode of defecation may well be a causative factor in the development of hiatal hernia. However, it is unwise to base the argument on such flimsy "not significant" evidence.

J A C THORPE

Thoracic Surgical Unit,
Papworth Hospital,
Cambridge

SIR,—Dr S S Fedail and others (13 January, p 91) put forward evidence to support the opinions of Muller¹ and Burkitt and James² that the pressure gradient between abdomen and thorax during straining at stool is an aetiological factor in hiatus hernia.

Although attractively simple, this does not withstand any detailed examination, since it fails entirely to explain why the oesophagus withdraws *pari passu* as the stomach advances up the mediastinum, as is the case in sliding hiatus hernia, where the oesophagogastric junction is always at the apex of the herniated stomach. The sliding type of hernia comprises the vast majority of all hiatus hernia. It fails also to explain the phenomenon of incompetence of the oesophagogastric sphincter,

usually found in association with sliding hernia, but often found in its absence.

Damage to the hiatus—for example, surgical trauma—results not in the sliding type of hernia but in the paraoesophageal type, in which the fundus of the stomach, or a part of it, together with the oesophagogastric junction, are herniated into the lower mediastinum, without disturbance of function of the sphincter, and without shortening of the oesophagus. One would expect this type of hernia to result from a raised pressure gradient across the diaphragm, as in straining at stool; but it does not.

The prevalence of hiatus hernia in the West and its rarity in Africa invite the conclusion that a principal aetiological factor is diet. With this I agree; but I feel that this factor induces a disorder of oesophagogastric motility, the important feature being over-activity of the longitudinal muscle of the oesophagus. This leads to oesophageal shortening, which draws the stomach into the chest, producing the characteristic sliding hernia. It is this shortening—and thickening—which distorts, elongates, and renders incompetent the oesophagogastric sphincter. It is this shortening which often renders repair of the hernia difficult, and sometimes impossible. Surgical procedures which overcome the shortening or the tension in the oesophagus relieve the symptoms, although they need not necessarily reduce the hernia.

Unfortunately we have no means of measuring the activity of the longitudinal muscle of the oesophagus, despite having had elegant methods of measurement of the circular muscle for many years. As we cannot measure longitudinal muscle activity, we prefer to ignore it. But the longitudinal muscle is by far the most prominent feature of the oesophagus.

In short, sliding hiatus hernia is not pushed up into the chest by abdominal pressure; it is pulled up by the oesophagus. I have previously elaborated this, basing my views on surgical experience.³

KENNETH S MULLARD

Bletchington, Oxon

¹ Muller, C J B, *South African Medical Journal*, 1948, **22**, 376.

² Burkitt, D P, and James, P A, *Lancet*, 1973, **2**, 128.

³ Mullard, K S, *Annals of the Royal College of Surgeons of England*, 1972, **50**, 73.

Physical therapy in chronic bronchitis

SIR,—Dr Margaret E Hodson and others (6 January, p 55) correctly observe that we did not separate the effect of intermittent positive pressure breathing (IPPB) and physiotherapy (2 December, p 1525). The similar response of our treated and control groups suggests that this was an economic and wise decision, unless one considers these forms of treatment mutually antagonistic.

I would be surprised if any alteration in physiotherapy technique or method of administration of bronchodilator would produce better results than we obtained in either of our treatment groups, but shall await the evidence.

Your correspondents quoted an excellent paper by Shenfield *et al* as evidence that a bronchodilator is more efficient by IPPB than by aerosol.¹ The actual title of the paper was "The fate of nebulised salbutamol (Albuterol) administered by intermittent positive respiration to asthmatic patients." As the title suggests, it was concerned with the pharmacodynamics of inhaled salbutamol, and blood

levels of the drug over eight hours and urine and faecal samples over three days. One could not suggest from the data presented in the paper that the use of IPPB with nebulised salbutamol was more effective than a pressurised aerosol alone, as the authors did in the summary. Furthermore, this paper referred to five convalescing asthmatics. We considered the paper by Berend *et al* more appropriate, as it concerned the disease we were studying and for that reason quoted it.²

The final point in the letter about the importance of physiotherapy and the use of a bronchodilator for those who have large volumes of secretions, is discussed in the *Lancet* editorial to which it refers.³

DUNCAN NEWTON

Service de Pneumologie,
Hôpital Saint Antoine,
Paris

¹ Shenfield, G M, *et al*, *American Review of Respiratory Diseases*, 1973, **108**, 501.

² Berend, N, *et al*, *British Journal of Diseases of the Chest*, 1978, **72**, 122.

³ Newton, D A G, *Lancet*, 1979, **1**, 49.

An unsavoury custom?

SIR,—It is one of the marks of a first-class hospital that its doctors and laboratory staff do not come to meals wearing their white coats. I know of no spread of an infection which was traced to this unsavoury custom; but why wait till something untoward occurs before discontinuing the practice where it still persists? Admittedly there are often administrative difficulties, notably in teaching hospitals where changing facilities would have to be found for numerous clinical students. The DHSS definitely frowns on this custom. If medical men do not feel concerned, perhaps the administrators in charge of hospital dining rooms and canteens ought to act and exclude clothing which might be contaminated from areas where food is served and consumed.

H LEHMANN

Department of Biochemistry,
University of Cambridge

Hazards of sulphinyprazone

SIR,—With reference to the letter on hazards of sulphinyprazone (23-30 December, p 1786), I would like to make some comments.

Sulphonyprazone being a pyrazole compound is related to phenylbutazone, which has been observed to potentiate the hypoglycaemic effect of the oral sulphonylureas. Hence sulphinyprazone should be used with caution in patients on concomitant sulpha drugs, the sulphonylurea hypoglycaemic agents, and insulin.

Sulphonyprazone is potent uricosuric agent, and alkalisation of urine and high fluid intake are necessary at first to avoid crystalluria. In patients on phenylbutazone oedema occurs, due to a renal effect in reducing sodium and water excretion. Since sulphinyprazone is closely related (a pyrazole) to phenylbutazone the salt retention as a side effect should also be borne in mind, for it may precipitate heart failure.

I endorse the note of caution of Dr D Mattingly and his colleagues that sulphinyprazone is contraindicated in patients on oral anticoagulants. This drug is also contraindicated in patients with history of hypersensitivity

to phenylbutazone and related compounds, sulfadiazine, and sulfisoxazole. Patients with an active peptic ulcer should not be given sulphinpyrazone. Finally it should be publicised that blood dyscrasias have been reported to be due to this drug, although rarely.

I hope these comments will heighten the clinical awareness of the side effects of sulphinpyrazone.

SAEED AHMAD

Fairmont General Hospital,
Fairmont,
West Virginia, USA

Tuberculin testing

SIR,—I refer to the note on tuberculin testing (13 January, p 108), in which reference is again made to the unreliability of the tine tuberculin test. The only alternative mentioned is the Mantoux test. This calls for accurate intracutaneous injection of the tuberculin. I find that young doctors have not been trained in this technique and do not trust themselves (nor are trusted by their seniors) to perform the test satisfactorily.

This causes me to ask what has become of the Heaf test. I used it with complete confidence for many years and advised its use throughout Northern Ireland in connection with the BCG vaccination programme of the Northern Ireland Tuberculosis Authority. The Heaf gun demands no special skill in its operation. Has it already been discarded as an obsolete instrument?

H G CALWELL

Carrickfergus,
Co Antrim

Nice people with no manners

SIR,—“Nice people with no manners” by Dr Stephen Lock (23 December, p 1774) aptly portrays problems which can slightly mar an enjoyable evening for the speaker. These are, as he shows, not confined to our profession. Indeed Stephen Leacock, humorist and academician, wrote an essay on “How it feels to be a lecturer.” He wittily described experiences he had when touring England and elsewhere: chairman had forgotten his name or got it wrong and one had to read out the subject of his talk from the programme. He wrote that “the first of the troubles . . . is the fact that the audience will not come to hear him.” Research could be done on what inspires doctors to attend meetings: how important is the subject, the speaker—or perhaps the meal? Some years ago as secretary to a medical society I got a speaker who had filled a vast hall in New York. Here the usual audience of about 19 attended. Shortly afterwards I invited an expert on income tax for doctors: the place was packed.

Most speakers could vie with each other in telling depressing tales—for example, an audience hardly more numerous than the speakers who had travelled many miles through snow and ice on a Sunday. Another time, after being surrounded by about 200 people when being given sustenance just before the lecture, as the time approached the crowd faded away leaving a handful of people, as the lecture had coincided with the hospital ball. Nevertheless, it was a pleasant evening, as the audience was one of quality rather than quantity.

Fortunately, the nice people with no

manners are, in my experience as well, rare. Usually the clearest details are given about the time, place, duration of talk, and type of audience—and hospitality is generous. The commonest difficulties are technical, particularly when showing slides. Some problems are avoidable, such as having a spare bulb, spare projector, and someone who understands the machine. Others are unpredictable, such as poor attendance or a power cut.

CLIFFORD HAWKINS

Queen Elizabeth Hospital,
Birmingham

Haemodynamic effects of buprenorphine after heart surgery

SIR,—It is impossible to find justification in the article by Dr F L Rosenfeldt and others (9 December, p 1602) for their conclusion that “Buprenorphine appears to be safer than morphine for use in patients with reduced cardiac reserve. . . .”

No evidence is presented by the authors to suggest that their patients, studied after open heart surgery, were in an unstable cardiovascular state or had a reduced cardiac reserve and the effects of morphine were not studied. The finding of a 24 mmHg fall in the mean arterial pressure in 18% of the patients studied suggests that the hypotensive effect of intravenous buprenorphine may be clinically relevant. The subsequent statement by the authors that “there was no overall change in mean arterial pressure” cannot be evaluated as no relevant data are provided.

One must conclude that buprenorphine may cause a fall in arterial blood pressure in some patients and that there is no justification on the evidence presented in the study for considering it to be safer than morphine as an intravenous analgesic.

D W BETHUNE

Thoracic Surgical Unit,
Papworth Hospital,
Cambridge

* * * We sent a copy of this letter to the authors, whose reply is printed below.—ED, *BMJ*.

SIR,—Thank you for allowing us to reply to the important issues raised by Dr Bethune.

The purpose of this paper was to report the haemodynamic effects of buprenorphine after cardiac surgery. In keeping with the style of the *BMJ* we trimmed our paper in many directions and are therefore delighted for the opportunity of supplying further information to Dr Bethune. He may judge from our previous publications on a similar group of patients^{1,2} that these patients do indeed have an unstable cardiovascular system. This is well recognised by most workers in the immediate period following open heart surgery, when such patients have to be treated in an intensive care unit to monitor closely the circulatory state.

Haemodynamic response in eight patients followed for 60 minutes after administration of buprenorphine
Values are mean $\pm$ 1 SEM

	Minutes after administration					
	0	5	10	15	30	60
Mean arterial pressure (mm Hg)	85 $\pm$ 6	85 $\pm$ 6	85 $\pm$ 5	83 $\pm$ 5	87 $\pm$ 5	82 $\pm$ 5
Cardiac output (l/min)	3.6 $\pm$ 0.3	3.6 $\pm$ 0.3	3.6 $\pm$ 0.3	3.5 $\pm$ 0.3	3.7 $\pm$ 0.3	3.5 $\pm$ 0.3
Heart rate (b/min)	103 $\pm$ 4	103 $\pm$ 5	100 $\pm$ 5	100 $\pm$ 5	100 $\pm$ 5	99 $\pm$ 4

Regarding Dr Bethune's query about the fall in arterial pressure, table printed below shows the change in mean arterial pressure, cardiac output, and heart rate, justifying our statement that “there was no overall change in mean arterial pressure.”

We carefully referred to our review of the haemodynamic effects of other analgesic agents³ and we have directly compared buprenorphine with Omnopon in the same patients, using the protocol we outlined in our paper. Considering the known haemodynamic effects of morphine in such patients we firmly believe that our conclusions are justified.

D J COLTART
D THOMPSON
N NAQUI

Department of Cardiology,
St Thomas's Hospital,
London SE1

¹ Lewis, G J R, et al, *European Journal of Cardiology*, 1978, 7, 283.

² Lewis, G J R, et al, *American Heart Journal*, 1978, 95, 301.

³ Malcolm, A D, and Coltart, D J, in *Pain—New Perspectives in Measurement and Management*, ed A W Hargreaves, R B Smith, and B A Whittle, p 41. Edinburgh, Churchill Livingstone, 1977.

Normality and abnormality in psychiatry

SIR,—The letter of Dr J G Edwards (4 November, p 1296) is a timely reminder of the problems of delineating normality in psychiatric practice. Indeed, it has been cynically noted in a university students' publication that those who are normal are “other people a psychiatrist has never met.”¹ That there may be an element of truth in this is attested to by the fact that “homoclitie” was coined to indicate a mentally healthy young man.² Furthermore, it has been suggested that those persons who claim to be free of any psychological symptoms may only be unusually capable of concealing their symptoms.³

“Normality and the psychiatrist” was the subject of a recent Australian paper.⁴ In addition to the statistical, clinical, and prognostic aspects referred to by Professor Geoffrey Rose and Dr D J P Barker (23 September, p 873) and the sociocultural component noted by Dr Edwards, this alluded to the fact that a Utopian ideal of health is sometimes regarded as normal and to the differing standards of normality taken in a developmental sense.

There is no doubt that normality must be considered according to many parameters, and one can only endorse Dr Edwards's comments that it is most important for doctors to recall that the designation of abnormality has far-reaching implications. For the individual it may involve expensive and uncomfortable investigations, isolation from the community, and uncertainty for the future. For society, it offers a means of categorising an individual.