prostacyclin production if the non-precursor fats are peroxidised.⁴ Certainly, to choose a margarine with a high content of C18:2ω6 in the cis form needs more information than is provided on the label, and none exists with a high content of eicosapentaenoic acid, which may be preferable.

From the point of view of a healthy cardiovascular system, it could be that we should be debating a fat intake based on fish rather than meat, instead of butter versus margarine.

JOHN VANE

Wellcome Research Laboratories,
Beckenham, Kent

¹ Moncada, S, and Vane, J R, *British Medical Bulletin*, 1978, **34**, 129.

² Hamberg, M, Svensson, J, and Samuelsson, B, *Proceedings of the National Academy of Sciences of the USA*, 1975, **72**, 2994.

³ D'Angelo, V, et al, *Thrombosis and Haemostasis*, 1978, **39**, 535.

⁴ Glavind, J, et al, *Acta Pathologica et Microbiologica Scandinavica*, 1952, **30**, 1.

⁵ Salmon, J A, et al, *Biochimica et Biophysica Acta*, 1978, **523**, 250.

⁶ Dyerberg, J, et al, *Lancet*, 1978, **2**, 117.

⁷ Raz, A, Minkes, M S, and Needleman, P, *Biochimica et Biophysica Acta*, 1977, **488**, 305.

⁸ von Lossow, T O, et al, *American Journal of Clinical Nutrition*, 1978, **31**, 1340.

⁹ Nordoy, A, et al, *Circulation Research*, 1978, **43**, 527.

¹⁰ Sanders, T A B, et al, *Lancet*, 1977, **1**, 751.

¹¹ Crawford, M A, et al, in *Dietary Lipids and Postnatal Development*, ed, C Galli, p 127, New York, Raven Press, 1973.

¹² Weihrauch, J, et al, *Food Technology*, 1977, **31**, 80.

SIR,—We should like to join the debate on fats and atheroma initiated by Sir John McMichael (20 January, p 173). Three points raised in that article need separating and clarifying: (a) the role of cholesterol in the atheromatous plaque; (b) hypercholesterolaemia as a risk factor for atherosclerosis; and (c) the effects of changing to a diet low in cholesterol and high in unsaturated fat.

The atheromatous plaque consists of a thickened area of the arterial intima and media due to an accumulation of proliferating smooth muscle cells surrounded by collagen, elastin, glycosaminoglycans, and fibrin. Cholesteryl ester is deposited in massive amounts around and within the smooth muscle cells.^{1, 2, 3} Although the histology of the plaque is complex there is evidence that it is the cholesterol in the lesion which plays a central role in its development. Severe atherosclerosis can be produced in nearly every species of experimental animal by feeding a diet that raises the plasma cholesterol level.⁴ Proliferative thickening of the arterial wall can be produced in animals by trauma, such as hypertension, or by destruction of the endothelium by physical or chemical methods; but the lesion formed is just a simple scar unless the animal has hypercholesterolaemia, in which case cholesterol is deposited in the lesion and severe atherosclerosis develops.¹

Studies on human populations have shown that symptomatic atherosclerosis does not frequently develop, even when there are predisposing factors such as diabetes mellitus, cigarette smoking, and hypertension, unless the mean concentration of plasma cholesterol in the population exceeds 4.1 mmol/l (158 mg/100 ml).⁵⁻⁸ Within any single Western population the probability of an individual developing myocardial infarction due to coronary heart disease increases in proportion to his plasma cholesterol level.⁹ Single gene-determined disorders such as familial hypercholesterolaemia, in which there is markedly elevated plasma and tissue cholesterol levels, are characterised by fulminant atherosclerosis

in childhood, without any other contributing factors such as hypertension, smoking, or diabetes mellitus being present.¹⁰

Thus there is good experimental and epidemiological evidence demonstrating the central role of cholesterol in the atheromatous plaque and linking hypercholesterolaemia to the development of atherosclerosis. This evidence, accumulated over several years, should not be cast aside lightly.

Recent evidence has shown that experimental atherosclerosis, especially in monkeys, develops more slowly or regresses in animals given diets which reduce serum lipid concentrations.¹¹⁻¹⁴ There have been three major primary prevention trials in human populations, and they have all shown that diets which reduce serum lipids significantly reduce the number of ischaemic events.¹⁵⁻¹⁷

Sir John McMichael points to the unknown hazards of a diet rich in polyunsaturated fats. For the patient with familial hypercholesterolaemia the prospect of death from atherosclerosis in childhood outweighs this consideration. For those members of a Western population who do not have this uncommon disorder but who have serum cholesterol levels well above 4.1 mmol/l, since 52% of all deaths in males aged 45-54 are due to cardiovascular disease¹⁸ it is perhaps prudent to suggest that they avoid a high cholesterol intake. Thus if they limited themselves to two eggs per week, stopped eating cream, reduced butter and milk intake, avoided fat on the meat, and reduced total meat intake, they might gain benefit from a reduced serum cholesterol level without the unknown dangers of excessive quantities of polyunsaturated fats.

K G TAYLOR

Department of Medicine,
The General Hospital,
Birmingham

D J BETTERIDGE

Department of Medicine,
St Bartholomew's Hospital,
London EC1

¹ Wissler, R W, in *The Myocardium: Failure and Infarction*, ed E Braunwald, p 155. New York, HP Publishing Company, 1973.

² Haust, M D, *Human Pathology*, 1971, **2**, 1.

³ Smith, E B, and Smith, R H, in *Atherosclerosis Reviews*, vol 1, ed R Paoletti and A M Gotto Jun, p 119. New York, Raven, 1976.

⁴ Roberts, J C, Jun, and Straus, R, (editors), *Comparative Atherosclerosis*, p 3. New York, Harper and Row, 1965.

⁵ Keys, A, *Circulation*, 1970, **41**, suppl 1, p 1.

⁶ Stamler, J, *Medical Clinics of North America*, 1973, **57**, 5.

⁷ Keys, A, *Atherosclerosis*, 1975, **22**, 149.

⁸ Epstein, F H, *Atherosclerosis*, 1971, **14**, 1.

⁹ Kannel, W B, et al, *Annals of Internal Medicine*, 1971, **74**, 1.

¹⁰ Fredrickson, D S, Goldstein, J L, and Brown, M S, in *The Metabolic Basis of Inherited Disease*, ed J B Stanbury, J B Wyngaarden, and D S Fredrickson. New York, McGraw Hill, (4th edn in press).

¹¹ Armstrong, M L, Warner, E D, and Connor, W E, *Circulation Research*, 1970, **27**, 59.

¹² Wissler, R W, et al, *Circulation*, 1971, **44**, suppl II, p 57 (abstract).

¹³ Armstrong, M L, and Megan, B M, *Circulation Research*, 1972, **30**, 675.

¹⁴ Vesselinovitch, D, et al, *American Journal of Pathology*, 1973, **70**, 75 (abstract).

¹⁵ Christakis, G, et al, *American Journal of Public Health*, 1966, **56**, 299.

¹⁶ Dayton, S, et al, *Circulation*, 1969, **40**, suppl II, 1.

¹⁷ Miettinen, M, et al, *Lancet*, 1972, **2**, 835.

¹⁸ Joint Working Party of the Royal College of Physicians of London and the British Cardiac Society, *Journal of the Royal College of Physicians of London*, 1976, **10**, 219.

Mentally disordered offenders

SIR,—I have studied with interest the letters (27 January, p 263) from Drs H R Rollin, S B Telford, K C Bailey, and D D R Williams on the subject of mentally disordered offenders.

Between them they have set out most of the historical steps by which such persons developed into a major social problem; but, apart from one comment by Dr Williams, they offer no practical solution to the problem.

Dr Williams recalls that one stimulus to the creation of asylums was the large number of mentally ill persons in prisons, since when the number of such persons in prisons has varied inversely with the accommodation made available for them in other institutions. He therefore is concerned lest the Government policy to run down large mental hospitals will aggravate the problem facing the prisons. I disagree with Dr Williams only in his equation of asylums with psychiatric hospitals. As the modern trend is to concentrate all psychiatric treatment in general hospitals and community centres, it stands to reason that the demand for such treatment in psychiatric hospitals must diminish; but that is no argument for ceasing to make any use of the buildings—indeed, there is a case, as mentioned by Dr Telford, for letting them revert to their former use as asylums. Dr Bailey would prefer them to be called "forensic hospitals," but that title suggests that they would provide treatment, and therefore would require the services of psychiatrists and other specialised staff. A well-run asylum does not need such staff. I should know, as I was born in an asylum and both my parents were asylum doctors. In my childhood most of the staff were attendants, not yet persuaded to seek the qualification of certificated mental nurse, which even then had been available to them for a generation. They did a good job of work, and handled the inmates with skill, understanding, and kindness.

Dr Rollin comments on the "open door therapeutic policy" now in vogue. But what, pray, is the connection between therapeutics and open doors? Before the 1930 Act initiated voluntary admission, all patients admitted were subject to judicial orders for their detention. Yet when the Woodilee Hospital in Scotland was opened as an asylum in 1877 it became immediately fully operative, although it had no locked doors right up to the time, I believe, of the first world war. Whatever caused the doors to be closed thereafter had nothing to do with therapeutics. I have no doubt that Woodilee, in its 30 years of open doors, dealt with the very same kind of difficult patients that are now refused admission on the paltry excuse of open doors.

The responsibility for creating the social problem of the generally rejected mentally disordered offender rests firmly, I regret to admit, on the shoulders of psychiatrists. Nothing in the 1959 Act authorised us to systematically destroy the main function envisaged for the large mental institutions at the time they were built—namely, to provide asylum. There is no law to stop us now from restoring conditions in any part of a psychiatric hospital so as to provide full security for those in need of it, and for negligible expense. Through a laissez-faire attitude on the part of public authorities, psychiatrists have been allowed so to abuse their powers as to make a mockery of the courts' procedure for dealing with mentally disordered offenders. Now even Dr Rollin has joined the ranks of those clamouring for the long-debated regional special secure units. Leaving aside the waste of public money demanded for such units when perfectly suitable buildings are already available, the psychiatrists have already announced their intention to use them in a

way which will do absolutely nothing to relieve the problem they have created for the courts. The great majority of the problem patients facing the courts are not in need of treatment—they are seeking asylum. Such people, unlike all the law-abiding mentally ill, will raise no objection to being housed in the vicinity of a building providing full security for criminals, for they did not object to it in the past.

As I was one of those who contributed suggestions for the Bill which subsequently became the 1959 Act, I must bear a share for its shortcomings. Our greatest mistake, which led to so much disruption of public services, was the delusion that the psychopath may respond to medical treatment. This purely English delusion reached its truly grandiose zenith in the setting-up of Grendon Underwood Prison, which has always specialised in the alleged treatment of psychopaths, to the exclusion of all prisoners with generally accepted forms of mental illness. The canny Scots committed no such folly when formulating their 1960 Act. Likewise the Northern Irish disclaimed medical responsibility for psychopaths in their 1961 Act. Alas, there is no United Kingdom in the field of psychiatry.

If the present-day psychiatrists wish to pursue the policy of running down mental hospitals as places unsuitable for the provision of treatment, then surely the logical step is for them to leave such places. In their own words, the present inmates do not need their services, and their hands are full in trying to meet the demand for their services elsewhere. Put bluntly, the so-called psychiatric hospitals could revert more smoothly to their former role of asylums if the psychiatrists would only get out of the way.

With some reservations because of ill-devised trends of policy in the social services, I would agree with Dr Telford that these now defunct psychiatric hospitals might suitably be taken over by social service departments. The only safeguard I would wish for is that the planning of asylums should not be placed in the hands of the psychiatric group of the medical profession (I hesitate to call it a discipline), because we have an appalling record of administrative mismanagement, and not all of us have regained insight.

O W S FITZGERALD

Salcombe, Devon

Emotion and empiricism

SIR,—We believe that your recent editorial entitled "Emotion and empiricism" (3 February, p 288) gave a muddled and misleading account of the scientific basis and ethics underlying the randomised controlled trial.

The clinical trial is of particular value when, as is usually the case, a new treatment is expected to be only a slight improvement over its predecessors. In such circumstances the difference in treatment effects is of the same order of magnitude as the biological variation in response between individuals and therefore it is essential that the clinical trial be of the best possible design. Any suspicion of systematic bias will lead to the results of the trial not being generally accepted and thus the investigators will have failed in their intention of settling the question concerning superiority or equivalence of the treatments. If from the outset, by the very nature of the trial's design,

there is doubt about whether or not it will gain widespread acceptance as an unbiased and objective experiment then it must be considered unethical. This we believe to be the case with clinical trials using historical controls. No matter how careful the investigator he will not be able to demonstrate that the historical controls are strictly comparable with the current cases. The process of randomisation guarantees that randomised controlled trials, as a class, are free from systematic bias. This does not excuse the investigator from checking the effectiveness of the randomisation in any particular instance, but the results of a well-designed randomised controlled trial will always gain greater credence than those from clinical trials employing historical controls.

If a clinician truly believes that one treatment is superior to alternatives then he should not enter his patients into a randomised controlled trial comparing those treatments. Furthermore, if a new treatment is obviously superior to its predecessors, as was the case when the antibiotics were first introduced, then a randomised controlled trial is unnecessary. No reputable statistician or clinician would ever subscribe to the philosophy "Randomise and let the individual go hang." We believe, however, that very few treatments show dramatic benefits and that in most cases of competing treatments no clinician can have a truly rational basis for preferring one treatment to another until a randomised controlled trial has been completed.

Randomised controlled trials of particular treatments are often claimed to be unethical merely because the treatments have been in general use for some time. This argument, if valid, surely makes a case for the speedy evaluation of new treatments as soon as they arise. Perhaps the greatest need for the randomised controlled trial lies in the weeding out of the many untested and probably ineffective treatments in current use.

A L COCHRANE

Rhoose, South Glamorgan

A S ST LEGER
P SWEETNAM

Medical Research Council
Epidemiology Unit,
Cardiff

SIR,—Your leading article (3 February, p 288) states: "Perhaps the most important thing for us to realise is that, after countless years of empiricism, the controlled trial has emerged as a sharp tool for dissection of certain problems. . . . Some guidelines are visible: when there is no rational or a priori basis for preferring one treatment to another the prospective controlled trial is wholly appropriate."

Empirical means (*Shorter Oxford Dictionary*) "Based on, or guided by, the results of observation and experiment only." The supreme error of our profession over "countless years" has been to derive treatment, not from observation and experiment, but from a priori reasoning—that is, from the antithesis of empiricism. We bled because the invalid's blood was believed to contain phlegma, which must clearly be eliminated by phlebotomy. We starved the feverish because fever is analogous to a fire, so just as a fire is damped down by depriving it of fuel, fever is damped down by depriving the patient of food. Septic foci were extirpated because toxins were being absorbed from them to cause sciatica, rheumatoid arthritis, peptic ulcer, etc. Analeptics were given to the poisoned because they antagonised hypnotics and therefore *must* be effective.

Women were kept in bed after labour because "if everyday activities are resumed when the uterus is as large as it is on the 10th day of the puerperium, it is clear that there must be a liability to prolapse, retroversion, and subinvolution."¹ And the victims of cardiac infarction were kept in bed for six weeks or more to "ensure firm cicatrization of the ventricular wall."² Examples could be multiplied ad infinitum. All these remedies, which we now know from observation and experiment—that is, on empirical grounds—to have been worse than useless were believed to be rational by those who recommended them.

The controlled trial is a particularly good example of empiricism because it is an experiment depending on unusually sophisticated observations. Rational and empirical are not, as you suggest, antitheses. The only rational ground I know for giving a remedy is the existence of empirical evidence that it is effective.

JOHN W TODD

Farnham, Surrey

¹ Percival, R, *Holland and Brews Manual of Obstetrics*, 13th edn. London, Churchill, 1969.
² Lewis, T, *Diseases of the Heart*, 3rd edn. London, Macmillan, 1942.

SIR,—Your leading article "Emotion and empiricism" (3 February, p 288) adds another useful point of view to the philosophy and conduct of clinical investigation. We agree that "some guidelines are visible" and would like to air two others, both concerning potentially difficult subjects: the ethics of informed consent and the nature of the study population.

"Informed consent" is an emotive issue, especially when the end-point of assessment is the survival or death of patients. It is important that the real objective of "trials" be clearly recognised. The argument sometimes goes as follows: "We know that no particular adjuvant treatment is successful (for the treatment of this malignant disease), nor is it likely to be, but a trial does give an opportunity for people to work together and to stipulate regimens for patient care, and also a reason to document outcome." We feel that this approach is ethically indefensible because of the harm done to patients when informed consent is requested; the individual's very existence may be seen to depend on this toss of a coin.

The wish to arrange collaborative studies to investigate the nature and outcome of treatment is legitimate and sometimes necessary, but to insist on a randomised trial format, merely to achieve this objective, is unnecessary.

Perhaps we should redefine the specific guidelines about informed consent as follows: (a) where the whole group can benefit from the result of the trial (for example, a new analgesic in the treatment of rheumatoid arthritis) then informed consent should pose no problems; (b) in contrast, when the relative frequency of death is the arbiter of success or failure, and this result can only be applied to *future* generations of patients, then the need for a randomised controlled trial should be established beyond doubt and, if it is, then the nature of informed consent should not necessarily involve the patient in the decision making.

Having decided that a randomised trial is necessary for a particular study, most clinical investigators try, in the period of trial design, to set exclusion clauses so that the population studied can be constrained to obtain greater uniformity. At first sight this seems a reasonable objective, but we believe it is a potential