prostacyclin production if the non-precursor fats are peroxidised.⁴ Certainly, to choose a margarine with a high content of C18:2ω6 in the cis form needs more information than is provided on the label, and none exists with a high content of eicosapentaenoic acid, which may be preferable.

From the point of view of a healthy cardiovascular system, it could be that we should be debating a fat intake based on fish rather than meat, instead of butter versus margarine.

JOHN VANE

Wellcome Research Laboratories,
Beckenham, Kent

¹ Moncada, S, and Vane, J R, *British Medical Bulletin*, 1978, **34**, 129.

² Hamberg, M, Svensson, J, and Samuelsson, B, *Proceedings of the National Academy of the Sciences of the USA*, 1975, **72**, 2994.

³ D'Angelo, V, et al, *Thrombosis and Haemostasis*, 1978, **39**, 535.

⁴ Glavind, J, et al, *Acta Pathologica et Microbiologica Scandinavica*, 1952, **30**, 1.

⁵ Salmon, J A, et al, *Biochimica et Biophysica Acta*, 1978, **523**, 250.

⁶ Dyerberg, J, et al, *Lancet*, 1978, **2**, 117.

⁷ Raz, A, Minkes, M S, and Needleman, P, *Biochimica et Biophysica Acta*, 1977, **488**, 305.

⁸ von Lossow, T O, et al, *American Journal of Clinical Nutrition*, 1978, **31**, 1340.

⁹ Nordoy, A, et al, *Circulation Research*, 1978, **43**, 527.

¹⁰ Sanders, T A B, et al, *Lancet*, 1977, **1**, 751.

¹¹ Crawford, M A, et al, in *Dietary Lipids and Postnatal Development*, ed, C Galli, p 127, New York, Raven Press, 1973.

¹² Weihrauch, J, et al, *Food Technology*, 1977, **31**, 80.

SIR,—We should like to join the debate on fats and atheroma initiated by Sir John McMichael (20 January, p 173). Three points raised in that article need separating and clarifying: (a) the role of cholesterol in the atheromatous plaque; (b) hypercholesterolaemia as a risk factor for atherosclerosis; and (c) the effects of changing to a diet low in cholesterol and high in unsaturated fat.

The atheromatous plaque consists of a thickened area of the arterial intima and media due to an accumulation of proliferating smooth muscle cells surrounded by collagen, elastin, glycosaminoglycans, and fibrin. Cholesteryl ester is deposited in massive amounts around and within the smooth muscle cells.^{1, 2, 3} Although the histology of the plaque is complex there is evidence that it is the cholesterol in the lesion which plays a central role in its development. Severe atherosclerosis can be produced in nearly every species of experimental animal by feeding a diet that raises the plasma cholesterol level.⁴ Proliferative thickening of the arterial wall can be produced in animals by trauma, such as hypertension, or by destruction of the endothelium by physical or chemical methods; but the lesion formed is just a simple scar unless the animal has hypercholesterolaemia, in which case cholesterol is deposited in the lesion and severe atherosclerosis develops.¹

Studies on human populations have shown that symptomatic atherosclerosis does not frequently develop, even when there are predisposing factors such as diabetes mellitus, cigarette smoking, and hypertension, unless the mean concentration of plasma cholesterol in the population exceeds 4.1 mmol/l (158 mg/100 ml).⁵⁻⁸ Within any single Western population the probability of an individual developing myocardial infarction due to coronary heart disease increases in proportion to his plasma cholesterol level.⁹ Single gene-determined disorders such as familial hypercholesterolaemia, in which there is markedly elevated plasma and tissue cholesterol levels, are characterised by fulminant atherosclerosis

in childhood, without any other contributing factors such as hypertension, smoking, or diabetes mellitus being present.¹⁰

Thus there is good experimental and epidemiological evidence demonstrating the central role of cholesterol in the atheromatous plaque and linking hypercholesterolaemia to the development of atherosclerosis. This evidence, accumulated over several years, should not be cast aside lightly.

Recent evidence has shown that experimental atherosclerosis, especially in monkeys, develops more slowly or regresses in animals given diets which reduce serum lipid concentrations.¹¹⁻¹⁴ There have been three major primary prevention trials in human populations, and they have all shown that diets which reduce serum lipids significantly reduce the number of ischaemic events.¹⁵⁻¹⁷

Sir John McMichael points to the unknown hazards of a diet rich in polyunsaturated fats. For the patient with familial hypercholesterolaemia the prospect of death from atherosclerosis in childhood outweighs this consideration. For those members of a Western population who do not have this uncommon disorder but who have serum cholesterol levels well above 4.1 mmol/l, since 52% of all deaths in males aged 45-54 are due to cardiovascular disease¹⁸ it is perhaps prudent to suggest that they avoid a high cholesterol intake. Thus if they limited themselves to two eggs per week, stopped eating cream, reduced butter and milk intake, avoided fat on the meat, and reduced total meat intake, they might gain benefit from a reduced serum cholesterol level without the unknown dangers of excessive quantities of polyunsaturated fats.

K G TAYLOR

Department of Medicine,
The General Hospital,
Birmingham

D J BETTERIDGE

Department of Medicine,
St Bartholomew's Hospital,
London EC1

¹ Wissler, R W, in *The Myocardium: Failure and Infarction*, ed E Braunwald, p 155. New York, HP Publishing Company, 1973.

² Haust, M D, *Human Pathology*, 1971, **2**, 1.

³ Smith, E B, and Smith, R H, in *Atherosclerosis Reviews*, vol 1, ed R Paoletti and A M Gotto Jun, p 119. New York, Raven, 1976.

⁴ Roberts, J C, Jun, and Straus, R, (editors), *Comparative Atherosclerosis*, p 3. New York, Harper and Row, 1965.

⁵ Keys, A, *Circulation*, 1970, **41**, suppl 1, p 1.

⁶ Stamler, J, *Medical Clinics of North America*, 1973, **57**, 5.

⁷ Keys, A, *Atherosclerosis*, 1975, **22**, 149.

⁸ Epstein, F H, *Atherosclerosis*, 1971, **14**, 1.

⁹ Kannel, W B, et al, *Annals of Internal Medicine*, 1971, **74**, 1.

¹⁰ Fredrickson, D S, Goldstein, J L, and Brown, M S, in *The Metabolic Basis of Inherited Disease*, ed J B Stanbury, J B Wyngaarden, and D S Fredrickson. New York, McGraw Hill, (4th edn in press).

¹¹ Armstrong, M L, Warner, E D, and Connor, W E, *Circulation Research*, 1970, **27**, 59.

¹² Wissler, R W, et al, *Circulation*, 1971, **44**, suppl II, p 57 (abstract).

¹³ Armstrong, M L, and Megan, B M, *Circulation Research*, 1972, **30**, 675.

¹⁴ Vesselinovitch, D, et al, *American Journal of Pathology*, 1973, **70**, 75 (abstract).

¹⁵ Christakis, G, et al, *American Journal of Public Health*, 1966, **56**, 299.

¹⁶ Dayton, S, et al, *Circulation*, 1969, **40**, suppl II, 1.

¹⁷ Miettinen, M, et al, *Lancet*, 1972, **2**, 835.

¹⁸ Joint Working Party of the Royal College of Physicians of London and the British Cardiac Society, *Journal of the Royal College of Physicians of London*, 1976, **10**, 219.

Mentally disordered offenders

SIR,—I have studied with interest the letters (27 January, p 263) from Drs H R Rollin, S B Telford, K C Bailey, and D D R Williams on the subject of mentally disordered offenders.

Between them they have set out most of the historical steps by which such persons developed into a major social problem; but, apart from one comment by Dr Williams, they offer no practical solution to the problem.

Dr Williams recalls that one stimulus to the creation of asylums was the large number of mentally ill persons in prisons, since when the number of such persons in prisons has varied inversely with the accommodation made available for them in other institutions. He therefore is concerned lest the Government policy to run down large mental hospitals will aggravate the problem facing the prisons. I disagree with Dr Williams only in his equation of asylums with psychiatric hospitals. As the modern trend is to concentrate all psychiatric treatment in general hospitals and community centres, it stands to reason that the demand for such treatment in psychiatric hospitals must diminish; but that is no argument for ceasing to make any use of the buildings—indeed, there is a case, as mentioned by Dr Telford, for letting them revert to their former use as asylums. Dr Bailey would prefer them to be called "forensic hospitals," but that title suggests that they would provide treatment, and therefore would require the services of psychiatrists and other specialised staff. A well-run asylum does not need such staff. I should know, as I was born in an asylum and both my parents were asylum doctors. In my childhood most of the staff were attendants, not yet persuaded to seek the qualification of certificated mental nurse, which even then had been available to them for a generation. They did a good job of work, and handled the inmates with skill, understanding, and kindness.

Dr Rollin comments on the "open door therapeutic policy" now in vogue. But what, pray, is the connection between therapeutics and open doors? Before the 1930 Act initiated voluntary admission, all patients admitted were subject to judicial orders for their detention. Yet when the Woodilee Hospital in Scotland was opened as an asylum in 1877 it became immediately fully operative, although it had no locked doors right up to the time, I believe, of the first world war. Whatever caused the doors to be closed thereafter had nothing to do with therapeutics. I have no doubt that Woodilee, in its 30 years of open doors, dealt with the very same kind of difficult patients that are now refused admission on the paltry excuse of open doors.

The responsibility for creating the social problem of the generally rejected mentally disordered offender rests firmly, I regret to admit, on the shoulders of psychiatrists. Nothing in the 1959 Act authorised us to systematically destroy the main function envisaged for the large mental institutions at the time they were built—namely, to provide asylum. There is no law to stop us now from restoring conditions in any part of a psychiatric hospital so as to provide full security for those in need of it, and for negligible expense. Through a laissez-faire attitude on the part of public authorities, psychiatrists have been allowed so to abuse their powers as to make a mockery of the courts' procedure for dealing with mentally disordered offenders. Now even Dr Rollin has joined the ranks of those clamouring for the long-debated regional special secure units. Leaving aside the waste of public money demanded for such units when perfectly suitable buildings are already available, the psychiatrists have already announced their intention to use them in a