adrenaline, or MA plus NMA are due to a chemical rather than to a metabolic effect of labetalol and represent falsely high values. Adequate methods should be employed in future studies evaluating the effects of labetalol on the sympathetic nervous system. Furthermore, our findings may prevent errors in the differential diagnosis of phaeochromocytoma.

RAINER KOLLOCH
LINO MIANO
VINCENT DE QUATTRO

Department of Medicine,
University of Southern California,
Los Angeles, California 90033

- ¹ Harris, P M, and Richards, D A, *British Medical Journal*, 1977, **2**, 1673.
- ² Chapman, B O, Veitch, A B, and Shepherd, B, *British Medical Journal*, 1978, **1**, 364.
- ³ Weidmann, P, et al, *American Journal of Cardiology*, 1978, **41**, 570.
- ⁴ Crout, J R, in *Standard Methods of Clinical Chemistry*, ed D Seligson, vol 3, p 62. New York, Academic Press, 1961.
- ⁵ Peuler, J D, and Johnson, G A, *Life Sciences*, 1977, **21**, 625.
- ⁶ Pisano, J J, *Clinica Chimica Acta*, 1960, **5**, 406.
- ⁷ Bigelow, L B, and Wil-Malherbe, H, *Analytical Biochemistry*, 1968, **26**, 92.
- ⁸ Pisano, J J, Crout, J R, and Abraham, D, *Clinica Chimica Acta*, 1962, **7**, 285.
- ⁹ Miano, L, Kolloch, R, and DeQuattro, V, submitted for publication.

Naming of drugs

SIR,—As a psychiatrist working in Germany and trying against unequal odds to correlate advances in drug therapy in this country and in Britain, I welcome the discussion stimulated by Dr G A MacGregor concerning the naming of drugs (18 November, p 1433).

Why not require the drug companies to display the approved name directly after the proprietary name and at least two-thirds as large? This would apply to advertisements as well as labels on containers, and to combinations of drugs as well as simple compounds. Even non-myopic doctors and nursing staff would then be able to readily associate the two names and pick out whichever is relevant. It would still be allowed to use the approved name with the manufacturer's name in brackets after it as an alternative, but I do not consider it realistic to expect drug companies to give up their cherished brand-names without a long struggle.

The system which I suggest is one which we could reasonably demand be used throughout the EEC.

M H BALL

W Germany

Plasma lipoproteins in renal transplant patients

SIR,—The report on plasma lipoproteins in renal transplant patients by Dr D B Evans and the late Dr P Ghosh (6 January, p 21) invites several comments.

They suggest that the reduction in plasma lipid concentrations was due to a reduction of the mean prednisolone dose from 21 mg/day to 11 mg/day. However, in their original paper¹ lipids were measured in 17 asthmatics receiving a mean dose of prednisolone of 9 mg/day. The cholesterol and triglyceride levels were 6.0 mmol/l (232 mg/100 ml) and 2.38 mmol/l (211 mg/100 ml) respectively, values very similar to those in the transplant patients on the higher dose of prednisolone rather than the lower one. Clearly, then, hyperlipidaemia is not related to the steroid dose alone.

The 17 long-term survivors analysed in 1978 constitute a different population from the original 32. The transplants with poorest function in 1971 would be likely to have failed by 1975, leaving a group of survivors with well-functioning grafts. (The fact that the rise in glomerular filtration rate over the four-year period was not statistically significant does not exclude this possibility.) It has been suggested² that some lipids, including triglyceride, especially in females, are related to graft function. With the progressive elimination of the worst-functioning grafts one would expect the mean triglyceride level of the group as a whole to fall.

The change in lipid profile could be further explained in part by the death of those of the 1971 cohort with the worst lipid profile. In one series of 101 transplant recipients,³ occlusive arterial disease was observed only in those with untreated hyperlipidaemia, while the 29 nonlipidaemic patients remained free of arterial disease over a mean three-year observation period.

It is inevitable that the long-term survivors of renal transplantation will tend to have low steroid requirements, good graft function, and absence of arterial disease; the other transplanted patients will have fallen by the wayside through death or graft failure. Other workers⁴ have followed renal transplant recipients for up to 13 years, and have found that the percentage of survivors with hyperlipidaemia progressively falls in the years following transplantation. However, the clustering of low steroid dose, good graft function, freedom from arterial disease, and normal lipid levels does not necessarily imply a causal relation between any of these factors as Drs Evans and Ghosh suggest.

A J NICHOLLS

Department of Medicine,
University of Aberdeen

- ¹ Ghosh, P, et al, *Transplantation*, 1973, **15**, 521.
- ² Abbott, L K, et al, *Atherosclerosis*, 1978, **30**, 97.
- ³ Ponticelli, C, et al, *Kidney International*, 1978, **13**, suppl 8, S72.
- ⁴ Thomas, F T, and Lee, H M, *Annals of Surgery*, 1976, **184**, 342.

Oesophageal reflux and its myths

SIR,—Your leading article "Oesophageal reflux and its myths" (6 January, p 3) commendably stresses the importance of treating reflux oesophagitis, rather than a sliding hiatus hernia, which may be no more than a chance radiological finding.

The editorial is perhaps guilty, however, of oversimplifying the significance of lower oesophageal sphincter pressure in the aetiology of reflux. As DeMeester and Johnson¹ have shown, 24-hour oesophageal pH monitoring is possibly the most sensitive method of monitoring reflux currently available. Their studies have also shown that reflux occurs in normal subjects and that, even in symptomatic patients, its occurrence depends on a number of factors other than posture. Price and Castell² point out that, while in general people with reflux have lower resting lower oesophageal sphincter pressures than those without, there is much overlap between populations with and without reflux. Furthermore, although the work of Cohen and Harris³ on the influence of gastrointestinal hormones on sphincter pressure has greatly increased understanding of the lower oesophageal sphincter mechanism, it remains true that hiatus hernia repair, by whichever method

may be chosen, prevents reflux by restoring the anatomical position and relations of the lower oesophageal sphincter without altering its hormonal sensitivity. It is probably the case that measurement of lower oesophageal sphincter pressure as currently performed is a relatively indirect indicator of reflux, which may be the result of prolonged periods of sphincter relaxation occurring in response to a variety of alterations in anatomical and hormonal environment.

It is also worth stressing the point that surgical procedures for hiatus hernia repair are, in fact, methods of treating reflux oesophagitis, and that without evidence of reflux oesophagitis there is no indication for repairing a sliding hiatus hernia.

WILLIAM FOUNTAIN

Department of Surgery,
Northwick Park Hospital,
Harrow, Middx

- ¹ DeMeester, T R, et al, *Annals of Surgery*, 1976, **184**, 459.
- ² Price, S, and Castell, D O, *Journal of the American Medical Association*, 1978, **240**, 44.
- ³ Cohen, S, and Harris, L D, *New England Journal of Medicine*, 1971, **284**, 1053.

SIR,—Following your leading article (6 January, p 3) entitled "Oesophageal reflux and its myths" I must express interest as well as concern. Myths usually stem from ancient fables and the article by Price and Castell¹ serves only to perpetuate these. It is well known that reflux can occur without evidence of hiatal hernia. "True oesophageal reflux" cannot be merely defined as a combination of radiological reflux and endoscopic abnormality.

There may be no radiological reflux and at endoscopy the oesophageal mucosa may look macroscopically normal. These tests are well recognised to be only 50-60% diagnostic.² You mention the acid perfusion test and intraluminal pH test as being useful in the detection of gastro-oesophageal reflux but fail to mention the most important, which is the standard acid reflux test.³ Only a thorough history, examination, elimination of biliary dyspepsia, and the appropriate use of oesophageal function tests can discriminate between "medical" and "surgical" reflux. There is no reason why in this modern era we should not apply our "sophisticated technology" to the diagnosis of oesophageal disorders and put the many "myths" on the shelf with the Cyclops and Hercules.

J A C THORPE

Thoracic Surgical Unit,
Papworth Hospital,
Cambridge

- ¹ Price, S, and Castell, D O, *Journal of the American Medical Association*, 1978, **240**, 44.
- ² Clark, J, Moossa, A R, and Skinner, D B, *Surgical Clinics of North America*, 1976, **56**, 1.
- ³ Skinner, D B, and Booth, D J, *Annals of Surgery*, 1970, **172**, 627.

Disrespect for language

SIR,—Pleased at being able to answer one or two of the questions in your Christmas Quiz (23-30 December, p 1778), I looked up the answers. You should not state that claudication is "thought to have originated from the limp of the emperor Claudius." *Claudicare*, to limp, was used often by Lucretius and by Cicero, both of whom were dead before Claudius was born.

And Claude Bernard was not the "discoverer