reward for dealing with specimens referred by general practitioners.

TONY KEABLE-ELLIOTT

Chairman, General Medical Services Committee

BMA House,
London WC1

Opportunity in health visiting

SIR,—Did your correspondents read my letter (9 December, p 1646)? I am sorry that Jill Thornton and 34 other health visitor students (6 January, p 58) should take such exception to my questioning whether health visiting should remain a secondary profession with entry only through nursing. I do believe it is sufficiently important to warrant its own separate training. To quote an analogy from medicine: dentistry is sufficiently different from medicine for there to be a separate but sometimes related training. This does not mean that no dentist may have a full medical training, but the majority do not; most dentists enter dental school intending to become dentists, not doctors. Most health visitors entered nursing training courses intending to become nurses, not health visitors. There are large parts of nurse training which are not relevant to health visiting and for this reason I proposed a basic health care course as the preliminary training of a health visitor.

Finally, I would not presume to suggest the form which such a training might take but I am sure that a profession of such importance would agree its own training programme. I am glad of the enthusiasm expressed by Miss Thornton and her fellow-students and hope that they will question whether health visiting should remain a secondary profession or move forwards to take a prime position. I believe that this is an important challenge.

MICHAEL HALL

Shebbear, Devon

Nursing at a crossroads

SIR,—In reply to Mr I G Schraibman's letter (13 January, p 121) I would like to make the following comments and proposals.

Alarmed by the understaffing of hospitals and the difficulty of recruiting new nurses, Mr Schraibman suggests various adjustments be made to the current nurse training schedule. As most nurses at present do not take A levels, the majority will be leaving school at 16 years of age. To ensure that enthusiasts are not lost in the 18-month period taking them to the minimum entry age of 17½ years, Mr Schraibman suggests accepting 16-year-olds into a 12-month cadetship, the official training beginning at 17 years. I believe that these proposals are both impractical and dangerous.

Firstly, after the second world war, the Wood report, in an attempt to swell the inadequate numbers of nurses, both lowered the entry age and shortened the course. These measures proved ineffective in improving recruitment. Secondly, the drop-out rate is notoriously high in the first 18 months of training, when the student is at her or his most young and immature stage. This is surely likely to be a more frequent occurrence if the age is lowered to one where most people are just beginning to take control of their lives. Thirdly, the only practical

change that has had any positive influence on recruitment is substantial rise in salary.

The other aspect of these proposals is the psychological effect of lowering the age of our nurses, which I believe has dangerous implications. Mr Schraibman appears to underestimate the enormous responsibility of the nurse in the psychological, social, and physical care of a patient. Regimented technical skills play a relatively small part in full nursing care. A comprehensive, sensitive approach to the patient is essential in her or his proper rehabilitation. A 16-year-old, taken straight from the sheltered environment of school, is unlikely to have the maturity and awareness integral to the role of a good nurse. Moreover, a year's cadetship, if effective at all, is likely to forge the standard glib attitudes prevalent in the nursing profession, rather than to lay the foundations for a true understanding of the highly complex role of the nurse.

Having entered nursing at 20 years of age, after two years' experience of working at home and abroad, I feel there are far more radical changes to be made to improve the situation. If mature, capable, and responsible nurses are sought, then attitudes to the students must be changed. The repressive, authoritarian approach to nurses in training is barely tolerable to anyone with any self-respect. If it is tolerated, it often instils a similarly insensitive and unquestioning attitude in the student.

If, as Mr Schraibman says, nurses are the indispensable foundations of patient care, they should be treated accordingly both in status and in financial worth. I can assure Mr Schraibman that when nursing sheds its image of dedicated slavery performed by young girls bidding time until marriage, and becomes a viable profession for intelligent, responsible adults, recruitment will be no problem. Understaffing is indeed a difficulty, severely exacerbated by the continuing public spending cuts; but short-sighted, short-cut measures offer no solution.

ALISON HADLEY

London, NW3

SIR,—I was interested in the letter by Mr I G Schraibman (13 January, p 121) about the restructuring of the nursing shifts.

As a divisional nursing officer I am concerned with the implications to the service of a 37½-hour week, and a lot of work is at present being undertaken to find a satisfactory solution. The shifts suggested do not allow an overlap, which is essential for continuity, and no provision has been made for coffee or meal breaks. The present 40-hour week is exclusive of meal breaks and therefore any restructuring must be based on 37½ hours plus meals.

The majority of my colleagues could introduce a most satisfactory shift system. Unfortunately, this could cause hardship to the nurses involved, as we have a high proportion of married women with domestic responsibilities, and early starts with late finishes could cause conflict—which would eventually lead to staff leaving the service. Another problem is public transport, especially of outlying areas, as not all nursing personnel are fortunate enough to have the use of a car and a 7 am start could mean staff leaving home before 6 am. In the "old days" when a large proportion of staff were single and resident greater flexibility was possible.

The nursing fraternity would welcome help and suggestions from the medical staff as

restructuring of the nursing shifts also entails restructuring of the medical staff's ward and department commitments.

E L THOMPSON

Newcastle General Hospital,
Newcastle upon Tyne

Prescribing—is a second opinion required?

SIR,—The Royal College of Nursing's view that it is illegal for their members and other nurses to write up a patient's oral contraceptive requirements in a delegation session at a family planning clinic seems to be derived, so far as I have been able to ascertain, from the *opinion* of the legal advisers of *their insurers* on an *interpretation* of a particular section of the Medicines Act 1978. But in matters legal there is surely more than one opinion?

The measures of adjustment to meet the situation created by this opinion in practice are cumbersome and time consuming. And inevitably there must be some ethical reservation about the wisdom of a doctor prescribing 13 packets of pills at a time, even when these are to be distributed at a nurse's discretion. What is *his* liability at law and with regard to indemnity? Could his action in so doing ever be interpreted as delegating prescribing? And so on. Furthermore, the new process, by the interruptions and pressure it creates, distracts the proper focus of attention from being on priorities such as changing pills where possible to the low-dosage type, etc, and altogether it disturbs the previous flexibility of the therapeutic contact.

In practice, the doctor is dependent on the nurse's verbal recommendation about required supplies for the patient she considers the doctor does not need to see. It seems to me a very fine point of law to consider that the nurse's writing this out for the doctor's consideration and approval is illegal but that her verbal communication of the same facts to him is (presumably) legal. Much has been made of the fact that under the old system the patient obtained her oral contraceptive supplies a little while (usually not much more than two hours) before the doctor actually signed the prescription and, of course, this is the point at issue. But would any court, in Britain at least, having regard to the safeguards, the high standard of nursing care, and the conscientious action of everyone in good faith, ever do more than simply observe that there was a technical impropriety at law but one for which no reasonable action at law or action for damages could properly be sustained?

Surely, in the present crisis of opinion, it is incumbent now on the medical defence societies, if they are not already doing so, to have defined at law what is meant by "prescribing medically"; to decide who is responsible in a delegation session (whether it is the doctor delegating or whether the nurse is responsible herself for any of the duties she carries out on a doctor's implicit or explicit instructions and, if so, which ones); what the legal risks are to a doctor in the circumstances and his liability in prescribing 13 packets of oral contraceptive pills (specifically with regard to negligence and the charge of delegating prescribing); and whether, in fact, the opinion on interpretation of the RCN's legal advisers is not open to question and dispute and may possibly even be untenable, depending on how prescribing medically is to be defined.