for the clinical convictions of the rescue services, since the inability of the laboratory-based research worker to reproduce the clinical results found in the field may be due to a fault in the design of the laboratory experiment and not, as is presumed at present, to errors of observation by the rescuers.

E L LLOYD

Princess Margaret Rose Orthopaedic
Hospital,
Edinburgh

¹ Kehlet, H, *et al*, *British Journal of Anaesthesia*, 1974, 46, 73.

² Oyama, T, *Canadian Anaesthetists' Association Journal*, 1969, 16, 361.

Rugby injuries to the cervical cord

SIR,—The letters published (20 January, pp 192 and 193) can only increase the concern already felt by the Medical Officers of Schools Association. We urgently need to know the incidence of cervical cord injuries due to rugby football, and hope that the suggestions made by Dr J R Jenner and his colleagues (p 193) for a central reporting system will bear fruit. Meanwhile the association is (conducting its own retrospective survey of these injuries in schoolboys in order to report to a meeting which has been called for 22 March, and I have already written to many colleagues who may know of cases. May I, through the courtesy of your columns, ask any readers who have not already heard from me to get in touch without delay if they can provide information about any cervical spine injury due to rugby football in a schoolboy?

T W HOSKINS

Honorary secretary,
Medical Officers of
Schools Association

The Infirmary,
Christ's Hospital,
Horsham, West Sussex

Rugby injuries involving the hand

SIR,—With reference to the article (23-30 December, p 1759), written by Drs J E Davies and T Gibson about injuries in Rugby Union football, we noted with interest the absence of any significant injuries to the hand reported in their studies of 185 players.

During our six months in a busy accident department in Newcastle upon Tyne, attached to a hand injuries unit, and because of our specific interest in sports injuries, we have noted eight cases of significant injury to the hand as the result of playing rugby.

Seven out of eight injuries resulted from a dislocation of the proximal interphalangeal joint during the tackling of opponents. The eighth injury was a fracture of the second metacarpal, which was the result of a hyper-extension force applied to the fingers during the act of catching the ball. Four out of the initial seven injuries resulted in a boutonniere-type deformity of the fingers as a result of rupturing of the central slip of the extensor expansion as it crossed the proximal interphalangeal joint; all were treated by external splintage.

Of particular interest was the observation that the boutonniere type injuries occurred because of the fingers being caught in the clothing of the opponents during the tackle. In two of the cases it was the pockets of the shorts of the opposition that were the culprits.

It raises the question of the usefulness of pockets in rugby shorts, particularly as we have seen injuries directly attributable to their presence.

We must also make the point that any manual worker who suffers a boutonniere injury to any of his fingers is not able to pursue his normal employment because of the injury and the splintage used in treating the condition. This results in both loss of productivity by employers and, if he is privately employed (for example, as a dentist, like one of our patients), loss of income. Clearly rugby injuries of this sort represent a not insignificant proportion of the upper limb injuries sustained in the game of rugby, and the morbidity resulting from these injuries causes both loss of playing time and loss of income.

R W NUTTON
M J GUY

Accident and Emergency Department,
Royal Victoria Infirmary,
Newcastle upon Tyne

"Trench foot" caused by the cold

SIR,—In this generally temperate climate localised cold injury is relatively uncommon. Because of this we think the following case of "trench foot," presenting at the end of the recent cold spell, to be of some interest.

The patient was an 18-year-old male student who lives in a comfortable, well-heated hall of residence. It was not until the sixth day after the initial exposure that he came for medical advice. The history was that six days previously, after a rather meagre breakfast, he went for a 20-mile cycle ride on a dry, clear day when the temperature was freezing or just below, wearing on his feet stretch-nylon socks and tightly laced training shoes, with a pair of tight bicycle clips round his ankles. He cycled quite slowly for about three hours, with the arches of the feet, not the toes, on the pedals, during which time he realised that his feet were very cold as after a short time he was unable to feel them. At home that evening his right foot began to be painful over the fourth and fifth toes. During the following two days the pain in the right foot increased and he noticed a macerated area between the fourth and fifth toes, but these discomforts did not prevent him walking several miles into university from his hall in the same footwear and in equally cold weather, the temperature averaging 2-4°C. The fourth day was again very cold (-3°C), and again he went cycling. His feet went cold and numb. When he was some four miles from hall his cycle broke down and he had to walk home. That evening, when sensation had returned, the right foot had become swollen and blue; indeed the area near his fourth and fifth toes was almost black, and he was unable to put his shoe or slipper on. Next morning, despite pain and great difficulty getting his right shoe on, he cycled three miles. By mid-day blisters had developed on his toes, and sufficient "sticky" fluid had oozed from his foot for him to be aware of a squelching noise as he walked. However, after this the shoe became easier on his foot and the pain was relieved.

Examination of the foot showed that the dorsum was moderately swollen and was a mottled bluish colour near the toes. There was blanching on pressure, but with poor capillary return. There were small blisters on the great and second toes. The thickly macerated area between the fourth and fifth toes was thought to represent a coincidental flare-up of tinea pedis. He was kept in sick bay and treated conservatively with bed rest, no specific "warming" being applied, and he made a slow and steady recovery. Interestingly, despite the absence of superimposed infection, the small blisters have taken over two weeks to heal.

In addition to the foot condition he developed an itching, macular rash, initially on the hands and

later involving the lower limbs and the umbilical area. Indeed, he primarily sought advice because of the rash. The rash faded without any specific therapy concurrently with the improvement of the foot. Unfortunately it was not possible to perform a cold agglutinins estimation until the eighth day, and the result was negative. We cannot be sure of the exact aetiology of the rash, but believe it on clinical grounds to be directly associated with the cold injury. A full blood count was normal; the erythrocyte sedimentation rate was 2 mm in the first hour.

This is a case of "trench foot" occurring in a cyclist. It demonstrates the necessity, even in our so-called temperate climate, of adequate protection of the extremities even in those who, by virtue of their physical activity, might otherwise expect to keep reasonably warm.

I C FRASER
JUDITH A LOFTUS

University Health Service,
University of Leeds

Epidural analgesic for bronchitic patients with fractured ribs

SIR,—The review of epidural analgesia relating to labour and delivery (13 January, p 72) brings to mind two recent patients who obtained remarkable benefit from this technique.

Both patients were middle-aged bronchitics who had fallen during the previous few days and fractured several ribs. One had been treated at home with bed rest and inadequate oral analgesia; this led to inability to cough owing to severe pain and consequently stasis of secretions resulting in hypostatic pneumonia. The other also obtained little benefit from oral analgesia and opiates were contraindicated because of his history of bronchitis. Both obtained immediate and remarkable relief of pain, were able to breathe and cough properly, and were rescued from impending respiratory failure by the timely relief of pain without any danger of respiratory depression by the use of epidural anaesthesia. I feel that this form of analgesia for the treatment of fractured ribs in the bronchitic patient is not practised sufficiently often.

PETER BAYLOR

St Charles's Hospital,
London W10

Terminology of enuresis

SIR,—I support Mr John R Hindmarsh (16 December, p 1714) in his plea for more precision in the terminology of enuresis. But I would prefer more precision than he shows in his letter. Firstly, it is important to qualify the term enuresis with "night-time" (or nocturnal) or "day-time" (or diurnal). Too often the day-time wetting problems get overlooked and yet are a serious problem to a great many schoolchildren.

The term enuresis is usually reserved for persons who void urine inappropriately at an age when continence is to be expected. I agree that it is inappropriate to use it for persons who have an organic lesion, such as a neuropathic bladder, and are suffering from overflow incontinence. However, to suggest that nocturnal enuresis should be applied only to persons who have "persistently wet the bed all their lives" will cause confusion. It is exceedingly uncommon to find anybody over the age of 5 who has wet the bed every night.