patients, associated with any evidence of drug-induced lupus.

It is possible therefore that extension of Dr Marshall's investigation to larger numbers of patients, with appropriate age-matched controls, could result in conclusions different from those reported. However, their finding that all prazosin-related ANF patterns were of the speckled variety is very unusual. When ANF did occur in patients on prazosin in our study it was not restricted to this morphological pattern. This suggests that there may be some unusual factors influencing results in Dr Marshall's patients.

The full data from which this summary is drawn will be presented elsewhere.

J D WILSON
R J BOOTH
J Y BULLOCK

Department of Medicine,
University of Auckland School of Medicine,
Auckland, New Zealand

¹ Wilson, J D, *et al*, *British Medical Journal*, 1978, 1, 14.

Treatment and survival in advanced breast cancer

SIR,—Dr T Priestman and his colleagues have reported the results of a comparative trial of endocrine versus cytotoxic treatment in advanced breast cancer (16 December, p 1673; 14 May 1977, p 1248). In their trial they administered oestrogen to postmenopausal patients with lung metastases and prednisone to postmenopausal patients with spread to liver or lymphangitis carcinomatosa and compared the therapeutic results with those of cytotoxic agents. In cases of extensive visceral involvement (the authors do not report details about the extension of metastases) the patients are at immediate risk. In these cases we believe it is not justifiable to treat them with oestrogens or only steroids (no matter what kind of trial they did) when it is well known that it takes at least three to four weeks for a hormonal response, whereas an immediate therapeutic response is needed. The treatment of choice is an intensive chemotherapeutic combination (including steroids).

S C TSILIAIOS
A E ATHANASIOU

Oncology Outpatient Clinic,
Idrima Kinonikon Asfaliseon,
Athens, Greece

Synovial biopsy in arthritis

SIR,—I was interested to read your leading article (10 February, p 363) on "Synovial biopsy in arthritis." I was more than a little surprised to find that it contained no reference to the use of arthroscopy in this procedure. Among its uses arthroscopy allows the clinician to comment on the macroscopic appearance of the synovium, examine the joint surfaces, and to find a suitably representative area of synovium for biopsy. The latter can reduce the "sampling errors" of a blind technique.

I am grateful that the leading article threw light on the types of synovitis and their histological differences, but I am sure that many of us involved in synovial biopsy, whether by needle or under vision with arthroscopy, are concerned by the number of times the report simply states a non-specific inflammation.

Surely it is only by building up experience

of macroscopic and microscopic features that a better differential diagnosis of synovitis can be obtained.

MICHAEL EDGAR

Royal National Orthopaedic Hospital,
Stanmore, Middlesex

Epidemiology and public health in American universities

SIR,—May I draw attention to a slip somewhere between pen and print which changed "without" to "with" in my recent article in your columns? (23-30 December, p 1737)? Neither W T Sedgwick, professor of biology and public health at Massachusetts Institute of Technology, or C-E A Winslow, first professor of public health at Yale, had medical degrees. Indeed a purpose of this passage was to draw attention to the fact that some of the leading early teachers of epidemiology in the United States, though bacteriologists, were *not* medically qualified.

Sedgwick embarked on a medical course (at Yale in 1877) but was so disillusioned with the unscientific approach to the subject there at that time that he left New Haven for Johns Hopkins, where he studied biology.¹ Winslow, who was a pupil of Sedgwick's, never studied medicine at all, and as a bacteriologist directed courses in his new department at Yale towards scientists who did not have medical degrees.²

ROY M ACHESON

Department of Community Medicine,
Addenbrooke's Hospital,
University of Cambridge

¹ Curran, J A, *Founders of the Harvard School of Public Health*. New York, Josiah Macy Jr Foundation, 1970.

² Acheson, R M, *American Journal of Epidemiology*, 1970, 91, 1.

Abortion and the NHS

SIR,—An analysis of the first decade of legal abortion in England and Wales is timely because the Wessex Regional Health Authority is now considering whether or not to introduce a regional day case abortion unit. Dr F G R Fowkes and others (27 January, p 217) present a case for more NHS abortions, but do not say whether such provision should be at a district or regional level.

The chief argument against the status quo is the "unfairness" of abortion provision, district by district and region by region. However, there is evidence that these disparities are lessening, not only from the paper by Fowkes *et al* but also from the Wessex abortion study.¹ A second argument is the different provision between the private sector and the NHS. In particular, single women aged 17-34 are less likely to get NHS abortions and more likely to be aborted in a private clinic. In practice, the private sector offers abortion almost "on demand," whereas the doctor in NHS practice is more likely to be selective. There are two reasons for this: NHS doctors on average tend to interpret the 1967 abortion act less liberally and, secondly, they are constrained by a shortage of resources for all forms of gynaecological care. The Wessex study has identified this fact and shown that districts where resources are less than average tend to do less abortions.

Like mixed sex wards, day care units are now the administrator's dream. Superficially it

seems that here is a way to save money and reduce waiting lists; but in practice day care units are very demanding and complex to run, and many of the savings are spurious.

There are no medical arguments in favour of increasing the number of NHS abortions vis-à-vis private abortions: the arguments are emotional and political. At a time when other types of private medical care are substantially increasing and when NHS waiting lists of all kinds are getting longer, it is difficult to understand why abortion should be singled out for priority care. If more money is spent on abortion units there must be less money for other patients.

Already NHS abortion practice is becoming more even-handed throughout the land. To develop regional units would polarise attitudes and distort the equitable and proper district developments for gynaecological services. It would be better to increase the allocation of money at each district gynaecological unit and encourage current trends.

ANTHONY NOBLE

Royal Hampshire County Hospital,
Winchester

¹ Ashton, J R, *et al*, *Second Report of the Working Group set up by the Wessex Regional Health Authority with the University of Southampton to Study the Provision of Induced Abortion and Abortion Related Services in Wessex*. University of Southampton, July 1978.

SIR,—The recent review of the first decade of experience of the Abortion Act (1967) (27 January, p 217) confirms what we have said in various places since 1971.^{1 2} The decline in mortality and morbidity rates, the use of concurrent sterilisation, and the shorter duration of pregnancy at abortion, particularly within the NHS, are also welcomed by us. However, it is not enough that we should merely accept this situation.

Comparison of the results of abortion obtained by the NHS and by the charitable and private sector shows so clearly how the results and the services could be improved immediately and considerably with almost no additional cost to the NHS. We can only conclude that the major factor responsible for the provision of poor abortion services within the NHS is reluctance on the part of the gynaecologists and health administrators who should be concerned. General practitioners are much more aware of the needs than the specialists, and the public has been unflagging in exerting pressure for the provision of humanly sensitive, safe, early, and efficient abortion.

An ill-informed spokesman for the DHSS made a hasty reply³ to our letter (19 August 1978, p 562) drawing attention to the differential mortality rates for abortion between the NHS and the private sector. We were promised early publication of the facts to rebut our suggestions that the NHS is still carrying out abortions too late, using inappropriate techniques, too often combined with sterilisation, and too often by relatively inexperienced operators. Later the DHSS suggested to us that our observations were a matter for the profession rather than the Department. We fail to understand how the profession can draw conclusions without the facts, which remain unpublished and which the DHSS surely have. As the delay in obtaining this information increases, we feel bound to ask the DHSS once again to provide the public and the profession with information that would enable the quality of our abortion services to be improved. We have personally asked the