repeated after one week. Four of the patients were postmenopausal; the other six were studied during the same phase of the menstrual cycle on both occasions.

Basal plasma prolactin levels were normal in all patients before and during treatment with diazepam. Metoclopramide produced a marked increase in plasma prolactin levels maximal 30 min after injection. An analysis of variance of the log transforms of the plasma prolactin levels showed no significant difference before and during diazepam treatment. The overall trend was also constant at each time before and during diazepam treatment. It is therefore concluded that diazepam does not affect basal prolactin levels or the release of prolactin in response to the dopamine antagonist metoclopramide. This finding compares with earlier reports, which showed that growth hormone production, pituitary-adrenal function,⁵ and thyroid function⁶ are unaffected by diazepam.

J D WILSON

Metabolic Unit,
Royal Victoria Hospital,
Belfast

D J KING

Department of Therapeutics,
Queens University of Belfast, and
Holywell Hospital,
Antrim

B SHERIDAN

Department of Biochemistry,
Royal Victoria Hospital,
Belfast

¹ Sachar, E J, *et al*, *Neuroregulators and Psychiatric Disorders*, p 242. New York, Oxford University Press, 1977.

² Horrobin, D F, *Prolactin: Physiology and Clinical Significance*, p 32. Lancaster, MTP, 1973.

³ Noel, G L, *et al*, *Journal of Clinical Endocrinology and Metabolism*, 1972, **35**, 840.

⁴ Judd, S J, Lazarus, L, and Smythe, G, *Journal of Clinical Endocrinology and Metabolism*, 1976, **43**, 313.

⁵ Havard, C W H, *et al*, *Journal of Endocrinology*, 1972, **52**, 79.

⁶ El-Hazmi, M A F, *Clinica Chimica Acta*, 1975, **63**, 211.

Drug compliance and the elderly patient

SIR,—Recent correspondence and criticism in the press of drugs prescribed for the elderly and the ways in which they are taken have not identified the two most important problems that certainly seem to affect the ways the elderly patients on my list take theirs.

Firstly, the current child-proof container seems also to be elderly-patient-proof as well. Arthritic thumbs make it quite impossible for a substantial minority of my elderly patients to remove the lids from such containers. My own recent investigations of this problem revealed at least six containers of which even I couldn't remove the lid.

Secondly, no more than 45% of 187 elderly patients on my list were able to read the label on the container giving the instructions. The small size of the label and the illegibility of the writing made the job of decipherment difficult. Instructing the chemist to label the containers clearly and in large writing has had no effect. Individual chemists whom I have directly approached tell me that the cost of typing labels would be prohibitive.

Surely the Department of Health and Social Security could recognise that these two problems exist and instruct all chemists to test whether the elderly recipient of the prescription can actually open the container in which the drug has been placed and can actually read the chemist's instructions. These two simple steps would be an excellent starting point in

ensuring compliance. At the moment the situation seems contrived to ensure the opposite.

G H CURTIS JENKINS

Ashford, Middx

Pneumomediastinum in anorexia nervosa

SIR,—We read with interest the report by Drs Anita J Donley and T J Kemple (9 December, p 1604). Recently we saw a similar case in which vomiting seemed an important aetiological factor.

A 16-year-old schoolgirl was referred with primary amenorrhoea, weight loss to 30 kg (height 1.63 m), anorexia, and the psychiatric features of anorexia nervosa. When admitted she was found to have surgical emphysema of the neck, confirmed by chest x-ray, which also showed pneumomediastinum but no pneumothorax. This feature had not been noted at outpatient attendance six days before. A Gastrografin swallow radiograph was normal.

She admitted to feeling nauseated and full after small meals and to one episode of retching and vomiting, perhaps self-induced, following an excessive food intake shortly before she was first seen. In the absence of demonstrable lung disease we feel vomiting to have been the likely precipitating cause of the surgical emphysema and pneumomediastinum.

A P BROOKS

C MARTYN

Metabolic Unit,
Western General Hospital,
Edinburgh

Toilet paper and spread of infection

SIR,—Dr R E W Fisher writes (28 October, p 1233) that "tiny sheets of toilet paper are disgusting and insanitary." It is not quite clear whether he objects to the size or the substance.

Several years ago, while I was working in the Middle East, a Moslem medical student followed me into an Armenian general store and saw me buy some rolls of strange pink paper. "Please, Mr Professor, what are those things?" he asked. "Toilet paper," I said. "What are they for?" he inquired politely. Patiently and rather patronisingly I explained that the sheets were used for personal cleansing after defecation. He was shocked in the extreme. "But, Mr Professor," he said, "in Iran we don't even clean our plates by rubbing them with paper?"

Since that time I have never been quite certain that the West was ahead in hygiene. It is worth noting, however, that the East is not particularly famous for the absence of dysentery.

JOHN H S PETTIT

Kuala Lumpur,
Malaysia

Rapid diagnosis of *Mycoplasma pneumoniae* infection: a reminder

SIR,—A common problem both in general practice and in hospitals, especially at this time of the year, is infection with *Mycoplasma pneumoniae* (MP), particularly involving the chest. This year has seen a markedly increased number of reports of MP infections in the community.¹

Diagnosis is often made in retrospect by serology when the decision about antibiotics has already been taken, perhaps wrongly.

Tetracycline and erythromycin are known to shorten the MP illness but are not ideal antibiotics for other forms of bacterial pneumonia. Ampicillin or amoxycillin, although often effective against the latter, are useless against MP and in fact may precipitate an exanthem in this infection.²

Agglutinins are found in well over 50% of patients with MP pneumonia, but testing for them is not easily performed, especially in general practice. A cold agglutinin screening test (CAST) done at the bedside has been previously reported as detecting titres of greater than 64.³ In the past three weeks we have seen three patients with lobar pneumonia in whom this simple CAST was positive and allowed us immediately to start appropriate treatment for the MP infections that were later confirmed by serology. This test can most simply be done by adding a few drops of blood to the citrate in a commercial prothrombin tube (making a 50/50 mix) and then placing the tube in the ice tray in the freezing compartment of a refrigerator. After a few minutes the cooled tube is removed, tilted, and rotated, when obvious agglutination will be seen on the wall of the tube if the test is positive. This agglutination goes by warming the tube in one's hand. This test is also ideally suited to general practice, as few homes are without a freezing compartment and prothrombin tubes can be carried by the visiting doctor.

We feel that this test deserves more publicity as an aid to quickly identifying this common and treatable infection and thus reducing morbidity and possible hospital admissions.

J T MACFARLANE

IAN A NEALE

Radcliffe Infirmary,
Oxford

¹ Communicable Diseases Surveillance Centre (PHLS), 1978, unpublished.

² Cherry, J D, Hurwitz, E S, and Welliver, R C, *Journal of Pediatrics*, 1975, **87**, 369.

³ Griffin, J F, *Annals of Internal Medicine*, 1969, **70**, 701.

Intermittent ophthalmoplegia

SIR,—Drs W F Kelly and G F Joplin (2 December, p 1534) rightly stress the difficulty in the differential diagnosis of intermittent ophthalmoplegia and obviously in a short report are unable to supply all the clinical data to enable the reader to assess their statement that the patient was clinically euthyroid, although this appears very likely. However, the possibility that the patient was mildly hypothyroid without clinical signs (obviously closely considered by the writers) appears to remain.

This patient had had a previous partial thyroidectomy, has thyroid antibodies, and in addition a borderline high plasma thyrotrophin (TSH) concentration with an exaggerated response to thyrotrophin-releasing hormone. The "normal" levels of T3 and T4 recorded do not exclude the possibility of mild thyroid deficiency since the normal range is a statistical one and it is now accepted that tissue requirements for thyroid hormones vary from person to person. The findings given appear at least compatible with a patient whose thyroid function is compromised and whose T4 level is maintained at a level below the ideal for this patient but still (just) within the population normal range by an increase in TSH activity.

It is a cornerstone of the treatment of ophthalmic thyroid disease that the patient be

kept unequivocally biochemically euthyroid. It is possible (and I feel worthy of serious consideration) that the addition of a small maintenance dose of L-thyroxine might enable the required dose of steroids (which of course also have a valuable role to play) to be reduced to a lower level than 10 mg of prednisone a day with an accompanying lessening of the risk of long-term steroid-induced complications.

J F FALCONER SMITH

Nuffield Department of Clinical
Biochemistry,
Radcliffe Infirmary,
Oxford

Possible environmental hazards of gas cooking

SIR,—Recent letters in your columns from Mr E A K Patrick and Dr M C S Kennedy (19 August, p 567) discussed possible hazards of indoor air pollution from gas stoves. In community studies of respiratory disease in Connecticut and South Carolina^{1,2} in which we obtained data on respiratory symptoms and lung function from about 7000 residents aged 7 years and over we compared residents in homes with gas stoves with those who used electric power for cooking. We took age, sex, race, height, weight, and smoking habits into account in these comparisons and found that respiratory symptoms were equally prevalent and lung function similar among men, women, and children living in homes with gas stoves and in homes with electric stoves. Using portable sampling equipment in selected homes³ we confirmed that nitrogen dioxide (NO₂) concentrations are higher in homes equipped with gas stoves than in those where electricity is used for cooking. Twenty-four-hour average NO₂ levels in 11 homes with gas stoves ranged up to 500 µg/m³, and short-term (2 h) peaks up to 3000 µg/m³ were observed in one such home. However, we have found no evidence that the use of gas for cooking is associated with increased respiratory illness among adults or among children aged 7 years and older. We did not investigate younger children, but any effect of indoor air pollution from gas stoves on their lungs would appear to be temporary, since no changes were detected after the age of 7.

H ROLAND HOSEIN

Department of Preventive Medicine
and Biostatistics,
University of Toronto,
Toronto, Ontario

AREND BOUHUYS

Department of Internal Medicine,
Yale University School of Medicine,
New Haven, Connecticut

- ¹ Mitchell, C A, Schilling, R S F, and Bouhuys, A, *American Journal of Epidemiology*, 1976, **103**, 212.
- ² Bouhuys, A, Beck, G J, and Schoenberg, J B, *Nature*, 1978, **276**, 466.
- ³ Binder, R E, et al, *Archives of Environmental Health*, 1976, **31**, 277.

Ankylosing spondylitis in HLA-B27-positive individuals: use in diagnosis

SIR,—We wish to take issue with several of the major points in your leading article (2 September, p 650) "HLA-B27 and risk of ankylosing spondylitis." You recognise the obvious conflict between the early and the more recent population studies of the prevalence of ankylosing spondylitis (AS) but, for reasons not given, prefer to believe the more recent

studies. In order to resolve this conflict we have recently conducted a study of the population of Busselton, Western Australia, and have found a prevalence of AS similar to that predicted by the early studies rather than the much higher prevalence suggested by Calin and Fries.¹ We HLA-typed some 80% of the adult population of this relatively isolated, stable, small country town. Of 2745 people, 168 (6.1%) were B27-positive. Careful questionnaire and clinical assessment was undertaken on 139 of these together with an age- and sex-matched control group from the same population. Radiographs, which were available from 22% of the B27-positive individuals, were reviewed. Not one case of AS was detected and there were no differences in the frequency and character of the back pain between the two groups. We used a questionnaire similar to that of Calin and Fries and obtained sacroiliac radiographs in a higher percentage of the study group. These data suggest a maximum prevalence of AS of 0.04% in the population as a whole and 0.7% in B27-positive individuals and are therefore consistent with the earlier "classical" surveys rather than some surveys¹ which may be explained either on the basis of the population studied or over-interpretation of symptoms, signs, and x-rays.

Your article states that in a suspicious clinical setting (for example, persistent backache) the presence of B27 will help in making a diagnosis. We contend that it is the absence of B27 which should be used to exclude AS, while the finding of B27 in a patient is of little help. Such a contention is based on the relative rarity of AS and the relative frequency of B27. Even if one accepts the figures from the study of Calin and Fries four out of five people with B27 will not have AS. Furthermore, recent studies² indicate that the incidence of B27-negative AS may be as low as 1-2%. Therefore it is clear that HLA-typing for B27 should be used as an excluding test rather than a confirming test.

Because the prevalence of AS in B27-positive individuals is nearer 1% than 100% genetic counselling will cause needless anxiety. In fact, with modern treatment, as outlined in your article, AS is nowadays an infrequent cause of significant morbidity.

F CHRISTIANSEN
PAUL ZILKO
R L DAWKINS

Department of Clinical Immunology,
Royal Perth Hospital,
Perth, Western Australia

- ¹ Calin, A, and Fries, J F, *New England Journal of Medicine*, 1975, **293**, 835.
- ² Joint Report on HLA and Disease, Seventh International Histocompatibility Workshop Conference, Oxford, 1977. In press.

* * * These new data from Dr Christiansen and his colleagues are of considerable importance and it will be of great interest to read a full report, when the size and age structure of the population and its ethnic origins will be clearer. The great variations which are known to exist in the frequency with which AS is found in different populations must be taken into account and it may be that a small, relatively isolated country town in Western Australia is not typical of the generality of Caucasians. Certainly the observations made by Calin and Fries are similar to those of Kidd *et al*¹ in suggesting a much higher frequency of clinical and subclinical AS in B27-positive individuals than Dr Christiansen has found. No doubt more studies will be completed and the

mystery resolved in time. We agree with Dr Christiansen that genetic counselling for ankylosing spondylitis should be approached with caution, and this was stated in the leading article. Also consistent with the leader is Dr Christiansen's point that the absence of B27 might be helpful in diagnosis.—Ed, *BMJ*.

¹ Kidd, K K, et al, in *HLA and Disease*, ed J Dausset and A Sveigaard, p 72, 1977.

Another hazard of pierced ears

SIR,—I should like to draw readers' attention to another hazard of the present fashion for pierced ears. Three children from the same family attended our casualty department 24 h after having their ears pierced at a reputable Sheffield jewellers. They had all received ethyl chloride spray to anaesthetise their ear lobes before piercing.

However, instead of the usual 20-30 seconds' worth of spray, each received several minutes of spraying to each lobe (the mother thought about seven minutes in one case). All the children had deep purple areas of chemical "frost bite" from their ears in a thick area 2 in (5 cm) wide across their necks running to the midline of their throats. These burns have subsequently completely healed with conservative treatment without leaving a scar.

D A NOBLE

Accident and Emergency Department,
Children's Hospital,
Sheffield

Polymyalgia rheumatica and primary biliary cirrhosis

SIR,—In reply to the comments of Drs A L Ogilvie and P J Toghill (25 November, p 1501) on our case reports (21 October, p 1128), we are aware that hepatic biochemical abnormalities occur in polymyalgia rheumatica (PMR) and, indeed, not uncommonly in the investigated population. However, the abnormalities we found were gross, consistently so, and were indicative of hepatic disease. The antimitochondrial antibody titres were high and remained so in our unbiopsied patients. Our patients did not have serositis, fever, or peripheral arthritis or arthralgia. Therefore we cannot agree that the systemic lupus erythematosus-like syndrome they mention should enter in the differential diagnosis.

Liver biopsy was not performed on two patients. One was a recluse and it reflects credit on her daughters and her general practitioner, Dr Nigel Shield, that we were able to investigate her as far as we did. A biopsy would not have altered management and neither would it have done so in our first case, in which it would probably not have contributed to diagnosis either. We believe that careful and repeated biochemical examination is sufficient to establish a probable diagnosis of chronic hepatitis in appropriate clinical circumstances. Walker *et al*¹ noted that in all of their biopsied cases the results confirmed the biochemical findings of significant hepatic abnormality. We also note that one of their patients, a doctor, refused liver biopsy because there were no hepatic symptoms.

With regard to Drs Ogilvie and Toghill's comments concerning our mention of a first case report of PMR and chronic hepatitis, we are surprised at their use of the word "categorically." Our reading of the text was that