of the aorta was found but no atheroma of the coronary arteries. Another of the eight patients, a woman aged 44, had a serum cholesterol concentration of 7 mmol/l (273 mg/100 ml) before thyroidectomy and a concentration of 6 mmol/l (230 mg/100 ml) before she died four years later.

These eight patients are as relevant to the debate on fat and atheroma as the flowers that bloom in the spring. It is time that this outworn reference was given a well-deserved rest.

P B S FOWLER

Charing Cross Hospital (Fulham),
London W6

¹ Blumgart, H L, Freedberg, A S, and Kurland, G S, *American Journal of Medicine*, 1953, 14, 665.

² McMichael, J, *Lancet*, 1974, 1, 1340.

³ Thompson, G, *Lancet*, 1974, 2, 102.

SIR,—With reference to the last comments in Drs K G Taylor and D J Betteridge's letter on "Fats and atheroma: an inquest" (17 February, p 485), I wish to point out that dietary limitation of cholesterol-containing foods is unnecessary and ineffective in reducing serum cholesterol. Over four-fifths of the body's cholesterol is endogenous and dietary cholesterol intake is compensated for anyway.¹ Restricting eggs to two per week is bad dietary practice—they are extremely nutritious, cheap, readily available, and easy to store and cook; at most I might consider a limitation of four per week. Reduction of milk would be unwise: research has shown that it has hypocholesterolaemic properties.² Limiting total fat and refined carbohydrates in the diet are the all-important factors.

F E DOBSON

Dietitian's Office,
Bromsgrove General Hospital,
Bromsgrove, Worcs

¹ Davidson, S, et al, *Human Nutrition and Dietetics*. Edinburgh, Churchill Livingstone, 1975.

² Howard, A N, and Marks, J, *Lancet*, 1977, 2, 255.

SIR,—I would like to deal with some of the supposed evidence in favour of an aetiological significance of cholesterol in atheroma, such as that quoted by your correspondents Drs K G Taylor and D J Betteridge (17 February, p 485).

Evidence from animal experiments is always a little suspect, but it is very much more so when vegetarian animals are made to take foods they never normally eat. Suppose we reversed the experiments and fed human beings the strict vegan diet of rabbits and rhesus monkeys. Such persons would develop muscular weakness and probably B₁₂ deficiency. This could be cured by giving essential amino-acids (first-class proteins) and vitamin B₁₂. Would Drs Taylor and Betteridge regard this as evidence that all rabbits and rhesus monkeys should have supplements of essential amino-acids and B₁₂?

Drs Taylor and Betteridge remind us that the probability that an individual will develop a coronary infarct increases in proportion to his serum cholesterol concentration. In myxoedema this relationship is not just a probability. The serum cholesterol is a direct measure of the severity of the disease. Do they suggest that this means that cholesterol causes myxoedema or it can be cured by lowering the cholesterol with polyunsaturated fats?

Most experiments purporting to show an improvement with polyunsaturated fats have been criticised because they do not exclude

other factors such as smoking, obesity, and hypertension (Professor J Yudkin would add sucrose), which by themselves or together may have been responsible for the result. While I cannot claim to have studied them all, it is reasonable to expect that after years of what can only be called determined research there would be little room left for argument if the significance of cholesterol in atheroma were anything like that of cigarette smoking in cancer of the lung.

Finally, the cholesterol of premenopausal women is higher than that of men, yet their incidence of coronary heart disease is lower. Would Drs Taylor and Betteridge recommend that men under 45 should eat more butter, cream, and eggs in the hope of lowering their incidence of coronary heart disease to that of premenopausal women.

We are all agreed on the desirability of a "sensible diet" but that this diet should deprive us of most of our eggs, butter, and cream with substitution by polyunsaturated fats is far from having been demonstrated. Even the plea that it may do good and cannot do harm is no longer tenable. Sir John McMichael has done well to ask for an inquest. I hope you will call on him to reply to any further criticisms from the peculiar entrenched form of logic which seems to have overtaken so much of the profession.

In the meantime may I (not for the first time) suggest that "a little of what you fancy does you good"—at least until it is proved to do otherwise.

A LEWIS

London, W1

Malnutrition in infants receiving cult diets

SIR,—We have read with interest and not a little concern the paper "Malnutrition in infants receiving cult diets" by Dr I F Roberts and others (3 February, p 296), which has excited considerable publicity and comment in the news media. We are concerned about the inaccurate references to diets in this paper and the unwarranted adverse comments about acceptable diets that have ensued.

The authors of this article have clearly confused different diets based on the sole consumption of plant foods. The result is that, while attention is quite rightly drawn to the serious consequences of feeding certain extremely limiting diets, the confusion about the nomenclature of diets is likely to cause unnecessary concern to parents who for a variety of reasons wish to feed their children a diet based on plant foods.

We saw the 13-month-old boy described in this paper at Kingston Hospital with the late Dr Frey Ellis before he went to the Hospital for Sick Children. This child was receiving a "fruitarian" and not a vegan diet. Indeed, the father of this child persistently refused to accept expert advice on a balanced vegan diet including the use of a soya-based milk and supplements of vitamin B₁₂. The other three infants described in the paper are not known to us, but it is evident that they were receiving macrobiotic diets. These are known to be dangerous,¹ and should be clearly distinguished from vegetarian and vegan diets.² The failure to distinguish between fruitarian and vegan diets has resulted in the statement that "Vegan and the more extreme macrobiotic diets are obviously inadequate for growing children. . . ." We agree entirely about the fruitarian and

macrobiotic diets, but cannot agree about a balanced vegan diet for we know that children have been, and are being, reared very successfully on such a regimen. Mothers who wish to feed their children on a vegan diet can obtain free advice and information from the Vegan Society, 47 Highlands Road, Leatherhead, Surrey.

J W T DICKERSON

Division of Nutrition and Food Science,
Department of Biochemistry,
University of Surrey,
Guildford, Surrey

ANN M FEHILY

Department of Haematology,
Kingston Hospital,
Kingston, Surrey

¹ American Medical Association Council on Foods and Nutrition, *Journal of American Medical Association*, 1971, 218, 397.

² Dickerson, J W T, and Fehily, A M, *Practitioner*, 1979, in press.

SIR,—Nutrient deficiency is more likely to occur if the variety of food in the diet is restricted, especially so when the diet is devoid of foods of animal origin. Even so, a properly constructed vegan diet, composed of a mixture of cereals, pulses, nuts, fruit, and vegetables, can meet all the requirements for nutrients provided that it is supplemented with vitamins B₁₂ and D.^{1 2}

Most vegan mothers breast-feed their babies well into the second year of life. Breast milk can meet all the nutrient requirements for the first six months of life provided that the supply is adequate. In the 20 vegan mothers we have studied the milk composition and yield, when estimated by test weighing of the infants, were normal; the milk was richer in linoleic acid than that from omnivores.^{2 3} Breast milk will continue to meet the nutrient requirements of the infant after six months provided that it is supplemented with a suitable weaning food such as Farex, which consists of a mixture of cereal and soy-bean flour. The infant can then gradually be introduced to a properly balanced vegan diet with suitable vitamin supplements. Over the past 15 years my colleagues²⁻⁴ and I have studied about 60 children born of vegan mothers and reared on vegan diets. Their development, as judged by their rate of growth and general health, seems to be quite normal.

We were involved with the first case described in the paper "Malnutrition in infants receiving cult diets: a form of child abuse" by Dr I F Roberts and others (3 February, p 296). The parents of this child were following a "fruitarian" diet, which excludes cereals and pulses as well as food of animal origin, and were attempting to wean the child on a similar diet. The authors, who omitted to mention that the child was eventually rehabilitated on a properly balanced vegan diet, incorrectly concluded that "Vegan . . . diets are obviously inadequate for growing children, with deficiencies of total energy, suitable protein, fat-soluble vitamins, and some minerals."

One of the most disturbing aspects of this paper was the unscientific manner in which it was presented. Assessment of nutrient intake was scanty; why one child developed kwashiorkor and another marasmus was not discussed. A more accurate assessment of nutritional state might have revealed that the children were being virtually starved. If this were the case then qualitative differences in diet become of secondary importance. The