casualty services. It will be a matter for local negotiation with health authorities to ensure that casualty services are available in all areas, and where these have to be provided by general practitioners they must be adequately paid for the work undertaken."—ED, *BMJ*.

Next year's pay

SIR,—It appears from several articles in the paramedical press that Dr R A Keable-Elliott is attempting to persuade the Government to give us the whole of the outstanding 20% increase in remuneration, including the third phase 10%, due in 1980. I hope he realises that this will cause a shortfall of somewhere around 13% on 1 April next year. The excuse given is that our negotiators are worried lest the Government impose a wage freeze. Surely this cannot apply to agreements already made by the Government. I fail to see why our negotiators have been panicked into such a manoeuvre. I personally would be quite happy to take my 10%, plus whatever additions the Review Body sees fit to give us this April and await next year's 10%, plus an element for inflation. If I was negotiating on behalf of the Government I would seize Dr Keable-Elliott's suggestion with both hands.

I would guess we will pick up somewhere around 22 or 23% on 1 April this year if everyone has done their homework properly. If inflation continues at its present level then a similar amount would be expected next year. Faith in the BMA among its members must have been badly shaken by the debacle of the clinical assistants' arbitration. Now I find Dr Keable-Elliott giving away a substantial part of my future income without so much as having discussed whether or not I would agree. Just whose side is he on?

MICHAEL WYNN

Buckie, Aberdeen

Short-term medical certification

SIR,—Those who listened to a recent edition of *Any Questions?* (Radio 4, 2 February) will have heard Mr Arthur Scargill say that in his opinion miners should be paid more than doctors. I do not share his view, but as a GP in a mining community I can understand how some might do so.

It is a lamentable but incontrovertible fact that a considerable part of my work could be done equally well by my receptionist, and my patients know it. I refer to the issuing of short-term medical certificates, a chore which in many cases requires no medical knowledge or skill whatsoever. It is obvious that in some practices this is far less of a burden than in others, but in a mining practice such as mine (I am in a partnership of three with just under 9000 patients) up to 20 "consultations" a day per partner are for the purpose of certification for "illnesses" which can be neither confirmed nor excluded by clinical examination. The arguments put forward in the DHSS letter published in this week's *BMJ* (17 February, p 501) are purely those of expediency and should not, I submit, be accepted by the profession. I cannot see that increased use of the form Med 5 would help appreciably.

I make no pretence to high academic ability, but I bitterly resent the fact that in much of my work my medical training is quite superfluous, and that a section of my patients judge me

not by my clinical ability, such as it is, but by the ease with which I can be "conned." It is surely quite illogical to advocate, as do the proponents of the new "charter," the training of more GPs, and the reduction in list sizes, without first ensuring that those already in practice do not have to waste so much of their time on such menial tasks.

The continuation of the present farcical system of certification, while no doubt very convenient for the DHSS, is resulting in the devaluation of general practice as a profession in the eyes of a significant section of the community, and Mr Scargill's statement is but a sign of this devaluation. This consideration must surely transcend all others in any further negotiations with the Department.

E W STURTON

Worksop, Notts

Normansfield and the NHS

SIR,—Dr T L Pilkington (3 February, p 348) appears to think that because psychiatrists have moved around and surgeons have not the former are more enlightened, although he does not give any reason why this should be so, whereas one could think of a number of explanations for this activity.

Unfortunately, only part of my letter (13 January, p 131) was published in the *BMJ*. Surgeons and those in other specialties usually have a rather longer course of training before they achieve consultant status than do psychiatrists, and most of them have moved many times by the time they achieve their final appointment and are older because of the increased length of training required before they are appointed. Perhaps, therefore, psychiatrists accept the change more willingly. To add to their troubles, those who practise surgery, medicine, and often even general practice find themselves moved about because of the rather bureaucratically imposed regulations from the royal colleges and other examining bodies and the need to have a much wider experience before they achieve the ultimate senior appointment. One would point out, however, that one of the reasons that the armed Forces find it difficult to recruit staff is the domestic insecurity which is involved and that the population at large thinks that it is entitled to have work brought to the workers rather than have the worker move his home. Higher appointments are expected to be very generously rewarded under the terms of disturbance which precede appointment. In my own case I was greatly discouraged from my family's choice of career in the Foreign Service by this very fact. If consultants are to be moved around in this fashion then I am sure the specialties will rapidly lose their popularity and an increasingly apparent shortage in some fields will reach a crisis level.

GEORGE T WATTS

General Hospital,
Birmingham 4

Careers for part-time doctors

SIR,—I have read with great interest the Talking Points (13 January, p 144 and 3 February, p 359) on "Revised career structure: first priority" and "The new consultant contract and women senior registrars". I have been considering the problems of doctors in non-

training grades who wish to work on a sessional basis in NHS hospitals and would like to proffer the following ideas for discussion.

At present the main categories of doctors in non-training hospital posts are consultants, hospital practitioners, medical assistants, and clinical assistants. As many medical assistants are full time my suggestions are confined to the other three categories. Consultants have a defined service commitment with ultimate responsibility for their patients, and need to have satisfactorily completed an approved course of training with a recognised higher qualification. Hospital practitioners are principals in general practice with a defined period of experience in the hospital work they undertake. Clinical assistants comprise the rest—those who either have not completed a full vocational training course or are unable to offer enough notional half days to be appointed as a consultant. No allowance is made for higher qualifications or experience, prospects for promotion are limited, and the salary scale has no increments.

Part-time doctors are unhappy at the lack of a defined mechanism for employment; they do not fit into the present system and special arrangements have to be made to employ them as individuals as they are frequently tied to a geographical area.

What is needed is a combination of the hospital practitioner and clinical assistant grades and the elimination of the principal in general practice qualification. There should be a pool of sessions that can be applied for and the applicant should be graded according to experience and higher qualifications and put at the appropriate point on an incremental scale. It is a matter for discussion whether the sessions should be funded nationally or by area health authorities, as there will always be the problem of a well-qualified doctor forced to move from one area to another because of personal and not professional reasons—and also the local service requirements have to be considered.

The name of the post needs careful consideration so that it neither implies a permanent subordinate status nor carries emotional overtones such as those evoked by the terms junior and senior hospital medical officer. Hospital practitioner is an inoffensive title if the restriction to principals in general practice is removed.

The number of doctors wishing to provide a service to the NHS on a less than full-time basis is likely to increase with the well-documented rise in female medical students. Although many of them should be able to complete their vocational training and become successful consultants provision should be made for those who are unable to fit into the present system.

MARGARET ELMES

Department of Pathology,
Welsh National School of Medicine,
Cardiff

Correction

Miniature peak-flow meters

The second sentence of the second paragraph of this letter by Dr B M Wright (17 February, p 489) should have read "In addition a correlation coefficient of 0.902 with the standard meter compared with 0.990 for the mini-meter, means that it is only about one-third as precise as the latter".