We believe that the most dramatic effect of preoperative and postoperative 1 α HCC in primary hyperparathyroidism is in the management of those few patients with really severe bone disease whose postoperative management in previous times remained stormy over several months. We have been accustomed to seeing about one such patient per year at Glasgow Royal Infirmary and believe that at least five of our 15 patients mentioned above would have fallen in this category (all had serum alkaline phosphatase and parathyroid hormone levels (where measured) some 10 times higher than the upper limits of normal). Such patients may not be encountered in Birmingham but still unfortunately turn up in Glasgow.

A body of evidence now exists³⁻⁵ to suggest that vitamin D deficiency may play a role in the genesis of the bone disease in these severe cases. On the one hand the vitamin D deficiency may cause an increase in parathyroid hormone secretion, while on the other hand the primary hyperparathyroidism may cause an increased requirement for vitamin D, possibly through the increased renal turnover of 25-hydroxy vitamin D₃. On this basis one might anticipate some amelioration of the clinical problem by giving 1 α HCC before operation, and in a few cases we have indeed demonstrated some reduction in the serum parathyroid hormone level. In addition, the prophylaxis of paraesthesiae and tetany, even in the less severe cases, seems preferable to awaiting their postoperative appearance, and preoperative administration of 1 α HCC for three or four days, in addition to continuing it postoperatively for a similar period, usually achieves this aim as well as probably contributing to bone healing.

Our present treatment schedule for 1 α HCC administration in primary hyperparathyroidism varies with the severity of the bone disease—1 or 2 μ g daily for five to 10 days preoperatively followed by 1 or 2 μ g daily for five to 50 days postoperatively.

As Dr Heath's series demonstrates, one must monitor serum calcium daily to identify any idiosyncratic sensitivity to 1 α HCC. It would be a great pity if the occurrence of hypercalcaemia in the one patient were to deprive the group with severe bone disease of the potential benefits of this therapy.

I T BOYLE
I FOGELMAN
G H BEASTALL
S JOFFE
B BOYCE

University Departments of Medicine,
Pathology, Clinical Biochemistry, and Surgery,
Glasgow Royal Infirmary,
Glasgow

- ¹ Boyle, I T, *et al*, *Clinical Endocrinology*, 1977, 7, 215S.
- ² Nordin, B E C, *Metabolic Bone and Stone Disease*, p105, Edinburgh, Churchill Livingstone, 1973.
- ³ Woodhouse, N J Y, *et al*, *Lancet*, 1971, 11, 283.
- ⁴ Stanbury, S W, *Clinical Endocrinology and Metabolism*, 1972, 1, 239.
- ⁵ Mawer, E B, *et al*, *Clinical Science and Molecular Medicine*, 1975, 48, 349.

Diet, sunlight, and 25-hydroxy vitamin D

SIR,—We read with interest the article by Dr E M E Poskitt and others (27 January, p 221) in which they discuss possible mechanisms for physiological control of circulating 25-hydroxy vitamin D (25-OHD). Bhattacharya and De Luca¹ showed evidence of suppression of hepatic D₃-25-hydroxylase in the rat liver and Haddad *et al*² proposed the same mechanism in humans.

We have reported previously on the use of vitamin D supplements and ultraviolet light in old age.³⁻⁴ In 15 subjects on supplements there was, after one month of treatment, a strong negative correlation between the initial plasma 25-OHD levels and the change in this ($r = -0.484$). In 12 subjects treated with ultraviolet light for one month there was a

similar association between the two variables ($r = -0.589$).

We would suggest that this provides further evidence of product inhibition in the control of hepatic D₃-25-hydroxylase in humans. It should be noted that the regulatory system appears to operate only for physiological amounts of vitamin D. The toxic effects from large doses of vitamin D are well recognised.

A P J SNELL

Department of Geriatric Medicine,
University of Manchester

W J MACLENNAN

Department of Geriatric Medicine,
University of Southampton

J HAMILTON

University of Southampton

- ¹ Bhattacharya, M H, and De Luca, H F, *Journal of Biological Chemistry*, 1973, 248, 2969.
- ² Haddad, J F, and Stamp, T C B, *American Journal of Medicine*, 1974, 57, 57.
- ³ MacLennan, W J, and Hamilton, J C, *British Medical Journal*, 1977, 2, 859.
- ⁴ Snell, A P, MacLennan, W J, and Hamilton, J C, *Age and Ageing*, 1978, 7, 225.

Registers for the prevention of genetic disease

SIR,—We read your leading article (27 January, p 215) with interest, and share your concern about confidentiality of genetic registers and the apprehension about "authoritarian intervention in this area." You very aptly stressed the great importance of support for affected families to whom we are the harbingers of bad tidings, saying that our "Genetic counselling should be a humane and cost-effective way of preventing intractable disease."

In the area of Huntington's disease, we wish to point out that although no preventive measures are currently available a register of affected families should be maintained for both research and a service commitment. In our seven years' experience of conducting a Huntington's disease clinic we have found in the register an invaluable clinical tool in the diagnosis and management of this disorder. Your argument that "some heterozygotes may take no notice of genetic counselling: they tend to be defiant towards the disorder, often refusing to let it interfere with their lives or their procreation" takes no real account of the empathetic process of genetic counselling and belies your advocacy of support for these families, as well as being contrary to the principle of families' affliction with severe genetic disorders. There has been a growth of public and family concern in this field. The concept of cost-effectiveness should be viewed in terms of alleviation of human suffering and assisting families to cope constructively, rather than just reducing the prevalence of the disorder in the future.

A properly maintained register, we have found, not only assists in the diagnosis of the disease by clinicians: it also provides valuable data in the planning of management now and in the future. It has provided us with a view of the family network system, from which we can begin to plan to make full use of family resources by support for the ill members of the family. The marshalling of supportive resources in the nuclear and extended family network is an integral part of counselling services for Huntington's disease families.¹

We therefore support the establishment of genetic registers for Huntington's disease by specialised units, with built-in, stringent safe-

guards in confidentiality and accountability, for both research and service commitments.

EDMOND CHIU
BETTY TELTSCHER
BRIAN DAVIES

Huntington's Disease Clinic,
Department of Psychiatry,
University of Melbourne, Australia

- ¹ Chiu, E, and Teltscher, B, *Medical Journal of Australia*, 1978, 2, 394.

Management of refractory oedema

SIR,—Dr J McCormack writes on the trick of combining a thiazide diuretic with a loop diuretic (10 February, p 412). I have also found this combination very useful (and effective in producing hypokalaemia).

Trials with combinations of diuretics made by Olesen and Sigurd have demonstrated the very impressive natriuresis and diuresis achieved when a small dose of bendrofluzide is added to frusemide and spironolactone, or bumetanide.^{1,2} For example, in patients with congestive heart failure given 3 g sodium chloride in their diet and a fluid intake of 1500 ml, bumetanide, 4 mg, and spironolactone, 100 mg, gave a natriuresis of 81 ± 13 mmol(mEq)/24 hours and a diuresis of 1628 ± 131 ml.² Doubling the dose of bumetanide did not increase diuresis. In contrast, combining bendrofluzide, 5 mg, increased natriuresis and diuresis significantly to 209 ± 50 mmol and 2533 ± 437 ml/24 h.

This effect is ascribed to loop diuretics supplying ample sodium at the distal sites of the nephron, inducing an accelerated reabsorption at these sites—which is effectively depressed by a thiazide.

PER FRANZÉN

Mjölbolsta Hospital,
Mjölbolsta, Finland

- ¹ Olesen, K H, and Sigurd, B, *Acta Medica Scandinavica*, 1971, 190, 233.
- ² Sigurd, B, *et al*, *American Heart Journal*, 1975, 89, 163.

Tuberculin testing

SIR,—Dr H G Calwell's observation (3 February, p 345) on the use of Heaf testing as an alternative to tine and Mantoux tests is noted with interest.

After 36 years' experience of a variety of tuberculin tests in chest clinics and hospital staff health departments (approximately 700 a year) I agree that the multiple puncture (Heaf test) is the method of choice. It is quick, painless, and easily administered even in struggling babies. If accurately used it is consistently reliable in its interpretation, differentiating clearly between non-specific (grade 1), post-BCG-positive, acquired immunity (grade 2), and possible occult infection.

The article by Drs J A Lunn and A J Johnson (3 June 1978, p 1451) and subsequent letters have indicated that the tine test produces false-negative results but have not clarified the reasons for this. In my experience the tines are often blunted at the tips and are not always at right angles to the base. Very firm pressure is therefore necessary to obtain adequate skin puncture. In a small series of 35 routine tuberculin tests on hospital staff, I used Heaf and tine tests on opposite arms. The comparative results were identical in every case. In some tests I purposely induced excessive bleeding from the tine punctures to exclude "washing out" the purified protein

derivative as being a cause for false negatives. The results were still identical.

It appears to be all a matter of technique—that is, firm pressure, careful sitting on the ulnar aspect of the mid forearm, adequate stretching of the skin, and checking that the tines have produced satisfactory puncture holes. However, I still prefer the Heaf test for routine screening of contacts and hospital staff as tine tests have certainly shown false negatives when used by the uninitiated.

J M PLATTS

General Hospital,
Newmarket, and
West Suffolk General Hospital,
Bury St Edmunds, Suffolk

Recording of blood pressure readings

SIR,—There seems to be increasing confusion over the recording of the diastolic blood pressure, with some physicians recording the fourth sound and others the fifth sound. Would it not be possible for all blood pressure readings to be recorded with all three readings on each occasion? Thus instead of writing 180/90 we would write 180/90/90 if the fourth and fifth sound occurred together, but if the fifth sound was at a lower level it would be recorded as 180/90/60.

If only the fourth sound were measured the physician would record 180/90/—, indicating non-measurement of the fifth sound; and likewise if only the fifth sound were measured he could record 180/—/60. With this simple technique in recording, consecutive blood pressure readings by different physicians would have greater significance than at present. Perhaps a new name should be coined for the fifth sound to further avoid confusion. Any suggestions?

M G SHELDON

Banbury, Oxon

Medical manpower in the year 2000

SIR,—Your leading article (3 March, p 573) concludes that the DHSS study¹ and our own study² indicate that the primary requirement for more sophisticated medical manpower forecasting is that the quantity and quality of data are improved. As is clear from our work, I agree very strongly with this conclusion. We have little idea about the level of women doctors' participation in the labour force (except for Dr Berenice Beaumont's excellent efforts). We know little about the career intentions of foreign doctors, although the Policy Studies Institute has recently completed some work on this for DHSS. We could "model" UK emigrant doctor flows but the DHSS does not bother. The manipulation of the data with regard to aspects of regional and career structure is to date very limited. Little account is taken of the RAWP report, *Priorities for Health and Personal Social Services in England, The Way Forward*, or other public policy pronouncements in DHSS forecasts. Much of the official forecasting seems to be blinkered and myopic.

We have had all these arguments before, in the Goodenough, Willink, and Todd reports of 1944, 1957, and 1968. This time round I hope that the DHSS will decide to (i) collect more and better data series on a regular basis; (ii) use this improved data stock to produce forecasts on a yearly or biannual basis for discussion with the profession; (iii) extend their forecasts to include regional and career

structure problems; (iv) integrate the forecasting of doctor manpower with the forecasting of nurse and other health personnel manpower; and (v) cost these forecasts in a detailed and sensible manner.

Forecasting will always be difficult but more sophisticated techniques could be used to manipulate the existing stock of data (and, one hopes, in the future a better stock of data). Clearly the inadequacies of the data stock and its manipulation by the DHSS in the recent past indicates that, implicitly or explicitly, the Department views such policy analysis as unimportant. Each doctor costs (according to DHSS estimates whose basis would incense any undergraduate economist by its crudity) £40 000 to produce and the medical schools are swallowing some of the brightest "person power" in the country. More careful data collection and forecasting might make us better able to use these scarce resources more prudently than we do at present.

If the DHSS fails to invest resources in forecasting perhaps a cost-effective allocation of BMA resources would be for the profession itself to collect and manipulate its own medical manpower data. Competing forecasts might be an efficient way of improving the quality of forecasts and the efficiency of resource utilisation.

A MAYNARD

Department of Economics and Related
Studies,
University of York,
York YO1 5DD

¹ Department of Health and Social Security (1978), *Medical Manpower—The Next Twenty Years*. A discussion paper. 1978.

² Maynard, A, and Walker, A, *Doctor Manpower 1975-2000: Alternative Forecasts and their Resource Implications*. Report for the Royal Commission on the National Health Service, Research Paper No 4. London, HMSO, 1978.

SIR,—Medical manpower is the subject of your leading article (3 March, p 573) and also a short paper (p 595). I have recently had to make a careful study of the DHSS paper on this subject and there are important points which are not brought out in either your editorial or Dr James Parkhouse's article.

The DHSS paper contains a great deal of valuable information and does not in my opinion show evidence of the hasty production which you infer. That aside, it does lead to the inescapable conclusion that it is impossible to calculate the number of doctors required in the next 20 years, for the following reasons. (1) There is no way of knowing what the population will be. (2) There is no way of knowing the number of foreign graduates that will be working to prop up the NHS. (3) Geographical maldistribution of doctors influences regional requirements. (4) Though the number of women doctors can be controlled, the amount of employment that they seek cannot.

Given these imponderables, any attempt to enumerate the doctors needed between now and the end of the century, which is the hope of the DHSS paper, is no better than a guess. Willink was wrong in 1957, Todd was wrong a decade later. The only concept that seems important to me is the doctor: population ratio; the current figure of roughly 1500 doctors per million population is certainly enough for our foreseeable requirements, provided good use is made of them; it is deployment rather than pure numbers that really matters, and I believe that our negotiators should hammer home the following six points to the DHSS.

(1) It is essential to improve the support services for family doctors, both staff and

physical facilities, so that doctors can make more efficient use of their time. (2) Domiciliary care of the sick requires far greater finance than it is receiving. (3) Because a GP spends so much time on social problems we should be exploring methods of training lay people as social counsellors, to release the GP for the more important clinical matters for which he is after all specially trained. (4) It seems likely that by 1990 50% of graduates will be women; it is therefore vital to provide adequate part-time career opportunities in the NHS. (5) It would be politic and prudent to try to reduce our dependence on foreign graduates, who currently account for 27% of NHS medical staff. (6) Surely the means are available to keep annual counts of the number of doctors on the NHS payroll and relate this to the population, so that any deviation from the appropriate doctor: population ratio can be detected promptly and acted on.

R M MILNE

Dechmont, West Lothian

Hazards at work

SIR,—I was most interested in the efforts of Dr B J Boucher and her committee to promote health and safety at work at a district level (3 March, p 599), particularly with regard to her comments about the lack of direct guidance given by the DHSS.

Perhaps correspondence in these columns could help to make clear how much we should rely on our own safety codes at district level and how much we can assume that accepted NHS practice automatically includes an adequate degree of safety cover for staff and patients. For instance, it would be reasonable to expect that equipment offered to hospitals in bulk would already have been vetted centrally by the NHS for possible hazards to users. That this is not a safe assumption is illustrated by the following example. During the recent industrial dispute an emergency supply of disposable paper surgeons' suits was obtained to meet a possible laundry shortage. These suits seemed a fire hazard and on testing I found that they were quite easy to ignite with a match, lighter, or cautery-probe. They burned vigorously, giving off dense, acrid smoke; and the material fused, shrank, and clung to surfaces in a frightening manner. The flames were difficult to extinguish with water, no doubt because the material had been partially waterproofed. While smokers pose the greatest risk, to themselves and others, in these suits any wearer would be at a serious disadvantage should an accident arise involving fire in the operating theatre.

Subsequently, I tested an array of disposable paper items for inflammability and found that surgeons' caps and masks, bed sheets, pillow cases, and triangular bandages also burned readily, with the production of choking fumes. In future these items should be over-stamped with a caution about inflammability by the manufacturers.

J G HANNINGTON-KIFF

Frimley Park Hospital,
Frimley, Surrey GU16 5UJ

The new consultant contract

SIR,—I was surprised to read the letter in your correspondence columns (17 March, p 755) from my friends at Northwick Park. It would seem a little late to attack the principles underlying the new consultant