CORRESPONDENCE

Experience in a child health clinic P R Williams, MRCP; B L Anderson, MB; Linda Downing, QN; Pat Savill, SCM, and Evelyn Pope, SRN.....	Differences between Leeds fractures and London fractures? R Wootton, PHD, and J Reeve, DM.....	Service for psychiatrically ill doctors? K N Hambly, MRCP.....
Upper gastrointestinal endoscopy V Moshakis, FRCS.....	Hypnosis H K Gooding, MB; E W Rees, FRCPsych.....	Clinical practice and epidemiology G C Hancock, MB.....
What shall we teach undergraduates? P A B Raffle, FRCP; R A Shinton, BA; Cicely D Williams, FRCP.....	Restrictions on NHS prescribing in Australia D de Souza, FRCPed; R C Hall.....	Difficulty in stopping lithium prophylaxis? M Yuce, MD.....
Fats and atheroma Sir George Pickering, FRCP, FRS.....	Detecting and managing hypertension B S Smith, FRCP.....	"Doctors' orders" I Capstick, FRCP.....
Myocardial infarction imaging P J Bourdillon, MRCP.....	Neglected indexes A S Thorley, FRCPsych.....	Keeping down the elephants H Weisl, FRCS.....
Detection of deep venous thrombosis by ^{99m}Tc-labelled red-cell scans D A P D'Auria, MD.....	Action on alcohol K Norcross, FRCS.....	Gas gangrene after burns H H G Eastcott, FRCS.....
Driving after anaesthetics W D A Smith, FFARCS.....	Vancouver style Paula J H Gosling, MB.....	Psychiatrists for the deaf? Sandra P Dowe.....
Sex determination and the H-Y antigen Ursula Mittwoch, DSC.....	Disinfection with glutaraldehyde G A J Ayliffe, FRCPATH, and others.....	Emotion and empiricism H A Wendel, MD.....
Polyunsaturated fatty acids in multiple sclerosis E J Field, FRCP.....	Safety of piped medical gases and electromedical equipment C S Ward, FFARCS.....	Microscopic words B J Freedman, FRCP.....
"Trench foot" caused by the cold I C Fraser, MFCM, and Judith A Loftus, MB.....	Non-motile sperms persisting after vasectomy: do they matter? I S Edwards, MB, and J L Farlow, FRCSed.....	Royal Medical Benevolent Fund D. Cotsell.....
Unnecessary skiing injuries K R Robak, MB, and A Benjamin, FRCS.....	Induction of labour and postpartum haemorrhage P W Howie, MRCP, and others.....	The World Medical Association H Hillman, MB.....
Alopecia areata Christine I Harrington, MD.....	Concurrent steroid and rifampicin therapy W van Marle, MB, and others.....	"Canvassing will disqualify" P L C Diggory, FRCP.....
		PPP plan for BMA members J G W Gelling, MBA.....
		The new consultant contract C E Astley, FRCP; A P J Ross, FRCS; R Hopkins, FDSRCS.....

We may return unduly long letters to the author for shortening so that we can offer readers as wide a selection as possible. We receive so many letters each week that we have to omit some of them. Letters must be signed personally by all their authors. We cannot acknowledge their receipt unless a stamped addressed envelope or an international reply coupon is enclosed.

Experience in a child health clinic

SIR,—Professor R S Illingworth's splendid article based on his experiences in an area health authority child health clinic (31 March, p 866) provokes me into writing down some of my own observations, based on three years' experience as a general practitioner running a well-baby clinic in our practice. His list of conditions and requests for advice from parents and his criticisms certainly made me think about the nature of the work we do. However, I am also struck by the dissimilarity of our experiences—perhaps, in part, based on the fact that I am a general practitioner with relatively little experience of hospital paediatrics, working exclusively with patients of my own or my colleagues.

My initial reaction after working in the clinic for a year was one of dissatisfaction—that I was being presented not with a wide array of paediatric clinical problems to which labels could be attached or firm advice or guidance given as to management but rather with a baffling array of questions, ill-defined problems affecting both mothers and infants which my medical training had ill suited me to answer, and a sense of bewilderment as to whether I was doing work of any worth at all. With time, and by using my knowledge of the families and the experience and help of my colleagues (including partners, health visitors, a visiting paediatrician, and a child psycho-

therapist attached to the practice), my attitude changed. I considered myself no longer as an expert in diagnosing leg clonus or a dab-hand at eliciting clicking hips (which I was not equipped to be in any case) but more as a sounding box for parents' concern about their babies and the difficulties they experience in just being parents. True, I became aware of the educational function of such clinics and I certainly had to revise my knowledge on immunisation and what constituted normal development. What also occurred to me was the extent of the maternal anxiety I met over relatively minor conditions affecting their offspring, the depressed mood of many young mothers, and the relief that was apparent on their faces when they saw mothers with similar problems exchanging views in the waiting area.

I would agree that expertise in diagnosing potentially troublesome clinical conditions and the offer of clear advice on management are important, but the value of such a clinic for me is also in recognising and bringing to light disturbance in mother-child interaction, and in offering time to a highly anxious and often depressed section of the community.

If I may refer to another point in the article, the decision about "ordering an investigation or merely seeing the child again" causes me less heartache now that I no longer work in

hospital—perhaps because the level of diagnostic uncertainty is higher in general practice and we are accustomed to facing such dilemmas time and time again in everyday work.

While I wholeheartedly agree with Professor Illingworth in his suggestion that paediatricians spend time in community child health clinics, I feel that it is even more imperative that general practitioners in training should be not only taught the full range of normal child development but also given an understanding of the imprecision and uncertainty common to such clinics.

PETER WILLIAMS

Kentish Town Health Centre,
London NW5 2AJ

SIR,—In these days when the trend seems to be toward general practitioner paediatricians I was surprised to read Professor R S Illingworth's article "Some experience in an area health authority child health clinic" (31 March, p 866). He has undoubtedly given an accurate précis of the types of problems experienced in this type of clinic, but what makes them so different from the ones experienced in general practice?

In my opinion a general practitioner with experience in paediatrics, rather than a "clinic doctor," is in an ideal position to allay parents' worries about physical growth, behaviour, and mental development, and guide them in their desire to help their child to achieve his best. The undoubted helpfulness of the health visitor to the clinic doctor applies equally to the general practitioner, but he has the added advantage of being able to experience the home situation and family circumstances on a per-

sonal basis. I do not accept that the "family doctor is too busy" for this type of consultation. Many of the conditions listed by Professor Illingworth can be handled adequately during routine consultation times; and in any case many interested general practitioners run similar "no-appointment" child health clinics as part of their usual working week.

I wholeheartedly agree with Professor Illingworth's "firm belief" that lecturers in paediatrics and paediatric registrars should have experience of paediatrics in the community, rather than total commitment to sterile academic paediatrics; but why not acquire the experience in a general practice setting?

BRYAN L ANDERSON

Perth

SIR,—I am delighted to learn that Professor R S Illingworth (24 March, p 797) has worked in child health clinics and welcome his suggestion that hospital-based paediatricians and medical students should have an opportunity to see the paediatric problems that present in the community.

It might be thought from Professor Illingworth's article that a health visitor is but a useful aid in a child health clinic, whereas in many child health centres it is a health visitor who takes the clinics, without a medical officer, and a health visitor who must deal with the many paediatric problems and queries mentioned in this article. Only 46.6% of under-5s ever attend such a clinic, according to figures in the Court Report; many of the paediatric problems of those who never attend are also dealt with by health visitors during home visits.

Professor Illingworth does not tell us on what evidence he bases his statement that health visitor training should be more realistic, beyond his reference to the examination papers. Perhaps his work in child health clinics did not allow him to observe the full range of the health visitor's work and skills? The health visitor training, part of which is undertaken with an experienced health visitor field work teacher, enables a nurse to work with families in the community carrying out health education, advising on child health, detecting deviations from normal, and helping parents to find and utilise the services their children need. Health visitors attend regular refresher courses, and continuing education through lectures and seminars on paediatrics helps them to remain up to date. I would agree that teaching by clinical medical officers is important too, as is the teaching carried out by doctors working in the primary health care team. After the basic health visitor training, as after every other basic training, continuing education and updating are essential.

LINDA DOWNING

Southampton

SIR,—Professor R S Illingworth suggests that health visitors have an important role in helping the clinic doctor, but that their training should be more realistic and appropriate facilities should be provided to keep them up to date in their work (31 March, p 866).

We suggest that health visitors are practitioners in their own right with a duty to liaise with all those concerned with health and welfare. Their training enables them to do this in a competent manner, and using the opportunities available both to attend lectures

and to study they are able to remain well informed about current ideas. A health visitor's training includes study of normal human development and deviations from normal, the social factors which influence behaviour, social policy, the social factors involved in health and disease, and the principles and practice of health visiting.

In relation to young children, health visitors are in a unique position in that they visit both mother and baby once the care of the midwife has ceased—usually between the 11th and 14th day after delivery. They usually make this visit having some knowledge about the mother's pregnancy, delivery, and subsequent health and also some information about the baby. The parents are seen subsequently with their children, either at the clinic or at home, by health visitors trained not only to look at individual problems but to observe the total situation. Health visitors learn to deal competently with many minor problems, but they also develop skill at knowing when to refer to others.

The role of the health visitor is to promote health in all its aspects through education. If Professor Illingworth is prepared to learn from health visitors I am sure that he will be able to score a few marks on one of their examination papers.

P M SAVILL
EVELYN POPE

Aldermoor Health Centre,
Southampton SO1 6ST

Upper gastrointestinal endoscopy

SIR,—I would like to comment on the paper "Upper gastrointestinal endoscopy: its effect on patient management" by Dr C D Holdsworth and others (24 March, p 775).

The authors state that 44 out of the 95 patients underwent a change in their management after endoscopy. This, I think, is a considerable number of patients and thus they correctly emphasise the usefulness of the procedure. However, what is more important is the fact that in their patient population there were at least 95 cases where the clinicians felt that endoscopy was needed in addition to the barium-meal examination. Therefore it seems that in these patients the conventional barium meal proved unnecessary and endoscopy should have been the primary investigation rather than radiology. It would be interesting to know how many of their 2500 endoscopic examinations yearly were preceded by an unnecessary conventional barium meal. I think the number must be high since the clinicians have changed to double-contrast barium meal in one of the centres since this study.

As experience with fiberoptic endoscopy increases all over the country, it seems that the conventional barium meal does not have a place in the investigation of the stomach and duodenum. Why should one bother with conventional radiology when in almost every radiological situation one needs endoscopy either for histological confirmation of the diagnosis (as when carcinoma is speculated), for assessment of progress of healing (as in duodenal ulcers and benign gastric ulcers), or in barium-meal-negative dyspepsia which is persistent?

May I point out that the statement that endoscopy has not been evaluated objectively in conditions other than acute upper gastrointestinal haemorrhage is wrong? There is at

least one study¹ that assesses fiberoptic endoscopy in the diagnosis of gastric carcinoma in relation to the conventional barium meal. The conclusion was that the barium meal has nothing to offer and thus endoscopy is the primary diagnostic tool.

Fiberoptic endoscopy therefore will be even more cost effective if it is not preceded by the expense of an unnecessary conventional barium-meal examination.

V MOSHAKIS

The London Hospital (Whitechapel),
London E1 1BB

¹ Moshakis, V, and Hooper, A A, *Clinical Oncology*, 1978, 4, 359.

What shall we teach undergraduates?

SIR,—I was interested to see the article by Professor V Wright and others (24 March, p 805) which set out the teaching choices of some 600 doctors. But I was sad to see no mention of occupational medicine. This is an expanding specialty, with probably the largest group of doctors practising predominantly outside the NHS, and is one of the specialties recognised as such by the Joint Committee on Higher Medical Training. Scarcely a year ago a new Faculty of Occupational Medicine was established within the Royal College of Physicians of London, which has already attracted over 500 members and associates to its ranks. In addition, the Society of Occupational Medicine, which dates its antecedents back to 1935, has now nearly 1500 members, of whom between 800 and 900 are thought to be full-time occupational physicians.

Training schemes in occupational medicine will be launched in 1979, but this article makes it all too clear that if the subject is not taught at undergraduate level it is simply not considered when the aspiring doctor chooses his career for a lifetime in medicine.

May I make yet another plea, by no means the first, that the health of people at work is of vital importance, provides a stimulating and varied medical career, and must receive its rightful place in the undergraduate curriculum?

ANDREW RAFFLE
Chairman, Specialist
Advisory Committee on
Occupational Medicine

Joint Committee on Higher
Medical Training,
London NW1 4LE

SIR,—In their most interesting paper "What shall we teach undergraduates?" (24 March, p 805) Professor V Wright and his colleagues appear to have neglected the views of the 600 doctors surveyed when they propose only the best taught specialties remain in the undergraduate curriculum. A majority of doctors did not recommend even one specialty for restriction to postgraduate study. Furthermore, some of the specialties the authors suggest are optional in the undergraduate curriculum were considered suitable for postgraduate study alone by under 1% of doctors.

A survey of 55 first- and second-year clinical students at King's College Hospital (including only those who had not read the above paper) was carried out to determine what priority students gave the specialties listed by Wright *et al.* Two points emerged. Firstly, 95% of students considered that 23 of the 28 specialties should be separately taught in the undergraduate curriculum. Secondly, the results of