Rate of remuneration according to number of sessions worked.

tion for each session worked will be only 92.3% of that for the eight scheduled sessions of the standard contract.

The CCHMS Negotiating Committee made the following comments on the new contract¹: "Those individuals who are presently whole time and work all the normal working hours from Monday to Friday are likely to need to maintain this pattern and will, as a result, be paid for 12 NHDs each week. The part-time consultant with significant private practice will be able to accept the standard contract and discharge it fully while retaining sufficient free time to carry on his private practice successfully." Thus consultants working a 5-day week will be remunerated at an overall rate which is 96% of that of the basic 4-day contract and those working a 5½-day week will receive less than 95% of the standard rate.

Only if the Review Body recognises this anomaly and prices timetabled NHDs in excess of eight at a rate 25% higher than those forming the "standard" contract will all consultants receive equal pay for equal periods of scheduled work.

R B PAYNE M K MASON
S I JACOBS G J HARDY
D L BARNARD L M SWINBURNE
J A DOSSETT

St James's University Hospital,
Leeds

¹ *British Medical Journal*, 1978, 1, 1300.

*. *The Secretary writes: "The authors have made the mistake of regarding the two unscheduled NHDs as involving no work. These NHDs are in recognition of the very heavy administrative load undertaken by consultants at unpredictable hours and of their obligation in accepting continuous responsibility for patients or departments under their care. The implication that there is no work involved in the unscheduled NHDs and consequently any NHDs above the basic 10 is quite fallacious." —Ed, *BMJ*.

Reduction of pay-beds in the NHS

SIR,—Your issue (13 January, p 146) carries information that the Health Services Board proposes to reduce the 35 private beds in the Bury, Oldham, Rochdale, and North Manchester Health Districts to nil on the grounds that there are alternative facilities at Highfield Nursing Home. The Health Services Board, in its letter to our health authority, suggests that it requires 75% occupancy at Highfield Nursing Home as being a satisfactory index of full occupancy and suggests that there is still spare occupancy at Highfield. We are reliably informed by the managers of Highfield Nursing

Home that the occupancy rate for the four full weeks in October was 80% and for the 21 weeks ending November was 67%.

We are most concerned that the Health Services Board is proposing to take action—and the board itself by its own admission never revokes its action—on the basis of inaccurate statements concerning the capacity of Highfield Nursing Home to replace the 35 remaining private beds in these four health districts as, in fact, there are only 36 beds altogether at Highfield Nursing Home.

Furthermore, while not deprecating the high standard of nursing care and the modern theatre facilities at Highfield Nursing Home, it should be noted that there are no x-ray department and no resident medical staff at Highfield. On clinical grounds, therefore, the board's statement that the beds at Highfield Nursing Home constitute alternative facilities to the private beds in the hospitals in the four health districts is incorrect as it would seem to us to be medically unsafe to manage patients with certain clinical disorders in Highfield Nursing Home with its present facilities.

O BAUEROVA O P LAKHANPAUL
C D BAUMBER S LAL
M E BENAIM M M P LEAHY
G S BHATT J C LOWRY
J C BRADLEY J MCLENACHAN
D J A BROWN D K MUKERJI
D M DAVIES F O'CONNOR
S G DEODHAR K W PEARSON
J P FRASER R PHILLIPS
M J GOODMAN A J SIMS
A E GREEN J F STONE
K S HAWORTH J B L TAYLOR
K KAUSHAL H W WILSON

Bury General Hospital,
Bury, Lancs

BUPA and private practice anaesthesia

SIR,—At the beginning of last year, when I heard that British United Provident Association (BUPA) was soliciting comments in preparation for a new edition of *A Guide to Private Consultant Practice*, I felt constrained to write to them about the lack of attention paid to the anaesthetist in private practice in the first edition. It seems to me that private practice is something of an abyss to the newly appointed consultant anaesthetist; he is unlikely to have been involved in assisting as a junior (in contrast to his surgical colleagues) and what he does learn about private practice is probably distorted and unfavourable. I wrote to BUPA suggesting that there should be a section devoted to private practice anaesthesia in the new booklet. I also suggested that 20% of surgeons' fees was a niggardly amount for the anaesthetist and that they should quote anaesthetists' fees separately in their patients' booklets.

I had no reply to this letter and six months later I wrote again enclosing a copy of my original letter. This time I was favoured with a reply which was slightly encouraging. Judge my disappointment in October when I was sent a copy of the new booklet and found that there was no section or even sentence about the problems of private practice anaesthesia, which are of interest to perhaps 1000 members of the medical profession, and that fees were still quoted inclusively with 20-25% as the usual ratio to the surgeon's fees. BUPA now has certain schemes in which the anaesthetist's fees are shown

separately, and makes the curious comment in the booklet that "it is accepted that anaesthesia should thus be seen to be recognised as a specialty in its own right." This is a very conservative statement 120 years after the death of John Snow, or even 30 years after the establishment of the NHS and appointment of consultant anaesthetists, but perhaps crystallises BUPA's attitude to the specialty.

ADRIAN PADFIELD

Department of Anaesthetics,
Hallamshire Hospital,
Sheffield

Trade unionism in medicine

SIR,—My thanks to Dr Richard Dreaper (24 February, p 558) for his support of my plea at the recent Honorary Secretaries' Conference for a more cautious approach to trade unionism by the BMA.

At a time when there are severe political pressures on the BMA to adopt trade union status and principles and yet tremendous apathy about the whole concept, it would appear to me (certainly in our division), I feel very uneasy about what the future holds for us all as a professional body. Do our leaders really feel convinced that this move will improve our status in society, our financial position, or, even more important, the service we are able to offer our patients? If the BMA is to have power as a trade union, does it not seem logical that it eventually adopts the closed-shop principle and, ultimately, the right to "withdraw its labour"?

If this is what is envisaged I suggest we place the facts fairly and squarely before our membership and invite them to express their views by postal ballot.

EGRYN M JONES

Wilmslow, Cheshire

SIR,—The nursing profession has shown us a splendid example in refusing, by a big majority, to contemplate strike action, as they would thereby "be betraying a trust." May we also stand firmly by our principles and refuse ever to consider industrial action or, as Dr Jean O Williams (17 February, p 494) warns, "be betrayed into such industrial action again." I heartily agree with Dr Richard Dreaper (24 February, p 558) that if we seek short-term advantage by such means we shall eventually cease to be regarded as a profession. Let us, as Dr D L McNeill (24 February, p 559) emphasises, keep our professional status and independence and give the TUC a wide berth.

MARGARET WILKINSON

Cardiff

College of Anaesthetists?

SIR,—The letter from Dr R S Atkinson and his colleagues (3 March, p 624) lists reasons why they believe anaesthesia should remain as a faculty within the Royal College of Surgeons, and questions how much support there is nationally for a college of anaesthetists.

While support nationally cannot be assessed without investigation, it is worth recording that at the annual general meeting of the Junior Anaesthetists Group held at Exeter in March 1978 there was overwhelming support for the establishment of a college (nobody voted against the formation and two voted as

uncertain). At least this helps to confirm that anaesthetic juniors want a separate college. In addition, many young anaesthetists feel that the establishment of a college can only serve to strengthen anaesthesia and establish it as an independent specialty. Perhaps then something may be achieved to correct the provident societies' view that treatment should be undertaken by consultants except in anaesthesia, where anyone will do. What effect has the "prestige, tradition, and esteem" of the faculty had on this? In the same issue of the *BMJ* (p 638) there is an assessment of the distribution of distinction awards to the different specialties. Does this represent the esteem in which anaesthetists are held? Or perhaps they do not deserve distinction awards for the quality of their work? A new college will not cure all the problems, but it may give the next generation of anaesthetists a fighting chance.

Many of the arguments have been succinctly analysed, though with a different emphasis, by Spence and Norman¹ and Zorab²; and I feel that the Anaesthetists' Academic Foundation may in fact achieve far more support once it becomes certain that the college will be established. Finally, the signatories of the letter are less than fair to the Association of Anaesthetists' offices, for all the benefits of the BMA are available to supply common rooms and food; and they also failed to describe the cramped offices of the Faculty of Anaesthetists, and the fact that it is only when you reach these offices that you realise that you are no longer in the Royal College of Surgeons but in the adjacent building.

M A THOMPSON

Department of Anaesthetics,
Guy's Hospital,
London SE1

¹ Spence, A A, and Norman, J, *British Journal of Anaesthesia*, 1978, **50**, 313.

² Zorab, J, *Anaesthesia*, 1979, **34**, 72.

Distribution of general medical practitioners

SIR,—Mr M J Buxton and Professor R E Klein (17 February, p 465) rightly point out that the nine Family Practitioner Committee areas (FPCs) with the most inflated lists are in the London area, with Kensington, Chelsea, and Westminster (KCW) at the top. They note that these FPCs (with one exception) also have low nominal list sizes. This is particularly true of inner London, with KCW again being in first place.

They then go on, however, to make the logical suggestion that nominal list size figures understate the inequalities in the distribution of general practitioners—with the implication that inner London is overdoctored in comparison with the rest of England and Wales. Although from the published figures this would appear to be the case, it is not the experience of those working in the area—for example, the annual reports of the three Community Health Councils in KCW have contained evidence of difficulties experienced by patients, particularly the elderly, in finding a general practitioner to accept them on his NHS list.

Average list sizes mask the considerable variations in list size of individual doctors, and the difficulty apparently experienced by patients is partly explained by a survey carried out by the administrator of the KCW FPC: it showed that 119 of the 260 doctors (46%) for whom the FPC was responsible were not

accepting new patients. The reasons for this apparently contradictory situation are various. KCW has the largest proportion of elderly GPs (13% aged 70 years and over compared with 3% nationally in 1977) and of single-handed GPs (56% compared with 16% nationally) in the country.¹ The patients they deal with have predominantly psychological and social problems.¹⁻³ There are high proportions of elderly living alone but, in an area with the highest population density in the country, there are difficulties with traffic and parking problems when visiting patients. It is particularly difficult to estimate the true population of the inner London areas owing to its rapid mobility, but this very mobility means that it is difficult for a GP to get to know his patients, for whom he may have no previous notes.¹

General practitioners in London are reported as having the lowest average net earnings under schedule D in the country.¹ Yet London is one of the most expensive places to live, and some GPs may need to maintain small NHS lists in order to carry out other professional commitments.

Even though the number of patients needing to be "allocated" to GPs is low (it being FPC policy to deal with the matter informally when possible), the number of telephoned and written inquiries dealing with assignment of patients to doctors' lists is high.¹ This evidence, together with that from the CHCs, tends to indicate that for patients looking for an NHS doctor in the KCW area the published statistics used by Buxton and Klein are misleading.

B JARMAN

MARGARET M LALLY

London NW8

¹ Jarman, B, *Journal of the Royal College of General Practitioners*, 1978, **28**, 598.

² Imber, V, *Statistical and Research Report Series*, No 16. London, HMSO.

³ Farmer, R D T, Preston, T D, and O'Brien, S E M, *British Journal of Preventive and Social Medicine*, 1977, **31**, 171.

⁴ *General Practitioner*, 21 April, 1978, p 1.

Nursing at a crossroads

SIR,—I enter the "Nursing at a crossroads" debate with some trepidation but, as a junior doctor who works daily with nurses below the rank of sister (and who is married to a qualified nurse), I feel that the views of those "on the ground" have not really been covered by the higher echelons of medical and nursing staff who have so far contributed.

Only one of your correspondents has mentioned the effect of salary on recruitment, yet everybody one asks feels that this is surely the crucial factor. The harsh fact is that people leaving school, however dedicated to nursing they may be, compare salary scales in different jobs; and nurses, like so many other workers in the National Health Service, have clearly fallen far behind.

The reasons for the near 30% drop-out of nurses before the completion of training are not so clear. Certainly in my experience (as other correspondents have suggested) the outdated, authoritarian attitude of many of those involved in training student nurses can be held partly to blame. Secondly, although much is made of the responsibility, many believe that general nurses are now allowed less responsibility than in the past—and indeed that decision-making is almost discouraged in the modern nurse (except, of course, on night

duty, when the coin is turned and the responsibility can be terrifying).

Following on from this is the question of educational requirements. Most of those I have spoken to feel that the acquisition of 5-6 O levels is necessary, but there is considerable scepticism over the requirement of A levels, which now seems to be the norm in some areas. Many believe that nursing is not an "academic" discipline but one needing common sense and practical ability (not that these are necessarily mutually exclusive); they feel that nursing managers are trying to change it in to something which it is not. At a time when initiative is apparently being restricted on the wards surely higher examinations are inappropriate, and nurses who have passed them are among those who are most disillusioned with practical nursing.

To sum up, the nurse's lot must be made a happier one. The training should be made more human; there should be less interference from nursing administrators; and, most importantly, the pay situation must be rectified. Unfortunately, Mr Ennals's continued lack of support for this cause, despite pleas from the Royal College of Nursing, only serves to publicise further the present deplorable situation and can do nothing but harm to nursing recruitment and thus to the National Health Service as a whole.

A J VALLANCE-OWEN

Accident and Emergency Department,
Royal Victoria Infirmary,
Newcastle upon Tyne

State of community medicine

SIR,—I have read with much interest the report (17 February, p 503) of the meeting of the Working Party on Community Health Doctors and the suggestions for training and career structure. I am delighted to learn of the positive progress being made to improve this aspect of the medical services provided by the National Health Services.

I read with great care the proposed training programme. While it appears that a comprehensive programme of general paediatric training is envisaged, I note that there is no reference to any postgraduate training in gynaecology. There will I think be an increasing demand for "well-women clinics" and there will be a need for family planning clinics in many areas, partly to meet patients' preferences and partly as necessary training centres for staff and foci for research projects. It is imperative for doctors working in these clinics to have postgraduate training and continuing education in non-surgical aspects of gynaecology, especially as the time spent on gynaecology by undergraduates in some medical schools has been so drastically reduced.

JEAN LAWRIE

Istana Darul Hana Office,
Brunei

Next year's pay

SIR,—You published a letter (3 March, p 626) from Dr Michael Wynn about the Review Body's recommendations for 1979-80.

The evidence submitted to the Review Body is confidential until the report is published, and I am therefore unable to comment on the detailed submissions made on this occasion. I would, however, like to point out that the Review Body in its eighth report said that at