diagnosis, or both. The biochemist then decides what tests, *if any*, will help. Moreover, no report would leave the department without an expert interpretation of the result. To implement such a proposal would require a drastic change in the philosophy of both clinicians and biochemists; but it is, of course, the only logical solution.

C SANDERSON

Biochemistry Department,
Airedale General Hospital,
Keighley, W Yorks BD20 6TD

Air embolism and intravenous catheters

SIR,—We read with interest the letter from Dr D P D Colqhoun (2 June, p 1489) reporting a further three cases of air embolism occurring via central venous catheters when the hubs became disconnected from the administration sets. We suspect that many similar incidents have occurred and have not been recognised.

When total parenteral nutrition is being administered in our general surgical ward, a Baxter-Trevenol "W" or "Y" administration set with the new Luer lock mechanism is used in conjunction with the central venous catheter. An incident has just been reported to the DHSS in which a patient receiving parenteral nutrition via a Vygon Surcath long-line and a "W" administration set almost suffered an air embolism. The thread of the Luer lock failed to engage the hub of the catheter and the two components slipped apart. Fortunately, the fault was recognised and corrected. We feel that this connection should be carefully inspected after assembly and at regular intervals during the period of infusion.

E METCALFE
D GRIFFITHS
J L PETERS
N M GREENWOOD

University College Hospital,
London WC1E 6AU

Captain Cook and scurvy

SIR,—May I suggest that Dr C C Booth is in error (2 June, p 1470) in suggesting that Captain Cook's success in making his historic scurvy-free voyages was due to Lind's work?

Cook's work was independent, and due to his using other antiscorbutics, such as malt and cabbage, and "scurvy grass" and wild celery when obtainable, together with his insisting on other measures such as strict cleanliness and adequate fresh water. It has even been suggested that his work delayed the introduction of lemon juice into the Navy. Another of the great accomplishments of this non-medical man was his distinguishing between yaws and syphilis.

BRYAN WILLIAMS

Chichester, Sussex

Vaccination and pregnancy

SIR,—We feel that the answer to the question on the vaccination of pregnant or nursing mothers in your "Any Questions?" column (5 May, p 1194) could be misinterpreted. The statement that "vaccines given to the mother in pregnancy will, of course, benefit the infant by the transplacental route whether or not he

is breast fed" fails to emphasise the dangers of giving live vaccines during pregnancy and the possibility of infection to the unborn infant.

In general, live vaccines are contraindicated in pregnancy unless the risk of getting the disease outweighs the risk of vaccination. Fetal vaccinia, although rare, is well documented; if the mother must be vaccinated this should be done under antivaccinial globulin cover. The risks of using live polio and yellow fever vaccine appear to be very small.¹

A pregnant woman intending to travel to a developing country should have the killed polio vaccine unless she has to travel immediately, when the live oral strain should be used. Similarly, if travelling to an endemic yellow fever area vaccination is advised, although it should be delayed if possible until beyond the first trimester. Rubella vaccination remains a contraindication in pregnancy; the seronegative mother should be vaccinated on the second to fourth day after delivery. At this stage in the puerperium the risk of pregnancy in the ensuing six weeks is extremely low. Nevertheless, contraception should be advised for at least three months.

G DU MONT
R C BEACH

Department of Paediatrics,
St Thomas's Hospital,
London SE1 7EH

¹ Levine, M M, Edsall, G, and Bruce-Chwatt, L J, *Lancet*, 1974, 2, 34.

An alternative to colostomy

SIR,—The introduction of circular stapling devices for intestinal anastomosis (the Russian SPTU gun and the American EEA gun) have enabled colorectal surgeons to perform safely very low rectal anastomoses in patients with carcinoma of the middle third of the rectum and at the same time preserve good anal sphincter control.¹ This obviates the need for abdominoperineal resection with a permanent colostomy in many of these patients. In our experience using the EEA gun the early results of this method of treating rectal cancer are excellent and the avoidance of a permanent colostomy saves the patient from many of the problems that we have previously commented on.² However, we are concerned about the economics of using the EEA gun and in particular the reluctance of some health authorities to sanction the purchase of these instruments because they are "too expensive" and "not a good buy."

The basic purchase price of the EEA instrument is £595 plus VAT and the cost of a set of staples is £47 per patient. The operating time to perform a sphincter-saving resection using this technique is about one and a half hours, the radiologically demonstrated leak rate is negligible, and the patients rapidly recover full continence after operation. The duration of postoperative stay in hospital is considerably less than that after an abdominoperineal resection.

By contrast, the patient having a conventional synchronous combined abdominoperineal resection has a larger and more blood-losing operation, a longer postoperative stay in hospital, and a permanent colostomy. We conservatively estimate the price of colostomy care over five years as between £597.50 and £2135. The low and high prices include home nursing costs and the cheapest or the more expensive conventional stoma

appliances, but exclude general practitioner, local authority disposal, and extra laundry costs. We have not attempted to quantify the social cost of the colostomy to the elderly patient, hospital and convalescent stay after abdominoperineal resection, or the many hidden costs of long-term care of stoma patients.

We therefore conclude that despite its high purchase price the EEA gun is a good economic alternative to a permanent colostomy.

N R PARROTT
H BRENDAN DEVLIN

North Tees General Hospital,
Stockton-on-Tees, Teeside TS19 8PE

¹ Goligher, J C, *Annals of the Royal College of Surgeons of England*, 1979, 61, 169.

² Devlin, H B, Plant, J A, and Griffin, M, *British Medical Journal*, 1971, 3, 413.

Cyclophosphamide in chronic active hepatitis

SIR,—Dr I T Gilmore and others, in reporting the results of a clinical trial of cyclophosphamide for the treatment of active chronic hepatitis (28 April, p 1120), conclude that because of persistent azoospermia "the benefits [of cyclophosphamide] would have to be striking to justify its use." While we agree with their conclusion, we believe there are more important reasons why this is so.

We are surprised that no mention was made of the association between alkylating agents (including cyclophosphamide) and cancer in humans. There are at least 64 case reports of cancer (mostly leukaemias, lymphomas, and bladder cancers) in patients treated with cyclophosphamide for a variety of malignant and non-malignant diseases.^{1,2} In addition, cyclophosphamide causes cancer in mice and rats when administered in doses similar to those used in clinical practice.¹

Cyclophosphamide is often considered together with other alkylating drugs in reports of adverse effects, since individuals are frequently treated with more than one of them. Six of these drugs, which have been evaluated in the monographs of the International Agency for Research on Cancer,³ have data on carcinogenicity from case reports, epidemiological studies, or both. Recently a group of experts in cancer research concluded that, of these six, two are causally associated with cancer in humans (chlorambucil and melphalan), and three (chlorambucil, cyclophosphamide, and thiopeta) are probably carcinogenic to humans but sufficient evidence to establish a causal relationship is lacking. The data for the remaining drug (triaziquone) are derived from four case reports and are inadequate to allow any conclusions regarding carcinogenicity.⁴

Unfortunately, most of the evidence on the carcinogenicity of these compounds comes from case reports, which do not allow incidence rates to be calculated. However, the few available epidemiological studies⁵⁻⁸ suggest that over a period of 10 years from 5% to 15% of patients given alkylating agents will develop cancer. This is substantially in excess of what would be expected in the absence of these drugs.

Investigators planning clinical trials using these drugs, particularly in adjuvant chemotherapy for early stages of cancer or in the immunosuppressive treatment of non-malignant disease, should balance this sizable