carcinogenic risk (along with other adverse outcomes) against the potential benefits.

RALPH ALTHOUSE
J E HUFF
LORENZO TOMATIS
J D WILBOURN

Unit of Chemical Carcinogenesis,
International Agency for Research
on Cancer,
Lyon, France

¹ International Agency for Research on Cancer, *IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Man*, vol 9, p 135. IARC, Lyon, 1975.

² Puri, H C, and Campbell, R A, *Lancet*, 1977, **2**, 1306.

³ Tomatis, L, et al, *Cancer Research*, 1978, **38**, 877.

⁴ IARC Working Group, *Chemicals and Industrial Processes Associated with Cancer in Humans*, submitted for publication.

⁵ Thiede, T, and Christiansen, B L, *Acta Medica Scandinavica*, 1969, **85**, 133.

⁶ Reimer, R, et al, *New England Journal of Medicine*, 1977, **297**, 177.

⁷ Einhorn, N, *Cancer*, 1978, **41**, 444.

⁸ Law, I P, and Blom, J, *Oncology*, 1977, **34**, 20.

Management and severity of tetanus

SIR,—The question raised by the article of Dr R S Edmondson and Mr M W Flowers (26 May, p 1401) is whether intensive care in the management of tetanus is causing unnecessary deaths. The mortality (10%) in their series of 100 must be compared with a mortality of 7% in the much larger series reported by Sanders¹ in a developing country where artificial ventilation and intensive care were not available.

My experience over two years in North India bears out these survival figures. Following instillation of antitetanus serum via lumbar puncture, chlorpromazine and diazepam were titrated against muscular spasms, respiratory complications were minimised by nursing in the head-down position, and locally trained nurses were specifically taught to watch for respiratory or cardiac difficulties. Feeding was usually possible by nasogastric tube at one week, once the worst of the spasms had passed.

Prognosis in tetanus should be related not only to severity of symptoms at presentation but to the site of entry of the tetanus bacilli. Women who had become infected during childbirth or during a normal menstrual period would be expected to have more severe symptoms, a longer convalescence, and also a higher mortality. This may be accounted for by the large area of the uterine wall available for infection.

Neonatal tetanus is virtually unknown in Great Britain. Although having many features of the adult disease, conventional treatment has remained unsuccessful and few centres can improve on a mortality of 80%.

JOHN RENNIE

Department of Surgery,
Charing Cross Hospital Medical
School,
London W6 8RF

¹ Sanders, R K, et al, *Lancet*, 1977, **1**, 974.

Medical reports and confidentiality

SIR,—The letter of Drs G B May and W P Garson (26 May, p 1427) raises issues which are certainly important, and which currently are much in the minds of doctors concerned about ethics. I would like to add one or two points for consideration—not necessarily because I believe Drs May and Garson to be wrong in their argument: it is just that, as

in most complex situations, every factor does not point towards the same conclusion.

I must first declare an interest, as a salaried adviser to an insurance company. My employer is concerned with health insurance, and Drs May and Garson write about the Life Offices; but the principles involved are, I think, the same. What I say now is my view, and not that of BUPA or of any health insurer.

It will help, at the outset, to distinguish between medical reports and the medical records on which they may be based. The latter, we would all agree, should—subject to the patient's consent—go only from doctor to doctor, although the law may compel wider scrutiny. But I do not think that the "medical attendant's report" is in the same category of confidentiality, even though both ethics and common sense suggest that it too go from doctor to doctor for interpretation and confidential storage.

As secretary, years ago, to the tripartite (Bar Council/Law Society/BMA) Committee on Medical Evidence I learned that the "privilege" attaching to medical documents and reports is vested wholly in the patient (or responsible relative), and not in the doctor or doctors who compile them. If the patient agrees to disclosure or reporting, the doctor—though he may feel obliged sometimes to advise against this—has no right of veto to refuse to implement his patient's decision. Drs May and Garson in that context raise two relevant issues: the patient may not fully appreciate what he is doing, and in any event is under pressure to agree. On the first of these, the medical attendant's duty is clear: he explains as best he can what are the implications, but the patient decides. The second point can be the subject of pejorative treatment, and by reference to "blackmail" Drs May and Garson—by their use of quotation marks—acknowledge this fact.

A contract of insurance is, in law, a matter of the "utmost good faith," and any proved departure from this—by the proposer—will usually invalidate the contract. If one seeks to insure one's health, or one's life, it is self-evident that one must make a true declaration of one's state of health. Sometimes this alone will suffice, but not always: medical verification (or explanation of something unclear in the declaration) may be properly required by the insurer. This is made necessary because, in insuring such things as goods and chattels or a holiday, the insurer has himself the knowledge to make full relevant disclosure, but he often has not this knowledge of his own medical history. It is notorious that we, as doctors, are not always fully communicative—either because we fail to get our meaning across or because we believe that the patient's best interest is served by a less than full disclosure to him of the nature of either investigations or findings.

Drs May and Garson suggest that the independent medical examination and report would serve as well as a report. Sometimes it does—but not always (quite apart from the added inconvenience and expense to the patient, who in the end pays, directly or indirectly). In a similar context, the interview of an applicant for a position of trust by an experienced personnel officer may suffice, but can never wholly do away with recourse to confidential reports from character referees. If one wishes to embark on a major financial transaction—or even to obtain a line of credit one usually is asked for, and agrees to, references from banker and employer. Is it

different in principle if the financial transaction is one in which the assessment of health or "life" is a major factor?

True, life assurance and health insurance can be, and sometimes are, provided without initial inquiries, but their total elimination would inevitably put the financial burden of "carrying" the unfit (and, alas, the dishonest) on the fit and honest. If premiums (or benefits) are not tailored to the individual and to the risk, then everyone must pay a higher premium, to meet the greater risk of the group and its members. This is, of course, the principle of the welfare state, and a perfectly respectable one. But it is by definition an arrangement which includes everyone, the good risk and the bad risk alike, and so is not comparable with voluntary insurance.

So far as the latter is concerned therefore, recourse to the "medical attendant's report" must in some cases continue to be a condition of securing personally selected cover, if the premiums charged are also to be personal and not on a communal basis.

DAVID GULLICK

BUPA
London WC2R 3AX

Isolated communities and their doctors

SIR,—I do not think Dr M A Wilson (2 June, p 1492) has really understood our letter about the rural practice fund (19 May, p 1358).

Firstly, the RPF may have started way back as a compensation for the necessarily small lists of the single-handed rural GP; but it is surely now regarded as a fund to offset the escalating costs of motoring, and as such should be based on these increased costs.

Secondly, he says that the 1959 inquiry was independent and fair. I had thought that the publication of Dick Crossman's diaries, covering the period when he was head of the Ministry of Health, had finally disposed of this fiction. As a matter of fact, a few years after the 1959 report a senior official in the Ministry admitted to me that the policy was to make rural practice an annexe of the nearest town—up to 20 miles radius was his yardstick. As our complaint is that the rural practice fund favours the town doctor extending out into the country, one surely should not be considered too paranoid to see a connection between them. In any case the capital costs for a young doctor taking over a rural village practice are now so prohibitive that the Department can really sit back and accomplish its purpose without further action.

The appearance of a few senior rural GPs arguing and fighting for what to DHSS officials must be a lost cause must seem strange to them; but perhaps they might understand that some of the profession dislike seeing valuable parts of British medical tradition destroyed by casual, secret, bureaucratic action, and are not prepared to let it pass without protest.

C G ELLIOTT

Lewes, Sussex

NHS certificates for hospital patients

SIR,—As my frequent protestations to my local hospital, LMC, medical staff committees, etc, have produced no result may I appeal through the medium of your journal for a change of