Emergency kit

SIR,—The general practitioner is entrusted with primary medical care, which if properly carried out is a formidable undertaking. For this reason he needs support from several quarters—patients, hospital, administration. Take emergency calls.

Some 250 000 attacks of myocardial infarction alone can be expected each year. Doctors have learned that the first two hours are crucial in such an attack, yet more often than not the vital hours are disregarded by patients themselves. What can the doctor do if he is not summoned in time, and why do prospective patients lack the necessary knowledge? Heart attacks are not the only emergencies. Doctors can be confronted with an acute respiratory failure, status asthmaticus, tension pneumothorax, anaphylactic shock, diabetic coma, dangerous convulsions, or other life-threatening conditions. Their efficiency and speed of action are essential in such circumstances; the suffering from acute retention of urine or renal colic can be immense. The rescuer has to be prepared and doctors involved in primary care have their emergency kits at the ready. But it is a rather haphazard affair. Different medical men have different ideas, some treatments are controversial, the general practitioner becomes confused. A doctor called out in the middle of the night may pick up a drug given to him the previous day by a pharmaceutical representative with a reassuring account. Which injections should be specially selected? Practolol? Atropine? Heroin? Corticosteroids? Which are the most essential appliances—an airway or a suction pump, a catheter or a stomach tube? An emergency kit must be handy, confined to essentials.

Despite the controversy about the standardised equipment for emergencies, or because of it, a panel of medical and pharmaceutical luminaries should come to the GP's aid and contrive one. It could be revised every four years or so. It worked with the *British National Formulary*; there is no reason why it should not work here. Family practitioner committees could help in supplying standardised emergency kits to individual practices at a non-profit-making price; I doubt if any GP would object to this.

C ROBERTS

Harrow, Middx HA2 6LY

Poisoning and enuresis

SIR,—We read with interest your leading article (17 March, p 705) on enuresis and the report by Dr A J Cronin and others on poisoning with tricyclic antidepressants (p 722). In the leading article you state that treatment should be reserved for older children and I would agree with this concept. When such a policy is adopted it is likely that by puberty less than 1% of children will require treatment.¹ Parents are unlikely to accept such delays with equanimity, and it should be remembered that enuresis may be an indication of possible urinary tract pathology. Where simple measures fail in obtaining a cure then a more detailed examination should be undertaken. One of the most useful measures is the buzzer alarm² but this instrument can result in the "non-enuretic" members of the family being awakened while the unfortunate sufferer remains asleep.

I would disagree with Dr A J Cronin and his colleagues when they state that "tricyclic

antidepressants are the only drugs which are of any benefit." We have used a regimen of diuretic therapy with success in the management of enuresis when all other methods short of surgical intervention have failed.³

R SCOTT

Urological Department,
Royal Infirmary,
Glasgow G4 0SF

L MORRISON

Stobhill Hospital,
Glasgow G21 3UW

¹ Gairdner, D, 1965, *British Medical Journal*, 1965, 2, 91.

² Meadow, R, *British Medical Journal*, 1977, 2, 1073.

³ Scott, R, and Morrison, L, *Journal of the Royal College of Surgeons of Edinburgh*, to be published.

SIR,—The solution to the problem of how to prevent the tragic and unnecessary deaths of enuretic children under the age of 5, discussed in your leading article (17 March, p 705), is basically simple but depends on the determination of those in whose power it lies to carry it out.

The solution is to treat the under-5s with the waking device, which, as you rightly say, renders 80% so treated permanently dry and yet is completely safe. To delay this treatment until 7 or later, as is often advised, will not save the younger children from the risks of poisoning; and in any case this delay is quite unnecessary, for young children respond to this treatment as well as the older ones.

Twenty years ago I started a clinic based on the waking device. It was open to enuretics of all ages and the results of the first 100 so treated were published (22 November 1958, p 1268). Among the under-5s 63% became completely dry within five weeks and 74% within 10 weeks. There were also two who were just under 3, both of whom became dry after only two rousings. These would have raised the cure rate to 80% but were not included since, by convention, they did not officially qualify as enuretics.

Thus the waking device can and therefore should be used at the earliest opportunity. It only remains to decide who should administer it. Not the children's hospitals—for they would be swamped with work, and besides they are too far away from many needing treatment. Nor should it be the GPs since many have already shown a reluctance to use it. The answer surely is that there should be clinics established in every district equipped with adequate apparatus and supervised by health visitors or nurses, and it should be widely known throughout the district that these clinics are prepared to treat enuretics of any age from 3 years old upwards. This would virtually abolish these iatrogenic poisonings, which have increased so greatly of recent years, and at the same time bring relief to countless families to an extent not previously achieved.

Paradoxically, the tricyclic compounds, which in excess can prove so lethal, should be employed at these clinics—but their use would be restricted to those who failed to be aroused by the waking device, and the dose given would tend to be lower than that often given to suppress bed-wetting (for example, for a child of 5 10 mg of imipramine rather than 25 mg). In some subtle way these preparations often seem to normalise the sleep pattern so that those who were deep sleepers and failed to respond to the alarm now do so and become permanently dry like any other bed-wetter. Thus the cure rate should rise even higher but

with minimum risk if the use of tricyclic compounds were confined to clinics fully aware of the potential dangers.

T H GILLISON

Ilkeston, Derbyshire DE7 5HL

SIR,—Yet again your pages (17 March, p 722) bear witness to the tragedy of poisoning with tricyclic antidepressants in childhood. The quantity ingested is likely to be larger when the tablets are prescribed not for the child but for an adult relative. The drug prescribed for the adult should be dispensed in childproof containers if young children have access to it.

In one of our cases recently, a girl of nearly 4 years ate all 35 of the 50 mg imipramine tablets prescribed for her mother's enuresis. Two hours later she arrived in hospital deeply comatose, hypotensive (systolic pressure 50 mm Hg), and hypothermic. Her electrocardiogram showed right bundle-branch block and two atrial premature beats in trigeminy with each sinus beat. Soon she progressed to grade I atrioventricular block (PR interval 0.36 s), with a sinus rate of 60 beats/min interrupted irregularly by atrial premature beats. To pre-empt complete heart block and to treat arrhythmias, an endocardial electrode was introduced into the right ventricle through a cannula inserted in the right subclavian vein. At no site could a threshold of less than 2 V be found and this was accepted. The pulse generator was switched to 3 V and to the demand mode. Although left bundle-branch block was recorded at one stage, complete heart block never occurred. The patient went into a supraventricular tachycardia at a rate of 170 beats/min, however, with an unrecordable blood pressure; but control was obtained by overdrive pacing and an elective subsequent pacing rate of 120 beats/min. She made an excellent recovery despite the large dose of drug taken and about five hours of poor cardiac output.

With appropriate measures, a favourable outcome has been reported¹ even in a child with a critically low cardiac output requiring continuous cardiac massage for two-and-a-half hours. We avoided using antiarrhythmic drugs because we feared that their negative inotropic effect on a depressed myocardium might be catastrophic. Fortunately, overdrive pacing was repeatedly successful. We would emphasise, however, the need to accept a moderately high pacing threshold in a poisoned heart. Frequent checks (every two to four hours, for example) of the threshold allow the pacing voltage to be reduced as recovery occurs.

G MCENERY
R S WINWOOD

Whipps Cross Hospital,
London E11 1NR

¹ South, D P, and Kilpatrick, S M, *British Medical Journal*, 1974, 4, 508.

Tricyclic antidepressant poisoning

SIR,—Dr A J Cronin and others (17 March, p 722) have rightly drawn attention to the dangers that can arise in children from the accidental ingestion of tricyclic antidepressant drugs. However, I feel that their case report and conclusions deserve further comment.

They describe the difficulties of treating massive overdosage with these drugs and indeed their patient received at least three different agents intended to reverse cardiotoxicity. No account was given, however, of the respiratory state of the child. The importance of this has been shown in two recent studies. Sutherland *et al*¹ compared 16 patients with tricyclic antidepressant poisoning with 26