improve the detection rate and possibly aid recovery as a result of early diagnosis and subsequent management.

GORDON L MILLS

University Department of
Geriatric Medicine,
Royal Liverpool Hospital,
Liverpool L7 8XP

¹ Goldman, A, et al, *Journal of the Royal College of Physicians of London*, 1977, 11, 291.

Rugby injuries involving the hand

SIR,—With reference to the letter (10 February, p 414) by Drs R W Nutton and M J Guy, it is not stated what kind of dislocations occurred. The common injury is the posterior one, which is usually no trouble; it is the anterior dislocation that is liable to cause rupture of the middle slip of the extensor tendon. This is not generally realised and is not in many of the textbooks. X-rays are not always taken after this injury, reduction being done on the spot; so it may not be known what type of dislocation occurred. So if after this injury a patient cannot fully extend the proximal interphalangeal joint, then he should be treated at once in a lively splint to prevent the Boutonniere deformity, which is a disabling and difficult condition to treat once it has developed.

R F PAYNE

Accident and Emergency Department,
North Devon District Hospital,
Barnstaple,
Devon EX31 4JB

Baragwanath Hospital, 1978

SIR,—With reference to the "Letter from Soweto" (17 March, p 739), allow me the following comments. First of all, I wish to state that I think this article is a reasonably fair account of the situation as it pertains at the present moment at Baragwanath, but certain statements do require revision in the light of recent events and perhaps others require some explanation.

I believe it is now a matter of record that salary discrimination among the medical profession in South Africa will soon be a thing of the past. I in fact expect that when this letter is published this will have already occurred to a large extent.

With regard to the hospital facilities, it is indeed true that there is an enormous demand for hospital beds for the black section of our community. In the light of what I have to say further on, the reason for this may become more apparent. Mr Sakalo is correct in stating that a new hospital has replaced the old Johannesburg General Hospital. What he has not mentioned, and perhaps did not know, is that a 2000-bed hospital is being built at the Medical University of South Africa, and that the University of the Witwatersrand medical faculty and also the University of Pretoria medical faculty in association with the Medical University of South Africa are involved in its implementation and its personnel development.

Finally, allow me to highlight several sentences in Mr Sakalo's final remarks. He states, "Many of the patients are treated for afflictions and diseases resulting from poverty and overcrowding." A little later he remarks "Malnutrition, in particular, is endemic among the black population and a study by the University of Pretoria has shown that 75 black children die of malnutrition every day." These two remarks should perhaps be seen in

the context of some other remarks he makes: "The maternity wards sprawl over 50 hectares of land and cope with a 2.7% birth rate, which is a rate higher than India's"; and a little further on he states, "Even in this modern day, in one of the most advanced black societies, witch doctors still retain a strong hold on the people."

While in no way trying to minimise our concern with the infant mortality rate and malnutrition (primarily a paediatric problem), the concerted effort by all the authorities to inculcate a concern for family planning programmes and for proper infant care and feeding has met with resistance from the major proportion of our black citizens who still cling to the old customs. I nevertheless believe that the programmes which have been instituted by our Department of Health and our faculties must ultimately bear fruit as the sophistication of our black peoples evolve.

J N DE KLERK

Chairman,
Federal Council,
Medical Association of South Africa

Pretoria 0081, South Africa

Schizophrenia or obsessive-compulsive neurosis?

SIR,—We were interested to read the case of Dettol addiction described by Dr J S Khan and others (24 March, p 791). The patient's psychiatric symptoms were a fear of acquiring syphilis, associated Dettol drinking to "purify" herself, and a morbid cleansing obsession, including use of Dettol-soaked pessaries. The patient was diagnosed as having a schizophrenic illness. Schizophrenia and obsessive-compulsive neurosis can be difficult to differentiate¹ and recent work² has suggested that resistance is not an essential feature of the latter, making the distinction even less clear. The limited description in the paper would certainly fit an obsessive-compulsive neurosis, which is usually amenable to behavioural treatment.³ If the patient died as a consequence of compulsive cleansing rituals—chronic ingestion of Dettol—this would be almost unique.

E D GREENBERG

I M MARKS

Maudsley Hospital and
Institute of Psychiatry,
London SE5 8AF

¹ Stengel, E, *Journal of Mental Science*, 1944, 91, 166.

² Stern, R S, and Cobb, J P, *British Journal of Psychiatry*, 1978, 132, 233.

³ Marks, I M, Hodgson, R, and Rachman, S, *British Journal of Psychiatry*, 1975, 127, 349.

Proteinuria at high altitude

SIR,—In your leading article (24 February, p 508) you discuss the physiological changes that occur on ascending to high altitude, with an emphasis on proteinuria. Some of the data reported are open to criticism.

Pines¹ reported studies on proteinuria in the early morning urine specimens of seven people climbing at heights of up to 5890 m. However, the urine protein was measured by an inaccurate method (Bili-Labstix), urine volumes were not taken into account, and there were insufficient data to allow any statistical correlation of proteinuria with altitude, exercise, or the development of acute mountain sickness. Rennie and Joseph² completed a more detailed study of proteinuria at altitude. They showed a good correlation of proteinuria and altitude, but again only early morning

urine samples were tested and an unsatisfactory correction for urine dilution had to be made.

During a recent expedition to the Himalayas³ 17 doctors and scientists collected 24-hour urine samples over a 14-day period during an ascent to 5500 m. In the group as a whole there was no correlation between proteinuria and altitude, although some individuals had an increase in proteinuria at altitude. Proteinuria was of no value in predicting and did not relate to the development of acute mountain sickness.

The effects of altitude on the body and the development of acute mountain sickness are difficult to study because of the large individual variation and problems associated with sample collection. However, progress will not be achieved by inadequate data collection and drawing unwarranted conclusions. For instance, there have been at least five controlled trials at altitude of the drug acetazolamide (which is widely considered to be of value in the prevention of acute mountain sickness); yet not one of these studies was well controlled under routine trekking conditions, and use of this drug cannot therefore at present be recommended.

At a recent symposium, shortly to be published,⁴ many clinical and biochemical aspects of altitude and acute mountain sickness were discussed. However, the mechanisms, early diagnosis, and prevention of the illness remain an open question.

A BRADWELL
J DELAMERE

Immunodiagnostic Research
Laboratory,
University Medical School,
Birmingham B15 2TJ

¹ Pines, A, *British Journal of Diseases of the Chest*, 1978, 72, 196.

² Rennie, I D B, and Joseph, B, *Lancet*, 1970, 1, 1247.

³ Birmingham Medical Research Expeditionary Society, October 1977.

⁴ *Postgraduate Medical Journal*, 1979, suppl, in press.

Nutritional intake, adiposity, and diabetes

SIR,—Professor Harry Keen and others (10 March, p 655) appeared surprised that, among their subjects, those with a high energy intake had less adiposity; they also had lower blood sugar concentrations and better glucose tolerance, and would be less likely to develop diabetes. High energy intakes had been associated with high intakes of total carbohydrate.

Professor J N Morris and his colleagues had also been surprised to record that among their subjects high energy intake had been associated with lower incidence of coronary heart disease; this had occurred in men eating more cereal dietary fibre, but not that derived from vegetables and fruit.¹ It would be helpful if Professor Keen gave us information concerning the dietary fibre in his diets: dietary fibre is an essential nutrient.

These findings do not surprise doctors who have worked in tropical Africa and observed a low incidence of obesity, diabetes mellitus, and coronary heart disease among those eating large amounts of lightly processed, high-fibre starch foods.² The dietary fibre hypothesis of diabetes mellitus has two aspects, even if only the first aspect is usually quoted. (1) Dietary-fibre-depleted energy foods tend to produce obesity and mature-onset diabetes in susceptible phenotypes. (2) Large amounts of high-fibre energy foods improve diabetes mellitus and obesity.

Metabolic ward studies have demonstrated that weight-maintaining diets of unrefined cereals and starch-containing vegetables (75% energy) produced within 14 days remission of mature-onset diabetes in 85% of patients; they discontinued antidiabetic drug therapy.³ After discharge patients were advised to continue a maintenance diet of unrefined cereals and starch vegetables (60% energy) and dietary fibre (neutral detergent fibre 50 g/d). All nine patients who elected not to continue this maintenance diet relapsed quickly; on the other hand, 15 patients who took the maintenance diet have been observed for 6-48 months without relapse (Anderson, personal communication, 1979). Dietary details were stated briefly in the cited article³; more detailed diets can be obtained from the authors, Veteran's Administration Center, Lexington, Kentucky 40507, USA.

D P BURKITT

Unit of Geographical Pathology,
St Thomas's Hospital Medical School,
London SE1 7EH

H C TROWELL

Woodgreen,
Nr Fordingbridge, Hants SP6 2AZ

¹ Morris, J N, Marr, J W, and Clayton, D G, *British Medical Journal*, 1977, **2**, 1307.

² Burkitt, D P, and Trowell, H C, *Refined Carbohydrate Foods and Disease: Some Implications of Dietary Fibre*, p 227. London, Academic Press, 1975.

³ Kiehm, T G, Anderson, J W, and Ward, K, *American Journal of Clinical Nutrition*, 1976, **28**, 895.

Altered bile in diabetic diarrhoea

SIR,—We would certainly agree with Dr J H B Scarpello (6 January, p 58) that cholestyramine has not always been helpful in the treatment of diabetic diarrhoea. There are many reasons why this is so and the following case history demonstrates that in an individual patient there may be more than one cause of "diabetic diarrhoea."

A 48-year-old man who had had diabetes for 18 years was referred with a long history of impotence and a 12-month history of poor control of diabetes and weakness of the legs. He had recently developed an ulcer on the left heel. Seven weeks before admission the patient started to complain of intermittent explosive watery diarrhoea, sometimes nocturnal, and sometimes this led to faecal incontinence. On examination he had diminished sensation in the feet and absent ankle jerks. The left foot was colder than the right with absent dorsalis pedis pulse and a necrotic ulcer on the heel. His blood pressure was 130/80 mm Hg standing and 130/90 mm Hg lying down. Barium enema, sigmoidoscopy, and stool culture for pathogenic organisms were negative. The results of the bile salt analysis are given in the accompanying table. There was a reduction in the fasting pool size and an increase in the faecal excretion of bile acids; 22.7% of the total bile acids were in deconjugated form. The patient responded to tetracycline therapy but the diarrhoea recurred. A ¹⁴C-glycocholate breath test, which is a non-invasive indirect test of bacterial overgrowth in the small bowel, was negative during this attack.

This patient had a considerable quantity of free bile acids in the small bowel, a situation strongly suggesting bacterial overgrowth. The patient responded dramatically to tetracycline but the response may have been fortuitous

since the condition of diabetic diarrhoea is one of spontaneous remission and relapse. A further attack was not associated with bacterial colonisation, as assessed by a ¹⁴C-glycocholate breath test; yet the patient again responded to tetracycline. This test has not been reported to give false-negative results in the contaminated-small-bowel syndrome.¹

In conclusion, although there are many causes of diabetic diarrhoea in the diabetic, our results² suggest that cholestyramine may be more appropriately used in prophylaxis between attacks of the diarrhoea, when the pool size is larger than normal, rather than in the diarrhoea stage, when the patients have small pool sizes.

A MOLLOY
GERALD TOMKIN

Metabolic Unit,
Adelaide Hospital,
Dublin 8

¹ Thaysen, E H, *Clinics in Gastroenterology*, 1977, **6**, 227.

² Molloy, A, and Tomkin, G H, *British Medical Journal*, 1978, **2**, 1462.

Shortening hospital stay for psychiatric care

SIR,—Professor S R Hirsch and others (17 February, p 442) described brief hospital stay for psychiatric patients. I note that he did not include relapse rate.

The records of schizophrenic patients admitted to Claybury Hospital in 1960 and 1961 were checked by me. In the subsequent 10 years the chances of readmission within six months of any admission were greater if they stayed in the hospital less than six weeks.

ROBERT J DOIG

Cane Hill Hospital,
Coudsdon, Surrey CR3 3YL

Epilepsy and learning

SIR,—The letter from Dr C R Haines (31 March, p 893) highlights the situation in which many school doctors find themselves where epilepsy is concerned. This association has urged in our reply to the Warnock Report that the problems presented by the trend towards confidentiality be overcome because the otherwise valuable role of the school doctor is hampered when information passes only between consultants and the general practitioner.

We have, therefore, endorsed any scheme whereby the specialist in community medicine (child health) would co-ordinate the flow of information between hospital, GPs, and the school doctors. We would hope that parents could be persuaded to give permission for dissemination of information in this way, and that each local education authority would promote such a scheme.

This association believes that a well-informed school medical officer may be a more appropriate person to hold the assessment files for children with special needs as envisaged by Warnock than the head teacher. In any event,

the school doctor should have access to these records as of right. We would hope that such an arrangement would provide teachers with the sort of practical help they need in meeting the special—and changeable—needs of children with epilepsy.

ANDREW GORDON CRAIG

Wokingham, Berks RG11 3AY

Schizophrenia-like psychosis after treatment with beta-blockers

SIR,—The cases reported by Drs J Steinhert and C R Pugh (24 March, p 790) of schizophrenic-like psychosis in patients treated with beta-blocking agents were particularly interesting, for although these drugs are widely used in cardiology they are being used increasingly for their peripheral effects. It is noteworthy that in the cases reported the onset of psychotic symptoms followed increase of dosage.

The authors make the point that it is uncertain whether beta-blockers have a central action, and this question has been asked ever since the drugs were discovered. Three possible modes of action can be envisaged. (1) By penetrating brain tissue these drugs may interrupt the flow of information channelled through β -adrenergic mechanisms. (2) They may have a direct action on sensitive neuronal networks independent of a specific beta-blocking property. (3) The possibility exists that a peripheral effect may be transmitted via neural or humoral channels, or both, to the central nervous system, becoming manifest in psychic and behavioural change.

There is evidence particularly from work on animals that central adrenergic systems exist which are involved in functional activities related to adjustment of awareness and arousal, learning processes, and regulation of muscle tone. There is also evidence that adrenergic receptors of the beta type exist in postsynaptic membranes, which are normally activated by adrenergic substances, but inactivated and made unresponsive by beta-blockers. Noradrenaline is released by and taken up by adrenergic terminals and because of its postsynaptic effects serves as an information carrier. Dahlström and Fuxe, by demonstrating the existence of monoamine-containing neurones in the central nervous system in mammals, have paved the way for the discovery of catecholamine-containing cell bodies, or "nuclei," which give rise to descending and ascending catecholaminergic fibres. While there is an extreme degree of divergence in these noradrenergic systems, they include in their territory a huge number of cells. Certain portions of these systems seem to be involved in the control of the level of vigilance, which is in keeping with the anatomical structure of the locus ceruleus, which has relatively few cells capable of establishing contact with numerous cells in the forebrain and neocortex. Electric stimulation of this area in cats induces electroencephalographic arousal, accompanied by dilatation of the pupil, contractions of the nictitating membrane, a rise in blood pressure, and increase followed by decrease in the heart rate.

While data thus far available do not yet permit the exact site and mode of action by which beta-blockers exert their influence on the central nervous system to be identified, it is interesting to note the work of Atsmon *et al.*,¹ who treated a psychotic woman with porphyria in Tel Aviv with large doses of propranolol for her hypertension. The treatment produced

Bile salt pool size and composition in the patient and a control (mean ($\pm$ SE of mean)) of recently diagnosed diabetic patients without complications

Glycine:taurine ratio	Trihydroxy: dihydroxy ratio	Fasting pool size (mmol)	Free bile acids (% of total)	Faecal bile acids (mmol/24 h)
3.3 (2.5 $\pm$ 0.4)	1.00 (1.25 $\pm$ 0.1)	5.544 (6.672 $\pm$ 0.389)	22.7 (none detected)	2.225 (0.900 $\pm$ 0.180)