more recently have contained small numbers (less than 24 organisms per ml) of spore-bearing *Bacillus* species.

While we have no evidence that any patient became infected by pseudomonads of these species, it would be unwise to assume that they may not take on an opportunistic role in patients who are immunologically impaired. We would therefore recommend that control of infection teams are alive to this possibility, that hospital water is from time to time monitored microbiologically, and furthermore that domestic hot water in general hospitals is maintained at a temperature adequate to kill vegetative bacteria.

H J BLACK
EDWARD J HOLT
K KITSON
M H MALONEY
D PHILLIPPS

Royal Infirmary,
Huddersfield, W Yorks HD3 3EA

¹ Department of Health and Social Security, *Hospital Services Engineer* No 30. London DHSS, 1976.

The wrong drug

SIR,—I read your leading article "The wrong drug" (12 May, p 1233) with interest. It seems to me that you left out the problem of drugs prescribed by hospital doctors at outpatient clinics and by general practitioners.

It has been my practice to have a look at the drugs I prescribe for my outpatients. Every time they come to see me I ask them and their relatives to bring the drug bottle. This has several advantages. The doctor gets an opportunity, firstly, to see whether the patient is getting the right drug or drugs and the right dosage and to find out what other drugs the patient is taking, as patients do receive treatment from more than one doctor for different ailments and the communication between doctors does break down; secondly, to see whether the patient is taking them as prescribed; thirdly, to discuss the side effects of the drugs with the patients and relatives and to point to which drug is causing what side effects (this is particularly useful when the patient is on more than one drug); fourthly, to be confronted by different types of drugs, which by their sheer number may have some sobering effect on the prescription habit; and, finally, to establish a better doctor-patient relationship by giving the patient the impression that he does care by checking the medicine he orders.

D CHAKRABORTI

Windsor Unit,
West Norfolk and King's Lynn General Hospital,
King's Lynn, Norfolk

Falciparum malaria despite chemoprophylaxis

SIR,—The letter from Dr S J Bentley continuing the discussion about partial suppression of falciparum malaria by chemoprophylaxis (19 May, p 1351) raises some important points concerning suitable prophylactic measures.

Having spent three and a half years in the Gambia, I fully endorse his statement that pyrimethamine in a dose of 25 mg weekly does not give adequate prophylactic cover. An increase of the dose to 25 mg every five days or 50 mg weekly is usually fully effective. There is also an increase in resistance to proguanil (Paludrine) in a daily dose of 100 mg. My wife has the dubious distinction of being the first

recorded case of malaria in the Gambia in a subject taking regular proguanil. I am not altogether happy about the suggestion of taking chloroquine as a prophylactic. This is a very valuable therapeutic agent for the treatment of overt attacks of malaria and its widespread use as a prophylactic agent is likely to speed up the development of resistant strains of plasmodia. These are already widespread in parts of the Far East.

A further point to consider is the question of chemoprophylaxis in women starting a family. Pyrimethamine, as a potent inhibitor of folate metabolism, is not to be recommended. I have seen the detrimental effects of chloroquine when continued throughout pregnancy. Many of the expatriate community in the Gambia used chloroquine and, although this was by no means a controlled study, many women had small babies, delivered early, and—in one case at least—with auditory defects, probably attributable to the drug. Martindale's *Extra Pharmacopoeia* recommends proguanil as the drug of choice for malaria prophylaxis in pregnancy, but what advice should be given to women in areas where there is marked resistance to this drug?

It is very important that those travelling abroad, whether on business or on holiday, should receive advice about malaria prophylaxis that is up to date. Such advice is very important but is not easily come by. Chemoprophylaxis is very effective, if the correct drug for the situation is taken in an adequate dose and continued for four weeks after exposure.

P MOODY

Pharmaceutical Department,
Preston Hospital,
North Shields, Tyne and Wear NE29 0LR

One man's schizophrenic illness

SIR,—I was very impressed by the excellent description presented by Mr Peter Wescott (14 April, p 989) in his personal paper "One man's schizophrenic illness." After reading his report a second time, I had the strong impression that Mr Wescott, despite the fact that he can write so well and present his symptoms in such a comprehensible manner, could possibly be doing even better if his medication regimen were modified.

Mr Wescott's description of the "four suicidal attempts and some period of suffocating depression" suggests that his diagnosis should possibly be modified to be that of schizoaffective illness, or what Dr Carlo Perris of Sweden prefers to call cycloid illness. If Mr Wescott had been treated at our medical centre, he would have been on an adequate course of lithium; small dosages of a neuroleptic would have been added only if he had shown an insufficient response to lithium. I am quite sure that the same treatment would have been rendered in Dr Perris's medical faculty. The reason for my writing this letter is that Mr Wescott sometimes feels "that the richness of my pre-injection days—even with brief outbursts of madness—is preferable to the numbed cabbage that I have become." He goes on to say, "... once I lived in a fascinating ocean of imagination, I now exist in a mere puddle of it." These excerpts are excellent examples of the side effects of neuroleptics and at the same time indicate that he should have the opportunity of an adequate exposure to lithium therapy, only being given neuroleptics in very small doses if the lithium falls short in the therapeutic goals.

Mr Wescott's comments about the "family GP" and about psychiatrists in general should be a teaching lesson for all of us psychopharmacologists. I hope that Mr Wescott has the opportunity to read a review of this problem by Carlo Perris.¹ I not only wish Mr Wescott the very best in the future but also hope that he has the opportunity to return to the richness of his pre-injection days.

DON GALLANT

Department of Psychiatry and Neurology,
Tulane University School of Medicine,
New Orleans, USA

¹ Perris, C, *Archives of General Psychiatry*, 1978, **35**, 328.

How to use an overhead projector

SIR,—Dr T S Murray's (3 March, p 602) "How to do it" article on the overhead projector missed one unique and very versatile feature. It is possible, by use of a xerox machine or similar copier, to produce projectable transparencies from books, drawings, tables, or EKGs in a matter of seconds. An 8½ × 11 inch sheet of acetate or similar material is simply substituted for paper in the machine. The resulting transparency is obtained quickly and inexpensively and is usually of high quality.

JOHN ROBBINS

Primary Care Centre,
University of California, Davis,
Sacramento,
California 95817, USA

Doctors and children's teeth

SIR,—I should like to raise an additional point regarding your recent leading article "Doctors and children's teeth" (12 May, p 1231).

In the last 25 years as a psychiatrist working with children I have often questioned a mother about the attitude of the father or stepfather towards the child, and only too frequently the answer is: "Very good, doctor. He always brings him sweets when he comes home." This social pattern of making sugar equate with affection must be altered in any fight against dental caries or obesity, or any other illness caused by excessive consumption of sugar.

J VINCENZI

Colchester, Essex CO6 2LE

Drugs for epilepsy

SIR,—I refer to the article on epilepsy by Drs D F Smith and J C Mucklow (14 April, p 1000). The statement that phenytoin is now generally regarded as the anticonvulsant of first choice in treating tonic-clonic epilepsy is not entirely accepted by most dealing with epilepsy. It is indeed a broad-spectrum anticonvulsant but there are many problems with its use in practice, particularly in younger patients. The disfiguring skin manifestations like acne, hyperkeratosis, hirsutism, etc, are certainly unacceptable.

In the case mentioned the combination of phenytoin, primidone, and clonazepam is, I think, against the modern practice of single-drug therapy. Clonazepam is a very weak anticonvulsant when used orally. It is very essential to classify the type of epilepsy in such cases and then a suitable single drug should be tried in

effective therapeutic dose before adding another drug. Sometimes it is also essential to check on biochemical insults like hypocalcaemia.

Carbamazepine is probably the drug of choice today because of its effectiveness in tonic-clonic and temporal lobe seizures and its wide range of safety in long use.

P S MEHTA

Stepping Hill Hospital,
Stockport, Cheshire SK2 7JE

Biochemical testing for medical emergencies

SIR,—The value of biochemical testing in medical emergency admissions was effectively discussed in the article by Drs B T Williams and R A Dixon (19 May, p 1313).

It will always be a problem deciding whether certain tests are "cost effective," but I feel that it would be detrimental to medicine as a whole if clinicians were only allowed to use certain biochemical investigations because statistically they are cost effective. It may be that certain tests are very costly to perform in terms of time and expertise, but I feel that it should be left to the judgment of the clinician to decide whether an investigation will help his particular patient.

Perhaps it is more important to train doctors to discriminate between the value of certain investigations rather than to enforce strict rules. Teaching hospitals are obviously at fault in that they tend to overinvestigate, as shown in the study; subsequently a junior doctor continues to follow this practice when he works as a house officer, and so it is perpetuated.

DAVID COOPER

University of Manchester
Department of Medicine,
Hope Hospital,
Salford M6 8HD

Pressure on the tracheal mucosa from cuffed tubes

SIR,—I would like to comment on two important aspects of the paper by Drs J M Leigh and J P Maynard (5 May, p 1173) relating to the use of endotracheal tubes for general anaesthesia.

Firstly, the seal point of a cuff can be very difficult to determine accurately. At that point the addition of small increments of air to the cuff of a Magill red rubber tube (or any other low residual volume cuff) may lead to a steep rise in pressure on the tracheal wall. It is much easier to detect when a small leak past the cuff is occurring than to detect the point at which it has just disappeared. If this leak is just present at an airway pressure of, say, 20 mm Hg then the mean pressure on the tracheal wall must be less than that figure, on the basis of the principle of the Starling resistor.¹ Equally, for fluid to pass the cuff from above, then the pressure in the pharynx must rise above 20 mm Hg, clearly a very rare event. If an anaesthetist is using a "minimal leak" to ensure low tracheal wall pressures, then disappearance of it would alert him to a possible rise in intra-cuff pressure, which can be due to the diffusion of nitrous oxide into the cuff.² Necessary measures can then be taken.

Secondly, Drs Leigh and Maynard consider replacing all red rubber tubes with Lanz

endotracheal tubes, but quite rightly raise the question of economics. Therefore we need to study the controlled laboratory and clinical trials of this new tube that clearly demonstrate its superiority over the original. To my knowledge, there are none and this is the same for most other new disposable endotracheal tubes. Yet if a drug company were trying to introduce a new potent analgesic we would insist on seeing the evidence of its superiority over morphine and other established drugs as shown by extensive laboratory and clinical trials. With invasive equipment the situation is different. We accept manufacturers' claims for their products, which are often extravagant and without any hard evidence of the products' being better than existing types. In fact there is work indicating that a Magill red rubber tube used correctly may be no worse than a disposable, high-residual-volume cuffed alternative.¹

Would it not be best, however, to teach people how to inflate cuffs correctly, thus eliminating the need for elaborate devices to prevent overinflation?

W G NOTCUTT

Department of Anaesthetics and Intensive
Care,
University of the West Indies,
Kingston 7, Jamaica

¹ Homi, J, et al, *British Journal of Anaesthesia*, 1978, **50**, 435.

² Revenas, B, and Lindholm, C E, *Acta Anaesthesiologica Scandinavica*, 1976, **20**, 321.

Immersion and thermoregulatory response

SIR,—We were interested to read the report by Dr M G Hayward and Professor W R Keatinge (5 May, p 1182) of an individual who showed little thermoregulatory response to immersion in cool water, but wonder whether evidence of this nature can account for unexplained diving accidents.

At a recent meeting of the Physiological Society¹ we presented data on three long-distance swimmers, obtained first during experimental immersion in a tank of cold water and then during a long-distance swim in Windermere. Two of the swimmers, who had trained in cold outdoor water, in the tank behaved like Hayward and Keatinge's subject (although the water temperature of 12°C was much colder); they showed relatively slight shivering and their deep body temperatures fell at 1.2 and 3.8°C/hour. The third, who had trained in indoor baths, showed a higher oxygen consumption and a slower rate of body cooling (0.6°C/hour). During the swim, the man who had trained in warm water cooled more rapidly than did the other two, although his oxygen consumption was still higher than theirs. He was forced to give up sooner. Our data suggest that people who have been repeatedly exposed to cold while swimming may show an advantage when tested in a similar way: they may not do so when tested immobilised in a tank of cold water. We also have unpublished data on individuals who show little metabolic response to immersion in cold water in a tank, but who cool slowly.

It seems curious to use the word acclimatisation, which usually implies improved responses, to describe a situation where heat loss on cold exposure was accelerated. It may be that testing in a more comparable situation would have shown improvement rather than

disadvantage; even if not, "disadvantageous acclimatisation" would seem likely to become apparent at an early stage, before serious hazard could arise.

We also wonder if it is the saturation diver who would be at risk since he is not usually exposed to severe cold, as Department of Environment regulations require external heat to be applied to the body in dives in excess of 50 m. Exposure to cold is commoner in the "bounce" (subsaturating) diver who dives in a dry suit to spend considerably shorter periods at working depth, and may lose heat in each of successive dives.

We have found great variation between individuals in their response to immersion in a tank of cold water. We are convinced that further study, and measurements in realistic physiological states, are needed to make predictions for the real situation.

I F G HAMPTON
DOUGLAS SMITH

Department of Physiology,
School of Medicine,
Leeds LS2 9JT

F St C GOLDEN

Institute of Naval Medicine,
Alverstoke, Hants

¹ Golden, F St C, Hampton, I F G, and Smith, D, *Journal of Physiology*, in press.

Medical representatives

SIR,—I read with interest your article (28 April, p 1134). While I agree with Drs T B Binns and Andrew Smith's enlightened comments on the subject, I would take issue on some points.

Firstly, in my company we do not discourage our representatives from "recommending someone else's product." The essence of good ethical representation (in pharmaceuticals or any other business) is to position your product to the needs of the individual. To start using a product on the recommendation of a medical representative demands trust and confidence in what the representative says—this in turn means believability. Under these circumstances if the representative, with his knowledge of his own and other competitive products, feels another product is more applicable then I would certainly expect him to refer to it.

Secondly, no doubt some medical departments will respond to Dr Binns's comments. My own experience in four major pharmaceutical companies is that medical information departments, whether they are part of the sales department or not, are totally impartial, since patients' lives may depend on their response. My previous comments regarding credibility are even more relevant in this case. Thirdly, rather than, as Dr Binns suggests, complain to the managing director of the company about representatives' misdemeanours, the clinician is likely to get a quicker response by contacting the sales manager.

Finally, we are all conscious that there is a patient at the end of all this. Unlike the commercial traveller or "quack doctor" we are not here today and gone tomorrow. The pharmaceutical industry is an integral part of community medicine and commercial development in this country and has to have, and rightly so, a "whiter than white" image at all levels.

P T HUGHES

Bristol Laboratories and Mead Johnson,
Slough, Berks SL3 6EB