Manipulation trials

SIR,—Every doctor and most laymen are well aware of patients who have been treated for backache for months, even years, within the NHS and have finally been adjured to "live with it." A proportion of such patients secured rapid relief from a lay manipulator. It was this well-recognised fact that let me teach spinal manipulation (among other techniques) to my physiotherapy students from 1940 to 1970. The intention was that those who needed this treatment should get it immediately, from skilled personnel, and in full view within the hospital rather than wander off in the end on their own to try their luck with some self-styled layman. This policy worked very well, many patients recovering swiftly—in half of all cases, backache recovered in one session of manipulation.

It is very noble of Dr H Sims-Williams and others (24 November, p 1318) to have put up the controlled trial that they describe. The result was negative; but, firstly, the patients were not selected as suited or unsuited to manipulation; secondly, they were treated by Maitland's adaptation of Balfour's¹ (1819) and Récamier's oscillatory techniques² (methods with scant resemblance to those taught by me³); and, thirdly, they were all assessed at the end of a month.

Physiotherapy (that is, heat and exercises) for backache is useless. Their article states, "The management of non-specific back pain is unsatisfactory and physiotherapy is often prescribed empirically." On what grounds? No patient should be treated until the cause of his symptoms becomes clear, and the measure prescribed should be appropriate to the diagnosis. If manipulation were confined to suitable cases, proper methods were employed, and the assessment were made after a few days a very different picture would emerge. Meanwhile, doctors are discouraged by papers like this one from adopting sensible measures and the jubilant lay manipulators cash in again.

JAMES CYRIAX

London NW1

¹ Balfour W. *Compression and percussion*. Edinburgh: Hill, 1819.

² Récamier M. *Rev Med Fr* 1838;1:74.

³ Cyriax J. *Textbook of orthopaedic medicine*, 10th edn. London: Baillière Tindall, 1980.

Assessing improvement of low back pain

SIR,—In their trial of mobilisation and manipulation for low back pain (24 November, p 1318), Dr H Sims-Williams and others have used as the principal criterion for improvement the patients' subjective assessment of pain. There are four categories: completely better, much better, slightly better, and no change (with two main categories if the pain is worse).

In 1974 Glover *et al*¹ published the results of a trial of manipulation for low back pain in which a different method was used. He asked the patients to consider their original pain as 100% and then asked them to say what percentage pain they had when it improved. He found that this correlated remarkably accurately with objective ways of measuring improvement (such as the angle of straight-leg raising as measured with a goniometer).

Since I first knew of Glover's work I have asked all my patients to express their improvement (or exacerbation) in percentage terms. It is surprising how readily patients will do this if

one persists. The most interesting observation I have made is that patients who describe their pain as "much better" give a figure of their remaining pain which varies from as much as 80% to as little as 15%. If this is used as the only criterion of improvement it quite obviously gives an invaluable guide for management as well as research.

M E BURLEIGH CARSON
Secretary General, International
Federation of Manipulative Medicine

London W1

¹ Glover JR, Morris TG, Khosla T. *British Journal of Industrial Medicine* 1974;31:59-64.

Rehabilitation

SIR,—If I may join in a debate on rehabilitation (8 December, p 1508), I would regard this as the final stage in any successful or semi-successful course of treatment in any specialty. Unfortunately, just as the mechanisms of, for instance, injuries attract little interest, the final stages of an illness also lack the glamour which appears to lie in the acute sector.

There has been far too little evaluation of physiotherapy and occupational therapy and I personally have my doubts about whether occasional visits to the physiotherapy department are of much value over a wide range of conditions, whereas I can see the value of "country house" rehabilitation where many of the inmates in fact, unknown to themselves, engage in rehabilitation activities.

Rehabilitation today should really be the interest of whichever specialty has been involved in the acute phase, but the biggest disasters which we face in this field are the lack of interest shown by our medical teachers in rehabilitation and the linkage of rehabilitation and rheumatology. It could be argued that rehabilitation in rheumatology is merely attempting to slow down a steadily degenerative process, whereas attempting to get patients back to their former level in activities which require a certain degree of recovery is true rehabilitation in its own right.

P R J VICKERS

Newcastle upon Tyne NE3 4AP

Intermittent venous sampling without heparinisation

SIR,—Drs M J G Farthing and Anna M Mattei (3 November, p 1112) report the use of a silicone elastomer cannula for intermittent venous sampling without heparinisation. Over the past three years we have successfully used a flexible polytetrafluoroethylene (Teflon) catheter (Abbocath-T, Abbott Laboratories) for this purpose in approximately 80 patients undergoing endocrine and platelet function tests.¹⁻³

A peripheral vein is cannulated with an 18- or 20-gauge Abbocath-T and a three-way tap is connected. The system is kept patent with sterile 0.9% saline. At the time of sampling the first few millilitres of aspirate are discarded before collecting the test sample. Even with 20-gauge catheters clotting in the lumen and haemolysis of the sample are extremely rare occurrences provided that the catheter is well flushed and filled with saline after blood collection. In one haemophilic patient such a catheter remained patent for six days despite

regular cryoprecipitate infusions and blood samplings.

These catheters are readily available and the technique is simple and reliable.

D P MIKHAILIDIS

Department of Chemical Pathology,
Royal Free Hospital,
London NW3 2QG

RONALD A HUTTON

Haemophilia Centre,
Royal Free Hospital,
London NW3 2QG

¹ Hutton, R A, *et al*, *Journal of Clinical Pathology*, 1979, 32, 434.

² Hutton, R A, *et al*, *Journal of Clinical Pathology*, 1979, 32, 1192.

³ Hutton, R A, *et al*, *Thrombosis Research*, in press.

Erythema something or other

SIR,—I was intrigued to read of yet another named disease associated with erythema nodosum (17 November, p 1263).

I feel that life in general medicine, as well as in my own specialty of dermatology, would be simpler if erythema nodosum, erythema multiforme (the definition of which in the United States includes the commonest of the erythemas—namely, toxic erythema), erythema marginatum, erythema annulare (whether centrifugal or fixed), and erythema you-name-any-of-the-dozens-of-others were to be considered causally as similar. Certainly this would be true when one considers most patients at the bedside.

I would like to challenge, moreover, someone to find an erythema that has *not* been associated with any of the common aetiological factors, which include infections of just about any micro-organism that one would care to name, drugs (most particularly the penicillins and sulphonamides), connective tissue disorders, neoplasms, various sera and inoculations—and one can probably complete the list by mentioning sarcoidosis.

PETER COPEMAN

Department of Dermatology,
Westminster Hospital,
London SW1P 2AP

A and E departments for children

SIR,—I read with interest your articles on accident and emergency services. I would like to support very strongly the letter from Dr Cynthia Illingworth regarding paediatric casualty departments (24 November, p 1364).

Since the closure of the Evelina Children's Hospital we have had a separate area of Guy's Hospital where only children are seen. The nursing staff are trained children's nurses and the casualty officer is always one of our senior house officers, who spend four months in the department as part of their year's rotation. We see approximately 14 000 children a year; half have had genuine accidents, the majority quite trivial; and the other half present predominantly medical and social problems. To help us deal with the latter we have very close liaison with health visitors. In our area the children's casualty department is adjacent to the adult casualty department. We have very good relationships with the adult services and frequently call on the casualty officer in the adult department to assist us with minor fractures, etc.

LEO STIMMLER

Paediatric Department,
Guy's Hospital,
London SE1 9RT