CORRESPONDENCE

Mortality of male doctors and reductions in cigarette smoking M R Alderson, MD.....	108	Incidence of malignant melanoma of the skin in England and Wales P W L Clough, PHD; N P Warren, MB; M Sutton, FRCP.....	112	Medically qualified preclinical academics N A Green, FRCS, and R E Coupland, MD... 116
Hypertension and general practice D G Beevers, MRCP; D M M Thomson, MB, and D C Rawlins, MB; S L Barley, MRCP... 108		Sunshine and malignant melanoma A A Epenetos, MRCP; A J Swerdlow, BM... 112		Revised consultant contract M K El Dars, DM; B B Milstein, FRCS.... 116
Peripheral nerve lesions in the muscle compartment syndrome J M S Pearce, FRCP.....	109	Supplies of Trilene A Barnes.....	113	Professional standards for consultant appointments B Thalayasingam, MRCP.....
Medical charities, prevention, and the media Jennie Popay, MA, and P A Draper, MFCM... 110		Poststreptococcal glomerulonephritis C G Isles, MB, and J M Boulton-Jones, MRCP 113		Short-term certification G M Watts, MRCS.....
Drug-induced oesophageal injury J D R Rose, MRCP, and G B Tobin, MRCP... 110		Epileptic drugs and the police J A Riddle, MB.....	113	Deference, authority, flirtation, and stealth A P J Ross, FRCS; R Pollard, FRCS..... 117
Reflux oesophagitis P I Reed, FRCP.....	110	Sustained-release lithium tablets and renal function P Vestergaard, MD.....	113	Health centres and private practice G M Coleman, LRCPT.....
The undiscovered country M W Flowers, FRCS.....	110	Psychoanalysis, psychotherapy, and health services S Gold, FRANZCP; Anne L Zweig, MD.... 113		Academy of medicine W J Stephen, MB.....
Manipulation trials J H Cyriax, MD.....	111	Paracetamol concentrations, hepatotoxicity, and antidotes J E Gilligan, FFARCS, and others..... 114		Health visitors and confidentiality B D M Williams, MRCP.....
Assessing improvement of low back pain M E B Carson, LMSSA.....	111	Labetalol and priapism M R Law, MRCP, and others.....	115	Multidisciplinary teams Lea Pearson, BA.....
Rehabilitation P R J Vickers, FRCS.....	111	Wanted: a new wound dressing J Bunyan.....	115	Participation in national quality control schemes D N Baron, FRCPATH.....
Intermittent venous sampling without heparinisation D P Mikhailidis, MB, and R A Hutton, PHD 111		Surgery in outpatients H G Calwell, MD.....	115	Points Blood pressure measurement (A J Cossor); A mild traumatic lesion (J H Gervis; D A Varvel); Recording the whoop (P H Tattersall); Subsidy of social functions by drug companies (D C Wilkins; C P Kent); Towards fewer handicapped children (M R Bahl); Accident and emergency services (O Ahmed); Once an alcoholic, always an alcoholic; BMA at Hong Kong (Doreen H Annear)..... 119
Erythema something or other P W M Copeman, FRCP.....	111	Parenteral nutrition before surgery? J F L Shaw, MB.....	115	Correction: Diagnostic kits and the clinical chemist (Warner)..... 119
A and E departments for children L Stimmler, MD.....	111	Diagnostic kits and the clinical chemist M S Walker, PHD.....	115	
Long-term urethral catheter drainage E S Glen, FRCS.....	112	Monitoring blood glucose R D Eastham, FRCPATH.....	116	

We may return unduly long letters to the author for shortening so that we can offer readers as wide a selection as possible. We receive so many letters each week that we have to omit some of them. Letters must be signed personally by all their authors. We cannot acknowledge their receipt unless a stamped addressed envelope or an international reply coupon is enclosed.

Correspondents should present their references in the Vancouver style (see examples in these columns). In particular, the names and initials of all authors must be given unless there are more than six, when only the first three should be given, followed by et al; and the first and last page numbers of articles and chapters should be included. Titles of papers are not, however, included in the correspondence section.

Mortality of male doctors and reductions in cigarette smoking

SIR,—The provocative paper by Mr P N Lee (15 December, p 1538) presents data on a very important issue and it is nice to read of attempts to utilise the Registrar General's publications. However, there are four points that should be borne in mind before attempting interpretation of his material.

Firstly, is there variation in the quality of the data, over time and between the groups being contrasted? Secondly, comparison cannot be made directly between standardised mortality ratios for different periods where the standard rates have altered—for example, the male doctors' mortality rates increased in the period 1931-51¹ and did not decline as suggested by Mr Lee. Thirdly, there have been changes in the recruitment both to the medical profession and to the categories making up social classes I and II between 1951 and 1971. Finally, there has been a reduction

in the average number of cigarettes smoked in social classes I and II between 1958 and 1971² and yet their suicide rates have not increased.

Mr Lee's conclusions have been extrapolated way beyond the material presented; he does not give any information on why people smoke or on what happens to them when they cease—though Fletcher and Doll³ concluded that doctors who stopped smoking were, in general, aware more of benefits than of any sense of deprivation, either physical or psychological. The above points do not lend any credence to the conclusions that giving up smoking has "adverse consequences."

MICHAEL ALDERSON

Division of Epidemiology,
Institute of Cancer Research: Royal General Hospital,
Sutton, Surrey SM2 5PX

¹ Adelstein AM. *Br Med J* 1978;ii:983-7.

² Todd GF. *Social class variation in cigarette smoking and in mortality from associated diseases*, Tobacco Research Council Occasional Paper No 2. London, TRC, 1976.

³ Fletcher C, Doll R. *Br J Prev Soc Med* 1969;23: 145-53.

Hypertension and general practice

SIR,—Dr M H F Coigley and Nurse Greta Barnes (15 December, p 1582) are to be congratulated on their impressive results in the detection and management of hypertension by participants in the Medical Research Council trial of mild-to-moderate hypertension. As one of the first contributors of patients to this study¹ I am glad that things are going so well.

The question which arises is whether such practices, like that in Stratford, are representative of general practice as a whole. Most practices are not involved in undergraduate

teaching or medical research and the picture is, sadly, very different when one looks at the state of affairs in more typical situations. Indeed, irony of ironies, in the very same issue of *BMJ* as the letter by Dr Coigley and Nurse Barnes is an important report from Brighton (p 1556); of the hypertensives detected in that screening survey, only half had previously been diagnosed. The experience in Brighton in 1979 is almost identical to ours in Renfrew in 1972,² and an even more woeful picture has been found in London.³ The "rule of halves" still holds in this country: of the hypertensives of this nation, about half are detected; of those who are detected, half are receiving treatment; and, of those who are being treated, only half have adequate blood pressure control. There remains therefore a serious deficiency in the present methods of detection and care of patients, many of whom would benefit from antihypertensive therapy.^{4 5}

Concerted action by the whole medical profession is necessary but hospital doctors cannot help much as only a minority of the population attend hospitals. Special screening units such as mine may help but they are expensive and are primarily for research. The third option is that as about 75% of the population visit their doctor at least once in three years⁶ this is the most feasible place to detect and treat hypertensives. Our studies in three general practices in Scotland⁷ have confirmed that this method of management is acceptable and efficient.

Despite the impressive findings of the Stratford group, I am afraid that we still have a long way to go before the national picture is half as good as theirs. Everything in the garden is not lovely, and I am unrepentant for my original comments (3 November, p 1137). Research-minded general practices are few and far between and their data on the state of detection must be biased, if only because of their abnormal excellence.

Why is it that one is always frightened to criticise general practice for fear of offending people despite the fact that, with notable exceptions, standards are often depressingly low? In hospital practice and in medical research one is constantly under a barrage of criticism both from without and from within, and this is to be welcomed as a stimulus and not denied hotly. We know our garden is not lovely.

It is a pity to see the pages of an international medical journal such as this containing personal comments about contributors to medical debates. I think this should be avoided. I do hope that the answer of William Pitt to Walpole when he was accused of the crime of youth does not obtain here (I am much older than Pitt was at that time): "The atrocious crime of being a young man . . . I shall attempt neither to palliate or deny, but content myself with wishing that I shall be one of those whose follies shall cease with their youth, and not of that number who are ignorant in spite of experience."

D G BEEVERS

University Department of Medicine,
Dudley Road Hospital,
Birmingham B18 7QH

¹ Beevers DG, Nelson CS, Duncan S, Padfield PL. *Postgrad Med J* 1976;32:683-6.

² Hawthorne VM, Greaves DA, Beevers DG. *Br Med J* 1974;iii:500-3.

³ Heller RF, Rose G. *Br Med J* 1977;i:1442-4.

⁴ Hypertension Detection and Follow-up Programme. *JAMA* 1979;242:2562-71.

⁵ Hypertension Detection and Follow-up Program. *JAMA* 1972;242:2572.

⁶ Beevers DG, Hawthorne VM, Sinclair RG. In: Berglund G *et al.* eds. *Pathophysiology and Management of Arterial Hypertension*. Molndal: AB Hassle, 1975:263.

⁷ Barber JH, *et al.* *Br Med J* 1979;i:843-6.

SIR,—We read with interest the account of the screening exercise carried out in Hove and reported by Dr J A P Trafford and others (15 December, p 1556).

In January 1977, with the help of the Medical Research Council, we screened patients in our practice between the ages of 35 and 64 years. This was in a caravan parked outside the practice premises. Unfortunately, snow at the end of the period made access from the surrounding villages more difficult and helped cut down attendances. The results were as follows. Patients attended—1067 out of 1174 in age range (90.9%); hypertensives (diastolic 90 mm Hg or over)—175 (16.4%); previously known hypertensives—85. By close co-operation with the unit and continuous updating of the addresses it was possible to contact the vast majority of the patients in the age group under study. This advantage would have been lost had strangers attempted the exercise.

The total cost was about £850 and less than £9.50 for each new hypertensive diagnosed. The majority of the new hypertensives were entered into the Medical Research Council's trial of the treatment of mild-to-moderate hypertension. So far as we know, there is little evidence to support routine treatment of symptomless hypertensives aged over 65 years. We are at present rescreening our practice, including new patients in the age group. The attendances at the unit are of a similar order. The costs have risen to about £1 for each patient screened. The patients diagnosed have been investigated and treated in general practice. This is a further large saving on the resources of the National Health Service compared with Hove, where they appear to have been referred automatically to hospital.

We understand that the MRC still has some vacancies for medium-sized practices in the trial and is willing to help screen patients from selected practices. We would suggest that if further funds are available for screening for hypertension the screening techniques developed by the MRC should be the basis for future developments as they are effective and economical.

D M M THOMSON
D C RAWLINS

Bath BA3 5PG

SIR,—The whole philosophy behind the paper of Dr J A P Trafford and others (15 December, p 1556) must be strongly challenged, for they are describing a foray into the community by a hospital-based team whose methods, assumptions, and findings run contrary to the way in which general practice is moving.

The original work of Hart¹ has shown convincingly that case finding in the course of normal general practice is the preferred method for patients with hypertension. What Hart did—and many others since have done—was to try to take the blood pressure of every patient (within set age limits) attending the surgery and after some time to contact the remaining patients who had no blood pressure reading in the notes. In this way he covered 98% of the age group 20-64 years.

We have recently done the same in our practice. After four years of trying to take the

blood pressure whenever patients attended, we were left with 707 who had no reading recorded in the notes. Our employed nurse organised the screening of these patients and within six months only 42 patients (2.1% of the total aged 30-64 years) had not come after three requests. The total cost of this final topping-up operation was £150 in 1977; the money was kindly provided by Glaxo Laboratories (to which we are extremely grateful) as part of a larger trial of the treatment of mild hypertension.²

Other reviews of the literature^{3 4} have also concluded that case finding in general practice is preferable. The most recent⁴ concludes that "where the purpose is to find and treat newly diagnosed hypertensive patients, then screening by general practitioners is likely to be a more cost-effective approach." There are, of course, other major advantages: "Follow-up and treatment are in the same hands, borderline values can be watched for a number of years, and an improved attitude to community welfare is generated amongst doctors and patients."⁵ General-practice-based case finding can also be a continuous process, which is a condition laid down as necessary for screening for disease.⁶

The implications for general practice in the Brighton team's work are considerable; particularly alarming is the commendation of a method of screening which detects hypertension "without undue burden on general practitioner services." The only thing which Dr Trafford and his colleagues have shown is that where general practice is badly organised (one-third of the records unusably inaccurate) the work of general practice will be taken over by energetic hospital teams, which will be prepared to spend far more money than is necessary or sensible in doing what can be almost entirely absorbed into the day-to-day work of a primary care team. It is quite clear that modern general practice can detect and treat almost every single patient with hypertension.⁷ Let us not lose general medicine by default and without good evidence, as we have lost domiciliary obstetrics and may well, if the proposals of the Joint Committee (15 December, p 1563) are accepted, lose much of paediatrics also.

S L BARLEY

Sheffield S10 3ED

¹ Hart JT. *Lancet* 1970;ii:223-6.

² Joesbury HE, Phillips CA, Garrett RT, Wilkes E, Smith AJ. *Br Med J* 1976;ii:1476-9.

³ Adler MW. *J Roy Coll Gen Pract* 1976;26:171-6.

⁴ Christie D. *J Roy Coll Gen Pract* 1979;29:597-601.

⁵ Coope J. *J Roy Coll Gen Pract* 1974;24:161-6.

⁶ Wilson JMG, Jungner G. In: *Principles and practice of screening for disease*. Geneva: WHO, 1968.

⁷ Parkin DM, Kellert RJ, Maclean DW, Ryan MP, Fulton M. *J Roy Coll Gen Pract* 1979;29:590-4.

Peripheral nerve lesions in the muscle compartment syndrome

SIR,—Your leading article (6 October, p 818) describes the compartmental syndrome of local damage to the contents of osteofascial compartments of limbs crushed by pressure sustained in a variety of ways. You rightly emphasise that alcoholics, drug addicts, and victims of accidental poisoning may be injured in this way, the critical factors being assumed to be immobility and coma, which subject the victim to the sustained weight of the head or trunk lying on the arm, or of one leg on the other. It is important to recognise that not only are muscle and blood vessels injured by