attention to sodium. We suggest that the changes in blood pressure that they saw were due to changing the dietary sodium, and that 24-hour urine collections would have revealed this.

Dietary fibre may not be irrelevant. Varying fibre amount and type might have altered the availability of cations for absorption in the bowel. This, however, is speculation and could not be detected by comparing tables of dietary sodium content. In any event, there is no case for relating this study to Burstyn's experiments with fat-enriched diets for rabbits (references 6-9 in the paper) until the major confounding variable of effective cation intake has been taken into account in both species.

The report also omits any mention of a search for an effect of age on blood pressure in the subjects, and does not refer to the age distributions of the various experimental groups. It is clear that blood pressure was measured in many different ways (at home, at work, at a health centre; with a standard and a random zero instrument; thrice weekly and on a single occasion) but we are not told how the various methods were distributed between the groups and during the crossover experiment.

L C WILLIAMS
J H TURNEY
VICTOR PARSONS

King's College Hospital
Renal Unit,
Dulwich Hospital,
London SE22 8PT

¹ Anon. *Lancet* 1975;ii:1325-6.

² Biachi G, et al. *Clin Sci Mol Med* 1974;47:435-48.

³ Dahl LK, et al. *Circ Res* 1974;34:94.

⁴ Parisi J, et al. *Amer Heart J* 1973;85:22-34.

⁵ Morgan T, et al. *Lancet* 1978;ii:227-30.

⁶ Magnani B, et al. *Clin Sci Mol Med* 1976;51:625-6S.

⁷ Freis ED, *Circulation* 1976;53:589-95.

SIR,—Angela Wright and others (15 December, p 1541) recently reported that a high-fibre diet exerted a hypotensive effect on the blood pressure of normal volunteers, the mechanism of which is not known. The diet of primitive societies with a low prevalence of hypertension is not only high in fibre content but low in sodium and rich in potassium. Modest modification of sodium intake may be helpful in the treatment of mild hypertension¹ and potassium supplementation in the diet may exert an ameliorating effect on the development of hypertension and protect from hypertensive complications.² Wholemeal bread has double the potassium content of white bread and All Bran has a much higher potassium content than cornflakes.³ It is possible that alteration of sodium and potassium intake by the volunteers in this study could have caused the blood pressure changes observed. In addition to information concerning the electrolyte content of the diets prescribed, any information on the changes in weight in these volunteers would be of interest, especially as potassium may have a natriuretic effect.⁴

It is surprising to observe such low standard errors in blood pressure measurement as those quoted in this paper when one notes the age range (18-60 years) of the volunteers and the fact that exercise was undertaken between each inflation of the cuff. On a low-fibre control diet the systolic blood pressure of 17 of their volunteers was 121.2 ± 1.6 mm Hg and the diastolic blood pressure was 78.5 ± 1.7 mm Hg. Using a London School of Hygiene "blind" sphygmomanometer we found that the supine systolic blood pressure

of 19 male medical students, aged 20-22 years, was 123.4 ± 2.4 mm Hg and the supine diastolic blood pressure was 66.4 ± 2.3 mm Hg. Bigger standard errors are normally found in groups of subjects with the age range given in the paper.

P PARFREY
KATE CONDON
P WRIGHT

Medical Unit, London Hospital
Medical College,
London E1 1BB

¹ Morgan T, Adam W, Gillies A, Wilson M, Morgan G, Carney S. *Lancet* 1978;ii:227-30.

² Meneely GR, Battarbee JD. *Am J Cardiol* 1976;38:768-85.

³ Paul AA, Southgate DAT. *McCance and Widdowson's the composition of foods*. 4th ed. London: HMSO, 1978.

⁴ Young DB, McCaa RE, Pan YJ. *Circ Res* 1976;38(6 suppl 2):84-9.

*We sent copies of these letters to Dr Burstyn, whose reply is printed below.—ED, *BMJ*.

SIR,—The letter of Dr Williams and his colleagues suggests that the effect we report (15 December, p 1541) may have been due to changes in the sodium intake of our subjects. Although we did not measure the sodium intake of the people participating in our study, we have no reason to suppose that they should synchronously change their food salting habits to confound our experiment. The high-fibre bread we gave our subjects would have supplied them with 31 mmol(mEq) sodium a day, while the low-fibre bread supplied 38 mmol/day (based on their mean intake of about 1 kg of bread a week). This difference of 7 mmol/day represents 8% of the rather low sodium intake recommended by the US Senate Select Committee on Nutrition¹ and less than 5% of the sodium intake typical of Western man, hardly sufficient to account for the effect we report.

Dr Williams and his colleagues are also dissatisfied with our blood pressure measurement techniques. All of the blood pressures apart from those of our 12 hypertensive patients were measured with Hawksley Random Zero sphygmomanometers in an office at the university. The two observers standardised their techniques carefully. Only one of these observers measured the blood pressures of the hypertensive patients at their own homes using an ordinary Accoson sphygmomanometer. We did not have large enough numbers of participants over the age of 30 to look for any effect of age on blood pressure. The average age of our subjects (with the exception of the hypertensive patients) was 26 and did not differ between experimental groups.

The letter of Dr Parfrey and his colleagues quite correctly asks how we managed to keep our standard errors small. The precision of our measurements was achieved by taking blood pressure readings three times weekly, each of which was itself the mean of three sphygmomanometer inflations. Between inflations subjects were asked to clench their fist sufficiently to prevent pooled blood from distorting subsequent diastolic pressure values. Furthermore, a group of 36 people whose systolic pressures were measured three times on a single occasion yielded the following result: first measurement— 128.7 ± 2.3 ; third measurement— 123.5 ± 2.2 . These were compared with the blood pressures of 36 of our experimental volunteers (who were age matched with the above group) towards the

end of their control periods (unfortunately, we had not preserved the three individual blood pressure values of our volunteers from the beginning of their control periods): first measurement— 117.8 ± 1.7 ; third measurement— 117.8 ± 1.6 . The systolic pressures of people who are unaccustomed to blood pressure measurements drops significantly ($p < 0.001$) in the four minutes between the first and third measurement. This is not the case with the "trained" volunteers who have had two weeks of blood pressure measurements. The systolic pressures of the "untrained" people were higher than those of the "trained" people (first measurement— $p < 0.001$; third measurement— $p < 0.05$), although the diastolic pressures of the two groups were very similar. Finally, the standard errors for the blood pressure readings are considerably larger for the "untrained" group (which are similar to those of Dr Parfrey and others) than the "trained" group. It appears that repeated blood pressure measurements over a period of days improve precision in two ways: (a) by allowing the observer to take mean values and (b) by accustoming the individual to the measurement and hence reducing the variability of this measurement (that is, producing lower systolic pressures in the third reading of "untrained" people and lower mean systolic pressures in "trained" people).

The suggestion by Angela Wright and her colleagues that potassium may have played a part in producing our results is an interesting one. Our bran-enriched wholemeal bread contained more potassium than our white bread and would have provided (with the 5 g bran a day given to the volunteers) 18 mmol(mEq)/day potassium, compared with 5 mmol/day provided by the low-fibre bread. This difference of 13 mmol/day amounts to 20% of the average daily potassium intake.² That this may have affected our results is undeniable, although it required 135 mmol/day potassium to reduce the blood pressure in the two cases of salt-induced hypertension cited by Meneely and Battarbee.³

PETER BURSTYN

University Department of Physiology
and Pharmacology,
Southampton SO9 3TU

¹ Select Committee on Nutrition and Human Needs. *Dietary goals for the United States*. Washington: United States Senate, 1977.

² Davidson S, Passmore R, Brock JF, Truswell AS. *Human nutrition and dietetics*. 6th edn. Edinburgh: Churchill Livingstone, 1975.

³ Meneely GR, Battarbee JD. *Am J Cardiol* 1976; 38:768-85.

Sclerosant treatment for hydroceles and epididymal cysts

SIR,—My experience of sclerosant treatment in the management of hydroceles and epididymal cysts differs widely from that reported by Mr H Thomson and Mr M Odell (22 September, p 704).

As a result of Moloney's¹ enthusiastic article¹ I set up a prospective study and treated approximately 75 patients between 1 January 1976 and 31 December 1978. The results in the first 56 patients were reported in detail earlier this year.² In summary, assessment one to two years after treatment showed 95% cure for hydroceles and 100% for epididymal cysts. Side effects and complications were minimal, none requiring surgical intervention. The few complications