trivial, I would like to point out that dracunculiasis is not acquired either by ingesting or by contact with snails but by drinking water containing infected *Cyclops* species, which are small (1-2 mm) freshwater crustaceans. In man the mature female worm discharges first-stage larvae (not eggs) at the surface of the skin, usually when the affected part of the body (commonly the foot and lower leg) comes into contact with water. These first-stage larvae are extremely active and can live for four to seven days; their further development, however, depends on their being eaten by a predatory species of *Cyclops*. The larvae penetrate the gut of the *Cyclops* and after two moults in the haemocoel reach the infective third stage.¹ Dracunculiasis today is mainly a disease of rural communities.

D ROLLINSON

Department of Zoology,
British Museum (Natural History),
London SW7 5BD

¹ Muller R. *A manual of medical helminthology*. London, Heinemann, 1975.

St Joseph's Hospice

SIR,—Unbalanced though it was, the programme on St Joseph's Hospice did not justify the strictures of your correspondent AS (15 December, p 1579).

Factual errors show that AS is out of touch with the running of St Joseph's and therefore with its singular lack of religiosity. Ministers, parsons, priests, and rabbis are familiar and welcome visitors, but proselytisation is unknown. If conversions occur, they do so in accordance with Johnson's dictum on imminent death. Certainly, there is nothing to suggest that Catholic or Christian patients are favoured more than those of other (or no) religion.

It would be very sad if AS succeeded in dissuading doctors, nurses, or patients from accepting what St Joseph's offers. I know because I have worked there; I am an Anglican with sceptical tendencies.

B SYMONDS

Stone, Staffs ST15 8LP

Resuscitation of the newborn

SIR,—Few articles and even fewer chapters in books of neonatal medicine give any specific details of the techniques that are necessary to resuscitate newborn babies effectively and safely. Dr H B Valman's article (24 November, p 1343) is exceptional in giving an outline of this subject with which all midwives and paediatricians should be familiar. However, there is one aspect of his instructions on endotracheal intubation which I think needs immediate correction.

The epiglottis is a sensitive and fragile tissue and the blade of the laryngoscope should not be placed over it or be used to press it against the tongue. Instead the tip of the laryngoscope should be advanced slowly over the tongue from the right side of the mouth and positioned in the vallecula, anterior to the epiglottis. In this position the epiglottis swings anteriorly when the laryngoscope is gently raised. There can then be no doubt about the site of the entrance to the larynx even when it is obscured by secretions or meconium. Failure to intubate is frequently due to unnecessary urgency, which leads to an incomplete identification of the anatomy of

the upper airway. This is made more difficult when the laryngoscope is advanced too far. Soft-tissue injury of the upper airway, including trauma to the epiglottis, is a common finding after frequent attempts at intubation by inexperienced personnel. Mechanical stimulation of the epiglottis can also cause reflex laryngeal spasm, making intubation even more difficult.¹

Another common mistake which adds to the difficulties encountered during intubation is the idea that the baby should be placed on a sloping surface with the head much lower than the feet. Dr Valman quite rightly states that the infant should be placed supine on a flat (horizontal) surface. In this position the head should be slightly elevated *without extending* the neck.² Unfortunately, the picture in Dr Valman's article shows a baby on a tilted surface on which some babies will slide, making intubation even more difficult.

ANDREW WILKINSON

Neonatal Unit,
Department of Paediatrics,
John Radcliffe Hospital,
Oxford OX3 9DU

¹ Johnson P. In: RR Robinson, ed. *Proceedings of the Francis E Camps International Symposium on Sudden and Unexpected Deaths in Infancy*. Toronto: Canadian Foundation for the Study of Infant Deaths, 1974:231-241.

² Klaus NH, Fanaroff AA. *Care of the high-risk neonate*. Philadelphia: WB Saunders Company, 1979:31-3.

Breathing difficulties in the newborn

SIR,—In his article in the *BMJ* of 8 December (p 1483), Dr H B Valman states that the cause of the respiratory distress syndrome is a deficiency of pulmonary surfactant.

While I would not dispute that a deficiency of surfactant *correlates* with the respiratory distress syndrome, precipitating *causes* are observable daily during hospital management of preterm infants. Every intervention, such as heel-pricking with needles, weighing on metal scales, insertion of nasogastric tubes, and especially insertion of rectal thermometers, causes an observable disturbance of respiration, and is frequently followed by a sustained increase in respiratory rate.

The distress which these regular and painful procedures cause the infants can only be expressed physically. This distress may be lessened by nurses who are aware of the babies' feelings, who talk and handle them sympathetically, and who stay a while with the babies when the procedures are over, talking and touching in a non-painful way and giving the babies something to suck—that is, holding the whole baby in mind.

If one's only contacts with other humans are painful, one's hope and one's will to survive are bound to diminish. No one who takes time and trouble to observe these babies will fail to see what I have described above.

AVERIL EARNSHAW

Adolescent Department,
Tavistock Clinic,
London NW3 5BA

Multiple sclerosis: what can and cannot be done

SIR,—The contribution of Professor E J Field and Greta Joyce (15 December, p 1571) is rightly published under the heading "For debate" and we hope that a lively discussion

will follow. Otherwise readers could get the wrong impression that multiple sclerosis has become a disease easy to diagnose and to prevent. Except for point (1)—that there is no cure for clinically established MS—all statements made by the authors are unproved or misleading.

Work with coded material has shown that the erythrocyte unsaturated fatty acid (E-UFA) test is non-specific¹ and by no means better than other laboratory parameters (immunoglobulin ratio, oligoclonal bands of the cerebrospinal fluid, etc). The results of other authors who could not confirm the test^{2,3} cannot be ignored. Seaman *et al*⁴—given as reference for the test's validity—could reproduce the results for clinically confirmed cases, but not for patients "with minimal neurological findings." That means that even if the test turns out to be reproducible in other laboratories we are still left with the difficulty of early diagnosis and often have to wait for dissemination in space and (indispensable not "wasted") time.

From several well-controlled studies we know that high doses of polyunsaturated fatty acids may mitigate or shorten a bout,^{5,6} but do not influence the overall prognosis.^{7,8} To speak of an "inborn anomaly in the handling of unsaturated fatty acids" as if this were a proved fact rather than a speculative hypothesis is not only a doubtful scientific manoeuvre: it creates a lot of anxiety in patients and their relatives. The pattern of familial multiple sclerosis is hardly compatible with Professor Field's assumption, apart from the fact that the process of myelin building is very little influenced by nutritional factors.

The E-UFA test is under investigation world wide because of its interesting pathogenetic mechanisms, which are far from being understood. The practical implications it might have are still doubtful.

S POSER

WOLFGANG POSER

University Department of Neurology,
34 Göttingen,
Federal Republic of Germany

¹ Field EJ, Joyce G, Bauer HJ, Wikström J. *Lancet* 1977;ii:658.

² Forrester JA, Smith WJ. *Lancet* 1977;ii:453-4.

³ Hawkins SA, Millar JHD. *Lancet* 1979;ii:165-6.

⁴ Seaman GVF, Swank RL, Tambllyn CH, Zukoski CF, IV. *Lancet* 1979;ii:1138-9.

⁵ Millar JHD, Langman MJS, Payling Wright H, *et al*. *Br Med J* 1973;ii:765-8.

⁶ Bates D, Fawcett PRW, Shaw DA, Weightman D. *Br Med J* 1978;ii:1390-1.

⁷ Bates D, Fawcett PRW, Shaw DA, Weightman D. *Br Med J* 1977;ii:932-3.

⁸ Paty DW, Cousin HK, Read S, Adlaka K. *Acta Neurol Scand* 1978;58:53-8.

SIR,—Professor E J Field and Greta Joyce (15 December, p 1571) quoted Seaman and his colleagues,¹ who, using a slightly modified Field's E-UFA (erythrocyte unsaturated fatty acid) test, were correct in all 65 patients with established multiple sclerosis. However, Seaman *et al* found the test negative in all four early cases, which would not support Field's claim of being able to identify latent MS and children at risk. A positive test would seem to be due to a secondary phenomenon rather than an inborn anomaly in the handling of unsaturated fatty acids.

Two trials^{2,3} have shown a marginal benefit for MS patients taking at least 17 g of linoleic acid daily. Eight capsules of Naudicelle daily were of no benefit whatever, possibly owing to the low dose of linoleic acid (3 g daily) and γ -linolenic acid (340 mg daily).³ The theoretical claim that γ -linolenic acid would be more

successful than linoleic acid has yet to be proved.

J H D MILLAR

Department of Neurology,
Royal Victoria Hospital,
Belfast BT12 6BA

¹ Seaman GVF, Swank RL, Tambllyn CH, Zukoski CF IV. *Lancet* 1979;ii:1138-9.

² Millar JHD, Zilka KJ, Langman MJS, et al. *Br Med J* 1973;ii:765-8.

³ Bates D, Fawcett PRW, Shaw DA, Weighman D. *Br Med J* 1978;ii:1390-1.

Retrospective versus prospective controls in clinical trials

SIR,—It has happened again. Many a time now, in my relatively short medical career, I have accepted as gospel truth medical knowledge without personally examining its foundations, only to find some time later that those foundations are almost non-existing. Examples include tonsillectomy, anticoagulant therapy for infarction, and cleansing the skin with alcohol before venepuncture. The most recent example was the attack on the holiness of prospective trials launched by Dr Lawrence Cranberg (17 November, p 1265). His article fills one's heart with fear, for no longer will one be able to stand up at Grand Rounds and put down some bright young whippersnapper: "Yes, but were those findings confirmed in a prospective trial?"

In spite of such disadvantages, Dr Cranberg's article has one great advantage: it leads us to examine the postulates on which clinical trials are based. In physics and chemistry there is no need for concurrent controls, for the variables are few and well known. Experiments are designed in serial fashion, altering one variable at a time. A careful record of the critical variables is kept and the experiment can be repeated time and time again, with identical results. In biology, however, we not only have a greatly increased number of variables, we also understand them less well. Indeed, sometimes we are not aware of variables that may effect the outcome of the experiment. This problem has been circumvented by the process of matching experimental preparations and then introducing a known controllable variable (administration of a chemical, the making of a lesion, etc) to one of the matching pair only, usually in a randomised fashion. The effect of the introduced variable is then determined by observing any difference between the pairs that did not exist initially. It is necessary only to match for those variables that might be affected by the experimental process. Thus sex, weight, age, concentration of certain metabolites, and the presence and severity of certain characteristics such as symptoms and signs of disease are among the more important variables. Duration of a disease may be important, but the point in time at which a variable is tested rarely is (provided ambient temperatures, etc, are treated as different variables). Yet the only advantage of the use of concurrent controls is to be able to say that the two experiments were done at the same point in time in the life of the universe. Other apparent advantages, such as ease of comparison for sameness, are important only in experiments where the variables are improperly recorded. Generally, the result of such work is as sloppy as the record keeping. Finally, it has been suggested by some reviewers of grant applications—not, of course, any associated with agencies to

which I apply for funds—that a temptation exists in retrospective trials to select the controls and experimental subjects on the basis of the expected or desired outcome. To those individuals I would say: "The null hypothesis states that there is no difference in the inherent honesty of researchers using retrospective and concurrent controls." Until the latter is disproved in a prospective, double-blind crossover trial, I shall feel justified in using retrospective or concurrent controls as dictated by the variables of the experimental situation.

In conclusion, thank you, Dr Cranberg.

GERALD TEVAARWERK

Department of Medicine,
University of Western Ontario,
St Joseph's Hospital,
London, Ontario,
Canada N6A 4V2

Rehabilitation

SIR,—May I take up some points from Dr D H Bossingham's letter (8 December, p 1508)?

It is certainly the responsibility of doctors, at both primary and secondary care level, to think about any necessary stages in rehabilitation that their patients might need during recovery, and to direct their patients in the desired direction and to the appropriate agencies for any help needed after (or during) clinical treatment. Unfortunately, rehabilitation is considered to be a "soft" subject and, perhaps partly for this reason, often induces an apparent occupational deafness in harassed clinicians.

If the subject is to become more scientific, thought and attention must be given to developing valid measures of outcome and evaluating different rehabilitation procedures. The recently founded Society of Research in Rehabilitation provides a forum for many disciplines (not only medicine but others including the remedial professions) to look critically at work that is being done not only in medical rehabilitation but in other fields, such as those of occupational assessment and employment rehabilitation.

Dr Bossingham points out that many doctors do not make use of the Manpower Service Commission's employment rehabilitation centres to help their patients return to work, if such an assessment is needed. While the employment rehabilitation centre does not provide training, a skilled occupational assessment can be given by a team which includes not only medical and nursing advice but also that of occupational psychologists and skilled workshop supervisors. Such an assessment may be followed by a placement directly into suitable work or possibly by training.

The number of referrals of patients to these centres from doctors has steadily declined over the last few years. There are many possible reasons for this, and to try to counteract this trend the Employment Medical Advisory Service (EMAS), which now provides medical and nursing services for 25 of the 27 employment rehabilitation centres, has recently agreed with the Employment Service Division (which manages the centres) the introduction of an additional method of direct medical referral of patients to the employment medical adviser at the centre. This referral could come from doctors working at either primary or secondary health care level.

I would like to draw your readers' attention to the further information about this new facility on p 193.

FELICITY C EDWARDS

Employment Medical Advisory Service,
Health and Safety Executive,
London NW1 5DT

Drug names that look or sound alike

SIR,—In the issue of 17 November (p 1297), Drs R N Palmer and F V Simpson cite instances of similar sounding drugs causing confusion. It is thus ironic that on the following page (1298) "Minerva" is confused by one of the very drug pairs listed by Dr H McNulty and Mr P Spurr (6 October, p 836). In the letter she quotes¹ Montello and her colleagues at the Children's Hospital of Philadelphia used clotrimazole and not co-trimoxazole to treat their patient's oral thrush. Dr W T Houlshy (p 1297) should note that these *are* the approved names—surely an unnecessary confusion but one which may lessen if trimethoprim alone replaces co-trimoxazole in common use.

An oral preparation of clotrimazole appears not to be available in this country^{2,3} but there is a suitable formulation of miconazole. Dakatrin Oral Gel (Janssen Pharmaceutical Ltd) is designed to persist in the mouth and might thus be a candidate for prescription "by thumb."

NEIL C FRASER

Royal Hospital for Sick Children,
Edinburgh EH9 1LF

¹ Montello JM, Darby MH, Faubel K, August CS. *New Engl J Med* 1979;301:1005.

² *Data Sheet Compendium 1979-80*. London: Pharmind Publications Ltd.

³ *MIMS* 1979;21, No 10. London: Haymarket Publishing Ltd.

Iatrogenic collapse

SIR,—I would like through your columns to discuss points brought up by Dr M J Boscoe (24 November, p 1367). A complication of profound respiratory depression is serious indeed, but I felt somewhat reassured after reading the references quoted in his letter.

In the fifth edition of *The Pharmaceutical Basis of Therapeutics*¹ respiratory depression is mentioned as a complication of Pamergan (pethidine hydrochloride 100 mg, promethazine hydrochloride 25 mg, and atropine sulphate 0.6 mg), but no statistics are given. In a large series² respiratory depression of short duration occurred in two patients out of 774 receiving the drug parenterally. This covers 10 320 doses of the drug, an incidence of 0.02%. One of these two patients was suffering from advanced carcinoma of the oesophagus and the second was markedly debilitated with generalised arteriosclerosis.

Comparing seven analgesic drugs, Prescott *et al*³ showed that all of them had caused some depression of respiration, a fact that is of great importance in obstetric practice owing to possible depressant effects on the respiration of the newborn. He quotes that, of the more established drugs, pethidine is the most useful obstetric analgesic. Of the eight medical students he investigated, respiratory depression was shown by asking volunteers to breathe a gas mixture containing 0.5% carbon dioxide in oxygen, the analgesics being injected intramuscularly after an initial record had been taken. No subject suffered profound respiratory depression.